

ANNALES DE L'I. H. P., SECTION A

J. MAGNEN

R. SENEOR

A note on cluster expansions

Annales de l'I. H. P., section A, tome 38, n° 1 (1983), p. 93-98.

http://www.numdam.org/item?id=AIHPA_1983__38_1_93_0

© Gauthier-Villars, 1983, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A », implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A note on cluster expansions

by

J. MAGNEN and R. SENEOR

Centre de Physique Théorique de l'École Polytechnique,
Plateau de Palaiseau, 91128 Palaiseau, Cedex, France
« Groupe de Recherche du C. N. R. S. n° 48 »

ABSTRACT. — We make, on a particular field theory model, an estimate of the weakest long range behaviour necessary to obtain easily a convergent cluster expansion.

RÉSUMÉ. — On estime, dans un modèle particulier de Théorie des Champs, le comportement à grande distance le plus faible permettant d'obtenir facilement un développement en « clusters » convergent.

1. INTRODUCTION

We prove the existence of the infinite volume limit, with a cluster expansion, of a φ_2^4 model with a covariance

$$C(x, y) = \int d^2p \exp i(x - y) \frac{|p|^{1,5+\varepsilon}}{(p^2 + 1)^2} \quad \text{for} \quad 0 < \varepsilon < 1/2$$

so that

$$|C(x, y)| \leq \frac{O(1)}{|x - y|^{3,5+\varepsilon/2} + 1}$$

Such models have been considered by Federbush [1].

In field theory the cluster expansion is a creation of Glimm Jaffe Spencer; the first version [3] used only characteristic functions, and Brydges gave of it a convenient form that we use [0]. Here the two estimates (lemma 1 and 2) are taken from Glimm and Jaffe [4], [2].

2. THE EXPANSION

For simplicity we consider a two point function:

$$\frac{S_\Lambda(f, g)}{Z_\Lambda} = Z_\Lambda^{-1} \int \varphi(f)\varphi(g) \exp\left(-\lambda \int_\Lambda \varphi^4(x) dx\right) d\mu$$

where

$$Z_\Lambda = \int \exp\left(-\lambda \int_\Lambda \varphi^4(x) dx\right) d\mu \quad \text{and} \quad \varphi(f) = \int \varphi(x) f(x) dx$$

where Λ is a space cutoff and $d\mu$ is the Gaussian measure of covariance $C(x, y)$ and mean zero.

We consider a unit lattice \mathcal{D} on \mathbb{R}^2 . The support of f (resp. g) is in $\Delta f \in \mathcal{D}$ [resp. Δg].

To a square $\Delta_1 \in \mathcal{D}^*$ we associate a variable s_1 and define

$$C(s_1) = s_1 C + (1 - s_1) C_1, \quad C_1 = \chi_{\Delta_1} C \chi_{\Delta_1} + (1 - \chi_{\Delta_1}) C (1 - \chi_{\Delta_1})$$

where χ_{Δ_1} is the characteristic function of Δ_1 .

Let $d\mu(s_1)$ be the measure of covariance $C(s_1)$ so that for some function Q of the field:

$$S(s_1) = \int Q d\mu(s_1)$$

A step of the expansion consists in the decomposition

$$S = S(s_1)|_{s_1=1} = S(0) + \int_0^1 \left(\frac{d}{ds_1} S(s_1)\right) ds_1$$

$$\frac{d}{ds_1} \int Q d\mu(s_1) = \sum_{\Delta_2} \int P(\Delta_1, \Delta_2) Q d\mu(s_1)$$

with (see [4]).

$$P(\Delta_1, \Delta_2) = \int \frac{dC(s_1)}{ds_1}(x, y) \frac{\delta}{\delta\varphi(x)} \frac{\delta}{\delta\varphi(y)} \chi_{\Delta_1}(x) \chi_{\Delta_2}(y) dx dy$$

In our case $Q = \varphi(f)\varphi(g)e^{-\int\varphi^4}$ so that, with $Q|_R$ = the part of Q with support in R , $C_\Delta = \chi_\Delta C \chi_\Delta$ and $S_{\sim\Delta} = \int Q|_{\sim\Delta} d\mu|_{\sim\Delta}$:

$$S = \int Q|_{\Delta_1} d\mu|_{\Delta_1} \cdot S_{\sim\Delta} + \sum_{\Delta_2} \int_0^1 ds_1 \int P(\Delta_1, \Delta_2) Q d\mu(s_1)$$

(*) We can take $\Delta_1 = \Delta f$ or Δg , but this is not necessary.

Then we define

$$C(s_1, s_2) = s_2 C(s_1) + (1 - s_2) C(s_1)_2$$

$$C(s_1)_2 = \chi_{\Delta_1 \cup \Delta_2} C(s_1) \chi_{\Delta_1 \cup \Delta_2} + (1 - \chi_{\Delta_1 \cup \Delta_2}) C(s_1) (1 - \chi_{\Delta_1 \cup \Delta_2})$$

and

$$\int P(\Delta_1, \Delta_2) Q d\mu(s_1) = \int P(\Delta_1, \Delta_2) Q \Big|_{\Delta_1 \cup \Delta_2} d\mu(s_1, s_2) \Big|_{\substack{s_2=0 \\ \Delta_1 \cup \Delta_2}} \cdot S_{\sim \Delta_1 \cup \Delta_2}$$

$$+ \sum_{j=1}^2 \sum_{\Delta_3} \int_0^1 ds_2 P(\Delta_j, \Delta_3) P(\Delta_1, \Delta_2) Q d\mu(s_1, s_2)$$

and so forth; we obtain

$$S = \int \varphi(f) e^{-\int_{\Delta_1} \varphi^4} d\mu \Big|_{\Delta_1} \cdot S_{\sim \Delta_1}$$

$$+ \sum_{\Delta_2} \int ds_1 \int P(\Delta_1, \Delta_2) Q \Big|_{\Delta_1 \cup \Delta_2} d\mu(s_1, s_2) \Big|_{\substack{s_2=0 \\ \Delta_1 \cup \Delta_2}} \times S_{\sim \Delta_1 \cup \Delta_2}$$

$$+ \sum_{i=3}^{\infty} \sum_{j_1=1}^1 \sum_{j_2=1}^2 \dots \sum_{j_{i-1}=1}^{i-1} \sum_{\substack{\{\Delta_j\} \\ i \geq j \geq 2}} \int P(\Delta_{j_{i-1}}, \Delta_i) P(\Delta_{j_{i-2}}, \Delta_{i-1})$$

$$\dots P(\Delta_{j_1}, \Delta_2) Q \Big|_{\cup \Delta_i} ds_1, \dots, ds_{i-1} d\mu_{(s_1, \dots, s_{i-1}, s_i)} \Big|_{\substack{s_i=0 \\ \cup \Delta_i}} \cdot S_{\sim \cup \Delta_i} \quad (1)$$

where $\cup \Delta_i$ means restricted to $\Delta_1 \cup \dots \cup \Delta_i$.

The formula giving $C(s_1, \dots, s_i)$ is the generalisation of the one giving $C(s_1, s_2)$:

$$C(s_1, \dots, s_i) = s_i C(s_1, \dots, s_{i-1}) + (1 - s_i) C(s_1, \dots, s_{i-1})_i$$

$$C(s_1, \dots, s_{i-1})_i = \chi_{\cup \Delta_i} C(s_1, \dots, s_{i-1}) \chi_{\cup \Delta_i} + (1 - \chi_{\cup \Delta_i}) C(s_1, \dots, s_{i-1}) (1 - \chi_{\cup \Delta_i})$$

with $\chi_{\cup \Delta_i} = \chi_{\Delta_1 \cup \dots \cup \Delta_i}$.

Then as noted above $C(s_1, \dots, s_i) |_{s_i=0}$ is a direct sum, so that $d\mu |_{s_i=0}$ factorizes and then in each term of formula (1) the Schwinger function in $\Delta_1 \cup \dots \cup \Delta_i$ is factorized out. We call such a term a tree, because the

propagators $\frac{dC}{ds}$ form no cycle (by construction):

$$\frac{dC}{ds_k}(s_1, \dots, s_k)(x, y) \quad \text{is non zero only if} \quad \begin{array}{l} x \in \Delta_1 \cup \dots \cup \Delta_k \\ y \notin \Delta_1 \cup \dots \cup \Delta_k \end{array}$$

so that at each step a propagator connects the tree formed by the squares of the tree (in formation) with some new square.

If we choose $s_i = 0$ then the tree is formed by $\Delta_1, \dots, \Delta_i$ and is factorized out, as said.

Then in each term of formula (1) we expand in the same way $S_{\sim \cup \Delta_i}$ choosing a Δ'_1 in $\mathbb{R}^2 \setminus \cup \Delta_i$ which plays the role of Δ_1, \dots . For given $\{j_1, \dots, j_{i-1}\}$ in formula (1), a vertex Δ_k is said of order n_k or

$$n(\Delta_k) = \# \{j_\alpha = k, \alpha \geq k\},$$

i. e. there are $n_k + 1$ propagators which have an extremity in Δ_k , except for Δ_1 where there are only n_1 .

Finally we obtain

$$S = \Sigma \Pi \text{ trees}$$

One tree contains Δ_f and Δ_g by parity.

For $\Delta_1, \dots, \Delta_i$ given we call $T(\Delta_1, \dots, \Delta_i)$ the sum of the trees whose vertices are $\Delta_1, \dots, \Delta_i$.

In the following we prove

PROPOSITION. —

$$\left| \sum_{\substack{\{\Delta_k\} \\ k \neq i, j}} T(\Delta_1, \dots, \Delta_n) \right| \leq 0(1) \frac{1}{(\text{dist}(\Delta_i, \Delta_j) + 1)^{3.5 + \varepsilon/4}} e^{-nK}$$

where K is as big as we want for λ small enough.

As a conclusion each tree is exponentially small like the number of squares that it contains. Moreover we can sum over the localization of the squares of each tree.

A standard argument of statistical mechanics [4, Chapter 6] gives then

Final result. — For λ small enough $\lim_{\Lambda \rightarrow \infty} \frac{S_\Lambda(f, g)}{Z_\Lambda}$ converges and is bounded by

$$\text{Cst}(f) \text{Cst}(g) (\text{dist}(\Delta_f, \Delta_g) + 1)^{-(3.5 + \varepsilon/4)}$$

3. THE BOUNDS

The form of each tree is given by the $n(\Delta)$'s. The sum over each $n(\Delta)$ is controlled using $\sum_{n(\Delta)} 2^{-n(\Delta)} \leq 0(1)$ so that $\sum_{n(\Delta)} \dots \leq 0(1) \sup_{n(\Delta)} 2^{n(\Delta)}$.

Now the sum over the order of the Δ 's in the tree is taken in into account by summing over all the localizations of the vertices of the tree using $[d(\Delta_1, \Delta) + 1]^{-2 - \varepsilon/4}$ as a combinatorial factor. We use

$$\sum_{\Delta} (d(\Delta_1, \Delta) + 1)^{-2 - \varepsilon/4} \leq 0(1)$$

for the $n(\Delta_1)$ squares linked to Δ ; then we iterate the process.

A chain of propagators links Δf to Δg in the tree containing both. Then using:

$$\sum_{\Delta_2} \frac{1}{(\text{dist}(\Delta_1, \Delta_2) + 1)^{3.5 + \varepsilon/4}} \frac{1}{(\text{dist}(\Delta_2, \Delta_3) + 1)^{3.5 + \varepsilon/4}} \leq \frac{0(1)}{(\text{dist}(\Delta_1, \Delta_3) + 1)^{3.5 + \varepsilon/4}}$$

we obtain the decrease in the distance of Δf to Δg .

Now we prove the proposition on trees.

For each vertex (localized in a square Δ) we apply the bound:

$$\int_{\Delta} \prod_{i=1}^{\alpha} |C(x_i, y) \varphi^{4-\alpha}(y)| dy \leq \left(\prod_{i=1}^{\alpha} \sup_{y \in \Delta} |C(x_i, y)| \right) \int_{\Delta} |\varphi^{4-\alpha}(y)| dy$$

So that the contribution of a square Δ is [for $\Delta = \Delta_1$ the product is up to $n(\Delta_1)$]:

$$I(n(\Delta)) = \left| \left(\prod_{i=1}^{n(\Delta)+1} \int_{\Delta} \frac{\delta}{\delta |\varphi|(x)} dx \right) e^{-\lambda \int |\varphi|^4} \right|$$

LEMMA 1. —

$$I(n(\Delta)) \leq 0(1) \lambda^{\frac{n(\Delta)}{4}} n(\Delta)^{\frac{3}{4}n(\Delta)} 0(1)^{n(\Delta)}$$

Proof. — The $\frac{\delta}{\delta |\varphi|}$'s derive either the exponential or already produced vertices. We use the Holder inequality:

$$\int_{\Delta} (\lambda^{1/4} |\varphi(x)|)^{\alpha} dx \leq \left(\lambda \int_{\Delta} \varphi(x)^4 \right)^{\alpha/4} \quad \alpha = 1, 2, 3,$$

then just by bookkeeping one obtains:

$$I(n) \leq 0(1) n^{\frac{n}{4}} \sup_{n'} \left(\left[\lambda \int_{\Delta} |\varphi(x)|^4 \right]^{\frac{4n'-n}{4}} e^{-\lambda \int_{\Delta} \varphi^4} n^{(n-n')} \right)$$

where n' is the number of vertices created by the $\frac{\delta}{\delta \varphi}$'s and $n^{(n-n')}$ bounds the number of terms produced by the derivations not acting on the exponential.

Using $x^l e^{-x} \leq l^l$ ($x > 0$), one obtains the lemma.

— We have still a non used decrease of the propagators, and if there

is $n(\Delta)$ distinct cubes then there is at least $\left(1 - \frac{\varepsilon}{100}\right)n(\Delta)$ cubes which are such that their distance to Δ is bigger than $0(1)\varepsilon^{1/2}n(\Delta)^{1/2}$ so that:

LEMMA 2. — Let $\Delta_1, \dots, \Delta_{n(\Delta)}$ be the squares linked to Δ and $x_i \in \Delta_i$:

$$\prod_{i=1}^{n(\Delta)} (C(x_i, y) (\text{dist}(\Delta, \Delta_i) + 1)^{2+\varepsilon/4}) \leq \prod_{i=1}^{n(\Delta)} (\text{dist}(\Delta, \Delta_i) + 1)^{-3/2-3\varepsilon/4} \leq 0(1)\varepsilon^{-n(\Delta)}n(\Delta)^{-3/4n(\Delta)}$$

The two lemmas prove then the bound of the proposition.

Remark. — Such an analysis can be made on a lot of models using the positivity of the potential or its equivalent.

REFERENCES

- [0] D. BRYDGES, *Comm. Math. Phys.*, t. **58**, 1978, p. 313. See also.
D. BRYDGES, P. FEDERBUSH, *J. Math. Phys.*, t. **19**, 1978, p. 2064.
[1] P. FEDERBUSH, *Commun. Math. Phys.*, t. **81**, 1981, p. 327.
[2] J. GLIMM, A. JAFFE, *Fortschr. Physik*, t. **21**, 1973, p. 327.
[3] J. GLIMM, A. JAFFE, T. SPENCER, *Ann. Math.*, t. **100**, 1974, p. 585.
[4] J. GLIMM, A. JAFFE, T. SPENCER, *Lecture Notes in Physics*, t. **25**; G. VELO, A. WIGHTMAN, Springer, 1973.

(Manuscrit reçu le 25 mai 1982)

« Toute représentation ou reproduction, intégrale ou partielle, faite sans le consentement de l'auteur ou de ses ayants droit ou ayants cause, est illicite (loi du 11 mars 1957, alinéa 1^{er} de l'article 40). Cette représentation ou reproduction, par quelque procédé que ce soit, constituerait une contrefaçon sanctionnée par les articles 425 et suivants du Code pénal. La loi du 11 mars 1957 n'autorise aux termes des alinéas 2 et 3 de l'article 41, que les copies ou reproductions strictement réservées à l'usage privé du copiste et non destinées à une utilisation collective d'une part et d'autre part que les analyses et les courtes citations dans un but d'exemple et d'illustration. »