

LEO SARIO

Completeness and existence of bounded biharmonic functions on a riemannian manifold

Annales de l'institut Fourier, tome 24, n° 1 (1974), p. 311-317

http://www.numdam.org/item?id=AIF_1974__24_1_311_0

© Annales de l'institut Fourier, 1974, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

COMPLETENESS AND EXISTENCE OF BOUNDED BIHARMONIC FUNCTIONS ON A RIEMANNIAN MANIFOLD

by Leo SARIO ⁽¹⁾

A.S. Galbraith has communicated to us the following intriguing problem : Does the completeness of a manifold imply, or is it implied by, the emptiness of the class H^2B of bounded nonharmonic biharmonic functions ? Among all manifolds considered thus far in biharmonic classification theory (cf. Bibliography), those that are complete fail to carry H^2B -functions, and one might suspect that this is always the case. We shall show, however, that there do exist complete manifolds of any dimension that carry H^2B -functions. Moreover, there exist both complete and incomplete manifolds not permitting these functions, and, trivially, incomplete manifolds possessing them.

We attach a Bibliography of recent work in the field.

1. Let C be the totality of complete Riemannian manifolds M , characterized by an infinite distance of any point of M to the ideal boundary. Denote by $\mathcal{O}_{H^2B}^N$ and $\tilde{\mathcal{O}}_{H^2B}^N$ the classes of N -manifolds, $N \geq 2$, for which $H^2B = \emptyset$ or $H^2B \neq \emptyset$, respectively.

THEOREM 1. — $C \cap \tilde{\mathcal{O}}_{H^2B}^N \neq \emptyset$ for every N .

Proof. — Take the N -cylinder

$$|x| < \infty, \quad |y_i| \leq 1, \quad i = 1, 2, \dots, N-1,$$

with each face $y_i = 1$ identified with $y_i = -1$, so as to obtain a covering space of the N -torus in the same manner as a conventional cylinder is a covering surface of the torus. Let T be this N -cylinder with the Riemannian metric

⁽¹⁾ The work was sponsored by the U.S. Army Research Office-Durham, Grant DA-ARO-D-31-124-71-G181, University of California, Los Angeles.

$$ds^2 = \mu^{-2}(x) dx^2 + \mu^{4/(N-1)}(x) \sum_{i=1}^{N-1} dy_i^2$$

where

$$\mu(x) = (2 + x^2)^{\frac{1}{2}} \log(2 + x^2).$$

To see that $T \in C$, it suffices to show, in view of the symmetry, that $\int_0^\infty \mu^{-1}(x) dx = \infty$. The verification is immediate :

$$\begin{aligned} \int_0^\infty (2 + x^2)^{-\frac{1}{2}} \log^{-1}(2 + x^2) dx &> \frac{1}{2} \int_0^\infty (2 + x)^{-1} \log^{-1}(2 + x) dx \\ &= \frac{1}{2} \int_0^\infty \log \log(2 + x) = \infty. \end{aligned}$$

We introduce the function

$$u(x) = \int_0^x \mu^{-3}(t) \int_0^t \mu(s) \int_0^s \mu^{-3}(r) dr ds dt.$$

The Laplace-Beltrami operator $\Delta = d\delta + \delta d$ gives

$$\Delta u = -g^{-\frac{1}{2}} (g^{\frac{1}{2}} g^{xx} u')' = -\mu^{-1} (\mu \mu^2 u')' = -\int_0^x \mu^{-3}(r) dr$$

and

$$\Delta^2 u = -\mu^{-1} (\mu \mu^2 (-\mu^{-3}))' = 0.$$

Thus u is nonharmonic biharmonic.

To see that u is bounded it suffices to show that it is so for $x > 0$. For all $s > 0$,

$$\int_0^s \mu^{-3}(r) dr = \int_0^s (2 + r^2)^{-3/2} \log^{-3}(2 + r^2) dr = o(1),$$

and for all $t > 0$,

$$\begin{aligned} \int_0^t \mu(s) \int_0^s \mu^{-3}(r) dr ds &< c \int_0^t (2 + s^2)^{\frac{1}{2}} \log(2 + s^2) ds \\ &< 2c \int_0^t (2 + s) \log(2 + s) ds \\ &= c \left[(2 + t)^2 \log(2 + t) \right. \\ &\quad \left. - \frac{1}{2} (2 + t)^2 + \text{const.} \right]. \end{aligned}$$

Here and later c is a constant, not always the same. We let $[\]$ stand for the expression in brackets and obtain

$$u(x) < c \int_0^x (2 + t^2)^{-3/2} \log^{-3}(2 + t^2) [\] dt.$$

The dominating term in the integrand is majorized by

$$\frac{1}{2} t^{-3} \log^{-3} t \times (2 + t)^2 \log(2 + t).$$

The integral from 1 to $x > 1$ is bounded, and consequently so is u for all x .

This completes the proof of Theorem 1.

2. The following simple example, valid for $N \geq 3$, is perhaps also of interest. Let

$$T : \quad |x| < \infty, \quad |y| \leq \pi, \quad |z_i| \leq 1, \quad i = 1, \dots, N - 2,$$

with the metric

$$ds^2 = dx^2 + e^{-x} dy^2 + e^{(2e^x - x)/(N-2)} \sum_{i=1}^{N-2} dz_i^2,$$

the opposite faces again identified by pairs. Clearly $T \in C$.

The function

$$u = \cos y$$

belongs to H^2B . In fact,

$$\begin{aligned} \Delta u &= -e^{-e^x+x} (e^{e^x-x} e^x) (-\cos y) \\ &= e^x \cos y, \end{aligned}$$

and

$$\Delta^2 u = -e^{-e^x+x} [(e^{e^x-x} e^x)' \cos y + e^{e^x-x} e^x e^x (-\cos y)] = 0.$$

Thus $T \in C \cap \tilde{\mathcal{G}}_{H^2B}$.

3. The reason that we are only interested in nonharmonic biharmonic functions is, of course, that completeness is known not

to exclude bounded harmonic functions (Nakai-Sario [6]). For $N \geq 3$, we insert here a simple proof of this fact.

Take the N -cylinder

$$T: |x| < \infty, \quad |y| \leq 1, \quad |z_i| \leq 1, \quad i = 1, \dots, N - 2,$$

with the metric

$$ds^2 = dx^2 + e^{2x^2} dy^2 + \sum_{i=1}^{N-2} dz_i^2.$$

Trivially $T \in C$. The function

$$h(x) = \int_0^x e^{-t^2} dt$$

is harmonic,

$$\Delta h = -e^{-x^2} (e^{x^2} e^{-x^2})' = 0.$$

It also is bounded and, in fact, even Dirichlet finite :

$$D(h) = c \int_{-\infty}^{\infty} e^{-2x^2} e^{x^2} dx < \infty.$$

4. We return to nonharmonic biharmonic functions.

THEOREM 2. — $C \cap \mathcal{O}_{H^2B}^N \neq \emptyset$ for every N .

Proof. — The Euclidean N -space $E^N \in C$. Every biharmonic function u has an expansion in spherical harmonics S_{nm}

$$u = \sum_{n=0}^{\infty} \sum_{m=1}^{m_n} (a_{nm} r^{n+2} + b_{nm} r^n) S_{nm}.$$

If $u \in H^2B$, then

$$\int_{|x|=r} u S_{nm} d\omega = c(a_{nm} r^{n+2} + b_{nm} r^n)$$

is bounded in r , hence $a_{nm} = b_{nm} = 0$ for all n , except for b_{01} . Therefore u is constant.

5. In view of $u = r^2 \in H^2B$ on the Euclidean N -ball, we have trivially $\tilde{C} \cap \tilde{\mathcal{O}}_{H^2B}^N \neq \emptyset$ for every N , with \tilde{C} the totality of incomplete Riemannian manifolds. It remains to show :

THEOREM 3. — $C \cap \mathcal{O}_{H^2B}^N \neq \emptyset$ for every N .

Proof. — Let E_α^N be the N -space $0 < r < \infty$ with the metric

$$ds = r^\alpha |dx|,$$

α a constant. It is known (Sario-Wang [19, 21]) that if $N \geq 4$, $E_\alpha^N \in \mathcal{O}_{H^2B}$ for every α ; $E_\alpha^2 \in \mathcal{O}_{H^2B}$ if and only if $\alpha \neq -1 \mp n/2$, $n = 1, 2, \dots$; $E_\alpha^3 \in \mathcal{O}_{H^2B}$ if and only if $\alpha \neq -1 \mp \left[\frac{1}{2} n(n+1) \right]^{\frac{1}{2}}$. On the other hand, $E_\alpha^N \in \widetilde{C}$ for every α , hence the theorem.

The author is sincerely indebted to Professor Cecilia Wang for a painstaking checking of the manuscript.

BIBLIOGRAPHY

- [1] D. Hada, L. Sario, C. Wang, Dirichlet finite biharmonic functions on the Poincaré N -ball, *J. Reine Angew. Math.* (to appear).
- [2] D. Hada, L. Sario, C. Wang, N -manifolds carrying bounded but no Dirichlet finite harmonic functions, *Nagoya Math. J.* (to appear).
- [3] Y.K. Kwon, L. Sario, B. Walsh, Behavior of biharmonic functions on Wiener's and Royden's compactifications, *Ann. Inst. Fourier (Grenoble)* 21 (1971), 217-226.
- [4] N. Mirsky, L. Sario, C. Wang, Bounded polyharmonic functions and the dimension of the manifold, *J. Math. Kyoto Univ.*, 13 (1973), 529-535.
- [5] M. Nakai, Dirichlet finite biharmonic functions on the plane with distorted metrics, (to appear).
- [6] M. Nakai, L. Sario, Completeness and function-theoretic degeneracy of Riemannian spaces, *Proc. Nat. Acad. Sci.*, 57 (1967), 29-31.
- [7] M. Nakai, L. Sario, Biharmonic classification of Riemannian manifolds, *Bull. Amer. Math. Soc.*, 77 (1971), 432-436.
- [8] M. Nakai, L. Sario, Quasiharmonic classification of Riemannian manifolds, *Proc. Amer. Math. Soc.*, 31 (1972), 165-169.

- [9] M. Nakai, L. Sario, Dirichlet finite biharmonic functions with Dirichlet finite Laplacians, *Math. Z.*, 122 (1971), 203-216.
- [10] M. Nakai, L. Sario, A property of biharmonic functions with Dirichlet finite Laplacians, *Math.Scand.* 29 (1971), 307-316.
- [11] M. Nakai, L. Sario, Existence of Dirichlet finite biharmonic functions, *Ann. Acad. Sci. Fenn. A.I.*, 532 (1973), 1-34.
- [12] M. Nakai, L. Sario, Existence of bounded biharmonic functions, *J. Reine Angew. Math.* 259 (1973), 147-156.
- [13] M. Nakai, L. Sario, Existence of bounded Dirichlet finite biharmonic functions, *Ann. Acad. Sci. Fenn. A.I.* 505 (1972), 1-12.
- [14] M. Nakai, L. Sario, Biharmonic functions on Riemannian manifolds, *Continuum Mechanics and Related Problems of Analysis*, Nauka, Moscow, 1972, 329-335.
- [15] L. Sario, Biharmonic and quasiharmonic functions on Riemannian manifolds, *Duplicated lecture notes 1968-70*, University of California, Los Angeles.
- [16] L. Sario, M. Nakai, *Classification Theory of Riemann Surfaces*, Springer-Verlag, 1970, 446 pp.
- [17] L. Sario, C. Wang, The class of (p, q) -biharmonic functions, *Pacific J. Math.*, 41 (1972), 799-808.
- [18] L. Sario, C. Wang, Counterexamples in the biharmonic classification of Riemannian 2-manifolds, *Pacific J. Math.* (to appear).
- [19] L. Sario, C. Wang, Generators of the space of bounded biharmonic functions, *Math. Z.*, 127 (1972), 273-280.
- [20] L. Sario, C. Wang, Quasiharmonic functions on the Poincaré N -ball, *Rend. Mat.* (to appear).
- [21] L. Sario, C. Wang, Riemannian manifolds of dimension $N \geq 4$ without bounded biharmonic functions, *J. London Math. Soc.* (to appear).
- [22] L. Sario, C. Wang, Existence of Dirichlet finite biharmonic functions on the Poincaré 3-ball, *Pacific J. Math.*, 48 (1973), 267-274.
- [23] L. Sario, C. Wang, Negative quasiharmonic functions, *Tôhoku Math. J.*, 26 (1974), 85-93.
- [24] L. Sario, C. Wang, Radial quasiharmonic functions, *Pacific J. Math.*, 46 (1973), 515-522.

COMPLETENESS AND EXISTENCE OF BOUNDED BIHARMONIC FUNCTIONS

- [25] L. Sario, C. Wang, Parabolicity and existence of bounded biharmonic functions, *Comm. Math. Helv.* 47, (1972), 341-347.
- [26] L. Sario, C. Wang, Positive harmonic functions and biharmonic degeneracy, *Bull. Amer. Math. Soc.*, 79 (1973), 182-187.
- [27] L. Sario, C. Wang, Parabolicity and existence of Dirichlet finite biharmonic functions, *J. London Math. Soc.* (to appear).
- [28] L. Sario, C. Wang, Harmonic and biharmonic degeneracy, *Kodai Math. Sem. Rep.* 25 (1973), 392-396.
- [29] L. Sario, C. Wang, M. Range, Biharmonic projection and decomposition, *Ann. Acad. Sci. Fenn. A.I.*, 494 (1971), 1-14.
- [30] C. Wang, L. Sario, Polyharmonic classification of Riemannian manifolds, *Kyoto Math. J.*, 12 (1972), 129-140.

Manuscrit reçu le 22 mai 1973
accepté par M. Brelot.

Leo SARIO
Department of Mathematics
University of California
Los Angeles, Calif. (USA)