

YOSHIHIRO MIZUTA

**On the boundary limits of harmonic functions
with gradient in L^p**

Annales de l'institut Fourier, tome 34, n° 1 (1984), p. 99-109

http://www.numdam.org/item?id=AIF_1984__34_1_99_0

© Annales de l'institut Fourier, 1984, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE BOUNDARY LIMITS OF HARMONIC FUNCTIONS WITH GRADIENT IN L^p

by Yoshihiro MIZUTA

1. Introduction.

Let u be a function harmonic in the half space

$$\mathbf{R}_+^n = \{x = (x_1, \dots, x_n); x_n > 0\}$$

and satisfying the condition :

$$\int_G |\text{grad } u(x)|^p x_n^\alpha dx < \infty \quad (1)$$

for any bounded open set $G \subset \mathbf{R}_+^n$, where $p > 1$ and $\alpha < p - 1$.
For $\xi \in \partial \mathbf{R}_+^n$, $\gamma \geq 1$ and $a > 1$, set

$$T_\gamma(\xi, a) = \{x \in \mathbf{R}_+^n; |x - \xi| < ax_n^{1/\gamma}\}.$$

The existence of nontangential limits of u , that is, the limit of $u(x)$ as $x \rightarrow \xi$, $x \in T_1(\xi, a)$, was studied by Carleson [1] ($n = p = 2$ and $0 \leq \alpha < 1$), Wallin [10] ($p = 2$ and $0 \leq \alpha < 1$) and Mizuta [6] in the present situation.

Recently Cruzeiro [2] proved the existence of $\lim u(x)$ as $x \rightarrow \xi$, $x \in T_\gamma(\xi, a)$, for a harmonic function u satisfying (1) with $p = n$ and $\alpha = 0$. The existence of such limits for Green potentials in \mathbf{R}_+^n was obtained by Wu [11]. Taking these results into account, we give the following theorem :

THEOREM. — *Let u be a function harmonic in \mathbf{R}_+^n and satisfying (1) with $p > 1$ and $\alpha < p - 1$.*

(i) *If $n - p + \alpha > 0$, then for each $\gamma > 1$ there exists a set $E_\gamma \subset \partial \mathbf{R}_+^n$ such that $H_{\gamma(n-p+\alpha)}(E_\gamma) = 0$ and*

$$\lim_{x \rightarrow \xi, x \in T_\gamma(\xi, a)} u(x) \quad (2)$$

exists and is finite for any $\xi \in \partial \mathbf{R}_+^n - E_\gamma$ and any $a > 1$.

(ii) If $n - p + \alpha = 0$, then there exists a set $E \subset \partial \mathbf{R}_+^n$ such that $B_{n/p, p}(E) = 0$ and (2) exists and is finite for any $\xi \in \partial \mathbf{R}_+^n - E$, any $\gamma \geq 1$ and any $a > 1$.

(iii) If $n - p + \alpha < 0$, then $\lim_{x \rightarrow \xi, x \in \mathbf{R}_+^n} u(x)$ exists and is finite for any $\xi \in \partial \mathbf{R}_+^n$.

Here H_d denotes the d -dimensional Hausdorff measure, and $B_{\beta, p}$ the Bessel capacity of index (β, p) (cf. [3]). In view of [3; Theorems 21 and 22], one notes the following results:

a) If $q > 1$ and $n - d \geq \beta q$, then $H_d(E) < \infty$ implies $B_{\beta, q}(E) = 0$;

b) If $q > 1$ and $n - d < \beta q$, then $B_{\beta, q}(E) = 0$ implies $H_d(E) = 0$.

Recently Nagel, Rudin and Shapiro [7] studied tangential behaviors of Poisson integrals of potential type functions (see Sec. 4, Remark 2). Their results are not applicable to our case unless $p = 2$.

Remark. — The same result as in the theorem is also valid for a domain Ω for which any function v satisfying

$$\int_{\Omega} |\text{grad } v(x)|^p \delta(x)^\alpha dx < \infty, \quad p > 1, \alpha < p - 1, \quad (3)$$

can be extended to a function satisfying (3) with Ω replaced by \mathbf{R}^n , where $\delta(x)$ denotes the distance from x to $\partial \Omega$. The special Lipschitz domains in [9; Chap. VI] are typical examples of Ω .

2. Lemmas.

First we note the following result, which follows readily from the fact in [4; p. 165].

LEMMA 1. — Let f be a locally integrable function on \mathbf{R}^n . For $\beta > 0$, we set

$$A_\beta = \{ \xi \in \partial \mathbf{R}_+^n ; \limsup_{r \downarrow 0} r^{-\beta} \int_{B(\xi, r)} |f(y)| dy > 0 \},$$

where $B(x, r)$ denotes the open ball with center at x and radius r . Then $H_\beta(A_\beta) = 0$.

By [3 ; Theorem 21] and the result of [4 ; p. 165], we have

LEMMA 2. — Let f be a locally integrable function on \mathbf{R}^n . For $p > 1$, we set

$$B_p = \{ x ; \limsup_{r \downarrow 0} (\log r^{-1})^{p-1} \int_{B(x, r)} |f(y)| dy > 0 \}.$$

Then $B_{n/p, p}(B_p) = 0$.

Next we prove the following technical result.

LEMMA 3. — Let $c_1 > 0$, $c_2 > 0$, $\gamma \geq 1$, $p > 1$ and $p - n \leq \alpha < p - 1$. Then

$$\left\{ \int_{\{y ; c_1 x_n < |x-y| < c_2 x_n^{1/\gamma}\}} |x-y|^{p'(1-n)} |y_n|^{-\alpha p'/p} dy \right\}^{1/p'} \leq C \begin{cases} x_n^{(p-n-\alpha)/p} & \text{if } n-p+\alpha > 0, \\ (\log x_n^{-1})^{1/p'} & \text{if } n-p+\alpha = 0, \end{cases}$$

where $1/p + 1/p' = 1$ and C is a positive constant independent of $x = (x_1, \dots, x_n)$ with $0 < x_n < 1/2$.

Proof. — Let $x^* = (0, \dots, 0, 1)$. By change of variables we see that the left hand side is equal to

$$x_n^{1-n-\alpha/p+n/p'} \left\{ \int_{E_1} |x^* - z|^{p'(1-n)} |z_n|^{-\alpha p'/p} dz \right\}^{1/p'} \leq C x_n^{(p-n-\alpha)/p} \left\{ \int_{E_2} (1 + |z|)^{p'(1-n)} |z_n|^{-\alpha p'/p} dz \right\}^{1/p'},$$

where $E_1 = \{z, c_1 < |x^* - z| < c_2 x_n^{1/\gamma-1}\}$,

$$E_2 = B(0, (c_2 + 1) x_n^{1/\gamma-1}),$$

C is a positive constant independent of x with $0 < x_n < 1$. The required inequalities are established by estimating the last integral.

In the same manner we can prove

LEMMA 4. — Let $p > 1$ and $\alpha < p - n$. Then

$$\left\{ \int_{\{y: x_n/2 < |x-y| < |x|/2\}} |x-y|^{p'(1-n)} |y_n|^{-\alpha p'/p} dy \right\}^{1/p'} \leq C |x|^{(p-n-\alpha)/p}$$

for any $x \in \mathbf{R}_+^n$, where C is a positive constant independent of x .

Finally we borrow a result from [6 ; Lemma 4].

LEMMA 5. — Let $p > 1$, $\alpha < p - 1$ and f be a measurable function on \mathbf{R}^n such that $\int_G |f(y)|^p |y_n|^\alpha dy < \infty$ for any bounded open set $G \subset \mathbf{R}^n$. If we set

$$E' = \left\{ \xi \in \partial \mathbf{R}_+^n; \int_{B(\xi, 1)} |\xi - y|^{1-n} |f(y)| dy = \infty \right\},$$

then $B_{1-\alpha/p, p}(E') = \emptyset$.

Remark. — If $p - \alpha > n$, then one sees that E' is empty.

3. Proof of the theorem.

Take q such that $q = p$ if $\alpha \leq 0$ and $1 < q < p/(\alpha + 1)$ if $\alpha > 0$. Then Hölder's inequality implies that

$$\int_G |\text{grad } u(x)|^q dx < \infty$$

for any bounded open set $G \subset \mathbf{R}_+^n$. Hence the function

$$v(x) = \begin{cases} u(x_1, \dots, x_{n-1}, x_n) & \text{for } x_n > 0, \\ u(x_1, \dots, x_{n-1}, -x_n) & \text{for } x_n < 0, \end{cases}$$

can be extended to a locally q -precise function w on \mathbf{R}^n in view of [8 ; Theorem 5.6] (for q -precise functions, see Ohtsuka [8 ; Chap. IV] and Ziemer [12]). Define

$$E' = \left\{ \xi \in \partial \mathbf{R}_+^n ; \int_{B(\xi, 1)} |\xi - y|^{1-n} |\text{grad } w(y)| dy = \infty \right\}.$$

Then we have $B_{1-\alpha/p, p}(E') = 0$ on account of Lemma 5. We also define A_β and B_p with $f(y) = |\text{grad } w(y)|^p |y_n|^\alpha$. By Lemmas 1 and 2, we see that $H_\beta(A_\beta) = 0$ and $B_{n/p, p}(B_p) = 0$.

First suppose $n - p + \alpha > 0$. Let $\gamma > 1$ be given. We shall show that (2) exists and is finite for any

$$\xi \in \partial \mathbf{R}_+^n - (E' \cup A_{\gamma(n-p+\alpha)})$$

and any $a > 1$. Let $\xi \in \partial \mathbf{R}_+^n - (E' \cup A_{\gamma(n-p+\alpha)})$. Take $N > 1$ such that $\xi \in B(0, N)$, and let ϕ_N be a function in $C_0^\infty(\mathbf{R}^n)$ such that $\phi_N = 1$ on $B(0, 2N)$ and $\phi_N = 0$ on $\mathbf{R}^n - B(0, 3N)$. Set $w_N = \phi_N w$. Then w_N is q -precise on \mathbf{R}^n and satisfies

$$w_N(x) = c \sum_{i=1}^n \int \left(\frac{\partial}{\partial x_i} R_2 \right) (x - y) \frac{\partial w_N}{\partial y_i} dy \quad \text{for } x \in \mathbf{R}_+^n,$$

where c is a constant, $R_2(x) = \log(1/|x|)$ if $n = 2$ and $R_2(x) = |x|^{2-n}$ if $n \geq 3$. In fact, since w_N is continuously differentiable on \mathbf{R}_+^n , the right hand side is continuous on \mathbf{R}_+^n and the required equality holds for any $x \in \mathbf{R}_+^n$ on account of Ohtsuka [8 ; Theorem 9.11].

For $x \in \mathbf{R}_+^n$, we write

$$w_N(x) = w_{N,1}(x) + w_{N,2}(x) + w_{N,3}(x),$$

where

$$w_{N,1}(x) = c \sum_{i=1}^n \int_{B(x, x_n/2)} \left(\frac{\partial}{\partial x_i} R_2 \right) (x - y) \frac{\partial w_N}{\partial y_i} dy,$$

$$w_{N,2}(x) = c \sum_{i=1}^n \int_{B(x, |\xi - x|/2) - B(x, x_n/2)} \left(\frac{\partial}{\partial x_i} R_2 \right) (x - y) \frac{\partial w_N}{\partial y_i} dy,$$

$$w_{N,3}(x) = c \sum_{i=1}^n \int_{\mathbf{R}^n - B(x, |\xi - x|/2)} \left(\frac{\partial}{\partial x_i} R_2 \right) (x - y) \frac{\partial w_N}{\partial y_i} dy.$$

Since w_N is harmonic on $B(0, 2N) \cap \mathbf{R}_+^n$, $w_{N,1}(x) = 0$ for $x \in B(0, N) \cap \mathbf{R}_+^n$. It follows from our assumption $\xi \notin E'$ that $\int |\xi - y|^{1-n} |\text{grad } w_N(y)| dy < \infty$. Hence Lebesgue's dominated convergence theorem gives

$$\lim_{x \rightarrow \xi, x \in \mathbf{R}_+^n} w_{N,3}(x) = c \sum_{i=1}^n \int \left(\frac{\partial}{\partial x_i} R_2 \right) (\xi - y) \frac{\partial w_N}{\partial y_i} dy.$$

For $w_{N,2}$ we apply Hölder's inequality to obtain by Lemma 3, $|w_{N,2}(x)|$

$$\begin{aligned} &\leq \text{const.} \left\{ \int_{\{y; x_n/2 < |x-y| < |\xi-x|/2\}} |x-y|^{p'(1-n)} |y_n|^{-\alpha p'/p} dy \right\}^{1/p'} \\ &\times \left\{ \int_{B(x, |\xi-x|/2)} |\text{grad } w_N(y)|^p |y_n|^\alpha dy \right\}^{1/p} \\ &\leq \text{const.} \left\{ x_n^{p-\alpha-n} \int_{B(\xi, 2ax_n^{1/\gamma})} |\text{grad } w(y)|^p |y_n|^\alpha dy \right\}^{1/p} \end{aligned}$$

for $x \in B(0, N) \cap T_\gamma(\xi, a)$. Since $\xi \notin A_{\gamma(n-p+\alpha)}$,

$$\lim_{x \rightarrow \xi, x \in T_\gamma(\xi, a)} w_{N,2}(x) = 0.$$

Thus
$$\begin{aligned} \lim_{x \rightarrow \xi, x \in T_\gamma(\xi, a)} u(x) &= \lim_{x \rightarrow \xi, x \in T_\gamma(\xi, a)} w_N(x) \\ &= \lim_{x \rightarrow \xi, x \in T_\gamma(\xi, a)} w_{N,3}(x), \end{aligned}$$

which is finite.

If $n - p + \alpha = 0$, then we can prove that (2) exists and is finite for any $\xi \in \partial \mathbf{R}_+^n - (E' \cup B_p)$, any $\gamma \geq 1$ and any $a > 1$.

If $n - p + \alpha < 0$, then E' is empty, so that similar arguments yield the required assertion with the aid of Lemma 4.

4. Remarks.

Remark 1. - If $p > n$, $p - \alpha - n > 0$ and u is a locally p -precise function on \mathbf{R}_+^n satisfying (1), then $u(x)$ has a finite limit as $x \rightarrow \xi$, $x \in \mathbf{R}_+^n$, for any $\xi \in \partial \mathbf{R}_+^n$.

Remark 2. — Nagel, Rudin and Shapiro [7] studied tangential behaviors of functions of the form

$$(P_{x_n} * (K * g))(x'), \quad x = (x', x_n) \in \mathbf{R}_+^n,$$

where P is the Poisson kernel in \mathbf{R}_+^n , K is a nonnegative kernel, which is radial and decreasing, and g is a function in $L^p(\mathbf{R}^{n-1})$. The function in our theorem has a boundary value in the Lipschitz space $\Lambda_{\beta}^{p,p}(\mathbf{R}^{n-1})$ with $\beta = 1 - (\alpha + 1)/p$ locally, provided $-1 < \alpha < p + 1$ (cf. [9; Chap. VI, §§ 4.3, 4.5]). We do not know whether functions f in $\Lambda_{\beta}^{p,p}(\mathbf{R}^{n-1})$ can be written as $f = K * g$, where K is an appropriate kernel function which is determined independently of f and $g \in L^p(\mathbf{R}^{n-1})$.

If $g \in L^p(\mathbf{R}^n)$, $\beta = 1 - (\alpha + 1)/p$ and $-1 < \alpha < p - 1$,

then

$$F(x') = \int g_{1-\alpha/p}((x', 0) - y) g(y) dy$$

belongs to $\Lambda_{\beta}^{p,p}(\mathbf{R}^{n-1})$, where g_{ρ} denotes the Bessel kernel of order ρ (cf. [9; Chap. VI, § 4.3]). Hence $u(x) = P_{x_n} * F(x')$ satisfies

$$\int_0^{\infty} \left[x_n^{k-\beta} \left\{ \int_{\mathbf{R}^{n-1}} \left| \left(\frac{\partial}{\partial x_n} \right)^k u(x', x_n) \right|^p dx' \right\}^{1/p} \right]^p x_n^{-1} dx_n < \infty,$$

where k is an integer greater than β , in view of [9; p. 152]. This implies, by the observation given after Lemma 4' in [9; Chap. V], that u satisfies

$$\int_{\mathbf{R}_+^n} |\text{grad } u(x)|^p x_n^{\alpha} dx < \infty. \tag{1}'$$

Thus our theorem is applicable to this function u .

Remark 3. — In case $p - \alpha - n = 0$, our theorem gives the best possible result as to the size of the exceptional set as the next proposition shows.

PROPOSITION . — *Let E be a compact set in $\partial \mathbf{R}_+^n$ with $B_{1-\alpha/p,p}(E) = 0$, where $p > 1$ and $-1 < \alpha < p - 1$. Then there exists a function u which is harmonic in \mathbf{R}_+^n and satisfies (1)' such that $\lim_{x_n \downarrow 0} u(x', x_n)$ does not exist for any $(x', 0) \in E$.*

Proof. – Since $B_{1-\alpha/p,p}(E) = 0$, we can find a nonnegative function $f \in L^p$ such that

$$F(x') \equiv \int g_{1-\alpha/p}((x', 0) - y) f(y) dy = \infty$$

for any $(x', 0) \in E$. As seen above, $P_{x_n} * F(x')$ satisfies (1)'. Take a_1, b_1 and c_1 such that

$$0 < a_1 < 1, \quad 0 < b_1 < c_1 < 1$$

and $P_{a_1} * F_1(x') > 1$ for $(x', 0) \in E$, where

$$F_1(x') = \int_{\{b_1 \leq |y_n| < c_1\}} g_{1-\alpha/p}((x', 0) - y) f(y) dy.$$

We proceed inductively and obtain $\{a_j\}, \{b_j\}$ and $\{c_j\}$ such that $0 < a_{j+1} < a_j, \quad 0 < c_{j+1} < b_j < c_j, \quad P_{a_k} * F_j(x') < 2^{-j}$

if $k < j$ and $(x', 0) \in E$ and $P_{a_j} * F_j(x') > j + \sum_{k=1}^{j-1} M_k$ if $(x', 0) \in E$, where

$$F_j(x') = \int_{\{b_j \leq |y_n| < c_j\}} g_{1-\alpha/p}((x', 0) - y) f(y) dy$$

and $M_k = \max \{F_k(x'); (x', 0) \in \partial R_+^n\}$.

Define

$$u(x', x_n) = \sum_{j=1}^{\infty} (-1)^j P_{x_n} * F_j(x').$$

Then one sees easily that $\{u(x', a_j)\}$ does not converge as $j \rightarrow \infty$ for any $(x', 0) \in E$. Since u satisfies (1)' and is harmonic in R_+^n , u is the required function.

Remark 4. – Let $p > 1, \quad p - n < \alpha < p - 1$ and $\alpha > -1$. Then we can find a function u harmonic in R_+^n and satisfying the following conditions :

- (i) u has a finite nontangential limit at 0 ;
- (ii) $\limsup_{x \rightarrow 0, x \in T_\gamma(0, a)} u(x) = \infty$ for any $\gamma > 1$ and any $a > 1$;
- (iii) $\int_{R_+^n} |\text{grad } u(x)|^p x_n^\alpha dx < \infty$.

To see this, let $x^{(j)} = (2^{-j}, 0, \dots, 0$ and define

$$f_j(y) = \begin{cases} a_j |x^{(j)} - y|^{-1} & \text{if } y \in B(x^{(j)}, 2^{-j-2}) - \mathbf{R}_+^n, \\ 0 & \text{otherwise,} \end{cases}$$

where a_j is a positive number determined later. Setting

$$u(x) = \int (x_n - y_n) |x - y|^{-n} f(y) dy,$$

where $f = \sum_{j=1}^{\infty} f_j$, we note the following facts :

a) $u(0) \leq \int |y|^{1-n} f(y) dy \leq C_1 \sum_{j=1}^{\infty} a_j$;

b) If $x = (2^{-j}, 0, \dots, 0, x_n)$ and $0 < x_n < 2^{-j-1}$, then $u(x) \geq C_2 a_j \log(2^{-j}/x_n)$;

c) $\int f(y)^p |y_n|^\alpha dy \leq C_3 \sum_{j=1}^{\infty} a_j^p 2^{-j(n-p+\alpha)}$,

where C_1, C_2 and C_3 are positive constants independent of x and j .

Now we choose $\{a_j\}$ so that (a) and (c) are finite but $\limsup_{j \rightarrow \infty} ja_j = \infty$. Then (a) implies that u has a nontangential limit at 0, and (iii) follows from (c) and [5; Lemma 6](*). By (b) and the construction of $\{a_j\}$, (ii) is fulfilled. Thus u satisfies (i), (ii) and (iii).

Remark 5. - If $p > 1$ and $\alpha = p - n > -1$, then for each $\gamma > 1$ there exists a function u harmonic in \mathbf{R}_+^n such that :

- (i) u has a finite nontangential limit at 0;
- (ii) $\limsup_{x \rightarrow 0, x \in T_{\gamma'}(\xi, a)} u(x) = \infty$ for any $\gamma' \geq \gamma$ and any $a > 1$;
- (iii) $\int_{\mathbf{R}_+^n} |\text{grad } u(x)|^p x_n^\alpha dx < \infty$.

In fact we modify above f_j by setting

$$f_j(y) = \begin{cases} a_j |x^{(j)} - y|^{-1} & \text{if } y \in B(x^{(j)}, 2^{-j-2}) - B(x^{(j)}, 2^{-\gamma j}) - \mathbf{R}_+^n, \\ 0 & \text{otherwise,} \end{cases}$$

(*) One notes that the conclusion of [5; Lemma 6] is true in case $p > 1$ and $-1 < \alpha < p - 1$ if g in the lemma has compact support.

and consider $u(x) = \sum_{j=1}^{\infty} \int (x_n - y_n) |x - y|^{-n} f_j(y) dy$. Then as in Remark 3 we can choose $\{a_j\}$ such that (i), (ii) and (iii) hold.

Remark 6. — Let $p > 1$ and h be a positive function on the interval $(0, \infty)$ such that $\lim_{r \rightarrow 0} h(r) (\log r^{-1})^{p-1} = 0$. Define for $\xi \in \partial \mathbf{R}_+^n$,

$$T_h(\xi) = \left\{ x \in \mathbf{R}_+^n ; \log \frac{|x - \xi|}{x_n} \leq h(|x - \xi|)^{-1/(p-1)} \right\}.$$

If u is a function harmonic in \mathbf{R}_+^n and satisfying (1) with $\alpha = p - n > -1$, then $\lim_{x \rightarrow \xi, x \in T_h(\xi)} u(x)$ exists and is finite for any $\xi \in \partial \mathbf{R}_+^n - (E' \cup B_h)$, where

$$B_h = \left\{ \xi \in \partial \mathbf{R}_+^n ; \limsup_{r \rightarrow 0} h(r)^{-1} \int_{B(\xi, r) \cap \mathbf{R}_+^n} |\text{grad } u(x)|^p x_n^\alpha dx > 0 \right\}.$$

BIBLIOGRAPHY

- [1] L. CARLESON, *Selected Problems on exceptional sets*, Van Nostrand, Princeton, 1967.
- [2] A.B. CRUZEIRO, Convergence au bord pour les fonctions harmoniques dans \mathbf{R}^d de la classe de Sobolev W_1^d , *C.R.A.S.*, Paris, 294 (1982), 71-74.
- [3] N.G. MEYERS, A theory of capacities for potentials in Lebesgue classes, *Math. Scand.*, 26 (1970), 255-292.
- [4] N.G. MEYERS, Continuity properties of potentials, *Duke Math. J.*, 42 (1975), 157-166.
- [5] Y. MIZUTA, On the existence of boundary values of Beppo Levi functions defined in the upper half space of \mathbf{R}^n , *Hiroshima Math. J.*, 6 (1976), 61-72.
- [6] Y. MIZUTA, Existence of various boundary limits of Beppo Levi functions of higher order, *Hiroshima Math. J.*, 9 (1979), 717-745.

- [7] A. NAGEL, W. RUDIN and J.H. SHAPIRO, Tangential boundary behavior of functions in Dirichlet-type spaces, *Ann. of Math.*, 116 (1982), 331-360.
- [8] M. OHTSUKA, Extremal length and precise functions in 3-space, *Lecture Notes, Hiroshima Univ.*, 1973.
- [9] E.M. STEIN, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, 1970.
- [10] H. WALLIN, On the existence of boundary values of a class of Beppo Levi functions, *Trans. Amer. Math. Soc.*, 120 (1965), 510-525.
- [11] J.-M. G. WU, L^p -densities and boundary behaviors of Green potentials, *Indiana Univ. Math. J.*, 28 (1979), 895-911.
- [12] W.P. ZIEMER, Extremal length as a capacity, *Michigan Math. J.*, 17 (1970), 117-128.

Manuscrit reçu le 2 novembre 1982.

Yoshihiro MIZUTA,
Department of Mathematics
Faculty of Integrated Arts and Sciences
Hiroshima University
Hiroshima 730 (Japan).