

ANNALES DE L'INSTITUT FOURIER

JEAN-PIERRE ROSAY

New examples of non-locally embeddable CR structures (with no non-constant CR distributions)

Annales de l'institut Fourier, tome 39, n° 3 (1989), p. 811-823

http://www.numdam.org/item?id=AIF_1989__39_3_811_0

© Annales de l'institut Fourier, 1989, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

NEW EXAMPLES OF NON-LOCALLY EMBEDDABLE *CR* STRUCTURES (WITH NO NON-CONSTANT *CR* DISTRIBUTIONS)

by Jean-Pierre ROSAY^(*)

Several years ago, L. Nirenberg gave a famous example of a strictly pseudoconvex, non-locally embeddable, *CR* structure on a manifold of dimension 3, ([7], [8] see also [4]).

This phenomenon of non-embeddability is important enough in order that it may be worthwhile to give new examples which may shed a new light and seem to present some improvement. In the previous examples it seems that only the existence of C^1 *CR* function could be investigated. We can just as well handle the case of *CR* distributions. There may be more than just a theoretical interest to it, since, for example, L^∞ functions arise quite naturally as weak limits.

Our example has the possible additional feature that it can be adapted so that the *CR* structure that we construct can be extended to a complex structure on the concave side of the manifold. This relates to a theorem by D. Catlin [14], and an example (in dimension 5, with degenerate Levi form) by D. Hill, who conjectured that such one sided extensions to the concave side always exist, and asked explicitly the question for Nirenberg's example [2]. Full details will not be provided, but some indications (that we hope to be sufficient) are given in a remark at the end of the paper. It seems also that our examples complete some examples by Jacobowitz [3], concerning the canonical bundle.

(*) Research partly supported by NSF grant DMS 8800610.

Key-word : *CR* structure.

A.M.S. Classification : 53B25 – 32E20.

In part I, we describe a simple “basic construction” from which everything follows easily. This is the main part of the paper. In II, we get immediately an example of a strictly pseudo convex *CR* structure on a manifold of dimension 3 (arbitrarily close to any given structure, as in [4]), for which there is a point so that every *CR* distribution defined in a connected neighborhood of this point, must be constant.

We get also examples of structures with only one “independent solution” (see further, for more details).

In III, we give a new approach to results by Jacobowitz and Trèves [4]. Jacobowitz and Trèves were the first to show that the phenomenon of non-embeddability occurs too in higher dimension for structures whose Levi form is non-degenerate and has only one positive eigenvalue. Then all *CR* functions, or distributions, must be smooth, if the *CR* structure is smooth ([12] page 493- , or [10]). In III we give examples of non-embeddable *CR* structure on manifolds of dimension $2m - 1$, whose Levi form has one positive and $(m - 2)$ negative eigenvalue (arbitrarily close to any given embedded structure), with locally, no non-constant *CR* functions. Or with only k “independent solutions” (k arbitrary $k \in \{0, \dots, m\}$) : z_1, \dots, z_k which satisfy $dz_1 \wedge \dots \wedge dz_k \neq 0$ and so that every *CR* function is a function of (z_1, \dots, z_k) . By *CR* structure we will always mean *CR* structure of hypersurface type, and we will consider only smooth (C^∞) structure.

Let us remind the reader that it is still an open question whether any strictly pseudoconvex *CR* structure of dimension 5 can be locally embedded (for dimension > 7 , this is known to be true [1], [5], [13]). Our example sheds some light on the difficulty to build a counterexample, from the point of view of function theory.

I. BASIC CONSTRUCTION

1. Lupacciolu’s theorem.

Although very simple, our construction relies entirely on a very nice result by G. Lupacciolu [6]. We are using only a very particular case, but it seems much more pleasant to state the result in some generality. A different and simple proof of Lupacciolu’s theorem is given in the appendix in [9].

“Let Ω be bounded domain in \mathbf{C}^n and K a polynomially convex compact set in $b\Omega$ (the boundary of Ω). If $b\Omega$ is smooth and strictly

pseudoconvex off K , then every CR distribution defined on $b\Omega - K$ has a holomorphic extension to Ω (i.e. in some sense K is negligible)."

Since every CR distribution on $b\Omega - K$ has a local holomorphic extension inside to Ω (due to strict pseudoconvexity), the case of CR distributions reduces easily to the case of smooth CR functions.

2. Construction of a CR structure.

We start with S a smooth strictly convex hypersurface defined in Ω , an open neighborhood of O in \mathbf{C}^2 . We assume that $O \in S$ and that $\text{Re } z_2 \geq 0$ on S (therefore > 0 on $S - \{0\}$). We fix $\beta > 0$, small enough in order that $S \cap \{\text{Re } z_2 \leq \beta\}$ is a compact set in Ω , and that the line segment from $(0, 0)$ to $(0, \beta)$ does not intersect S except at $(0, 0)$. For $0 < \alpha < \beta$ we set :

$$\begin{aligned} \Sigma &= S \cap \{\text{Re } z_2 < \beta\} \\ \Sigma^- &= S \cap \{\text{Re } z_2 < \alpha\} \\ \Sigma^+ &= S \cap \{\alpha < \text{Re } z_2 < \beta\} \\ S_\alpha &= S \cap \{\text{Re } z_2 = \alpha\}. \end{aligned}$$

Then, we consider $z_1^\#$ a smooth function defined on $\overline{\Sigma^-}$ with the following properties :

- (i) $\bar{\partial}_b z_1^\#$ vanishes to infinite order along S_α .
- (ii) $z_1^\#$ is sufficiently close to z_1 (say in the \mathcal{C}^2 topology) in order that the map $\tau : (z_1, z_2) \mapsto (z_1^\#(z_1, z_2), z_2)$ maps $\overline{\Sigma^-}$ into a strictly convex surface (diffeomorphically).

(iii) For $|y_2|$ small $z_1^\#(z_1, \alpha + iy_2) = z_1 + \frac{\epsilon}{z_1}$ for some positive ϵ , $\epsilon < \frac{d^2}{25}$, where d is the distance from $(0, \alpha)$ to S .

(iv) For $\text{Re } z_2 < \frac{2\alpha}{3}$ $z_1^\#(z_1, z_2) = z_1$.

To construct such a function $z_1^\#$ is extremely easy. One can set $z_1^\#(z_1, z_2) = z_1 + \frac{\epsilon}{z_1} \chi(z_2)$. For χ one picks a smooth cut-off function, defined in \mathbf{C} , which is identically 1 in a neighborhood of α , but has small support around α , and such that $\bar{\partial} \chi$ vanishes to infinite order along the

real line $\{\operatorname{Re} z = \alpha\}$. Notice, as a crucial fact, that for z_2 close to α , $z_1 \neq 0$ if $(z_1, z_2) \in S$. Once χ is chosen, it is enough to take ϵ small enough. Observe that one can therefore take $z_1^\#$ as close as wanted to z_1 (in the C^∞ topology).

It follows from (ii) that there is a unique CR structure on Σ^- such that $z_1^\#$ and z_2 are CR functions. It is the pullback of the structure induced from \mathbf{C}^2 on $\tau(\Sigma^-)$ and it is therefore a strictly pseudoconvex structure. It follows from (i) that this structure patches smoothly with the standard CR structure along S_α . On Σ^+ we use the standard CR structure (induced from \mathbf{C}^2).

This finally gives rise to a smooth CR structure on Σ which is strictly pseudoconvex, and for which we have as independant CR functions :

$$\begin{cases} \text{(a) on } \Sigma^-, z_1^\# \text{ and } z_2 \\ \text{(b) on } \Sigma^+, z_1 \text{ and } z_2. \end{cases}$$

As a last piece of notation, let $\Sigma' = \Sigma \cap \{\operatorname{Re} z_2 > \frac{\alpha}{2}\}$. For some applications we will need to consider functions defined only on Σ' . The only idea in our construction is in the following claim.

CLAIM — *It follows from (iii) that every CR function defined on Σ or Σ' , (equipped with the above CR structure) is a function of z_2 only. The same holds for CR distributions. In addition, any CR distribution on Σ (or Σ'), that is constant on Σ^+ , is constant.*

3. Proof of the claim.

3.1. For simplicity let us first treat the case of continuous CR function on Σ (or Σ'). The restriction of f to Σ^+ is a CR function (for the usual structure). By 1, it extends to a holomorphic function on the open set bounded by Σ and the hyperplanes $\{\operatorname{Re} x_2 = \alpha$ or $\beta\}$.

Let γ be the curve in the z_1 plane constituted by the points z_1 such that $(z_1, \alpha) \in S$. We conclude from what precedes that the function $z_1 \mapsto f(z_1, \alpha)$, defined on γ , extends holomorphically inside the convex curve γ , and is therefore the uniform limit on γ of a sequence of polynomials in z_1 .

The same holds, for $|y_2|$ small, on the corresponding curve

$\{(z_1 \in \mathbf{C}, (z_1, \alpha + iy_2) \in S)\}$, for the function $z_1 \mapsto f(z_1, \alpha + iy_2)$.

Together with $\tau(\Sigma^-)$, the hyperplane $\{\text{Re } z_2 = \alpha\}$ (or the hyperplanes $\{\text{Re } z_2 = \alpha \text{ or } \frac{\alpha}{2}\}$) bounds a domain to which we can again apply Lupacciolu's theorem, since CR functions on Σ^- for our structure are usual CR functions on $\tau(\Sigma^-)$.

The conclusion is that on the curve γ the function $z_1 \mapsto f(z_1, \alpha)$ is a uniform limit of polynomials in $z_1 + \frac{\epsilon}{z_1}$ ($= z_1^\#$). And, again, the same holds, for $|y_2|$ small, on the corresponding curves. Notice that 0 is inside γ .

It now follows from the above facts and from the Lemma below (which we immediately state for distributions), that for $|y_2|$ small $f(z_1, \alpha + iy_2)$ does not depend on z_1 . So the holomorphic extensions of f , from Σ^+ and $\tau(\Sigma^-)$ respectively, do not depend on z_1 . Hence f depends only on z_2 , as claimed. And it is also clear that if f is constant on Σ^+ then f is constant.

It remains to state and prove the following Lemma.

3.2. LEMMA. — *Let γ be a smooth closed curve in \mathbf{C} , so that $O \notin \gamma$ and O does not belong to the unbounded component of $\mathbf{C} - \gamma$. Let d be the distance from O to γ (or any smaller positive number). Let $\epsilon \in \mathbf{C}$, $0 < |\epsilon| < \frac{d^2}{25}$. Then no non-constant distribution on γ can be both the limit in the sense of distributions of :*

- (a) a sequence of polynomials in z
- (b) a sequence of polynomials in $z + \frac{\epsilon}{z}$.

Proof of the Lemma. — We can assume ϵ to be real and positive. Let τ_1 be the mapping defined on $\mathbf{C} - \{0\}$ by $\tau_1(z) = z + \frac{\epsilon}{z}$. If $z \in \gamma$ $|\tau_1(z)| \geq d - \frac{\epsilon}{d} > \frac{d}{2}$ and τ is 1-1 on γ . Indeed $\tau(z) = \tau(z')$ implies $z = z'$ or $z' = -\frac{\epsilon}{z}$, but $|zz'| \geq d^2$ for z and $z' \in \gamma$. Set $\Gamma_\epsilon = \{z \in \mathbf{C}, |z| = \frac{\sqrt{\epsilon}}{2}\}$. τ maps Γ_ϵ to Γ'_ϵ the ellipse with vertices at $\pm \left(2\sqrt{\epsilon} + \frac{\sqrt{\epsilon}}{2}\right)$ and $\pm \left(2\sqrt{\epsilon} - \frac{\sqrt{\epsilon}}{2}\right) i$, which lies inside the curve $\tau_1(\gamma)$, since $\frac{5\sqrt{\epsilon}}{2} < \frac{d}{2}$. But a crucial fact is that as a

map from Γ_ϵ to Γ'_ϵ , τ_1 reverses orientation. If f is a distribution on γ which is on γ the limit of a sequence of polynomials, f extends holomorphically inside γ . Set $g = (\tau_1)_* f$ (i.e. $f \circ \tau_1^{-1}$ if f is a function) on $\tau_1(\gamma)$. If f is a limit of polynomials in $z + \frac{\epsilon}{z}$, g extends holomorphically inside $\tau_1(\gamma)$.

Denote by \tilde{f} and \tilde{g} the above holomorphic extensions. By unique analytic continuation, we have $\tilde{g} \circ \tau_1 = \tilde{f}$ on the region of the plane delimited by γ and Γ_ϵ (which is indeed mapped, by τ_1 , inside $\tau_1(\gamma)$).

If \tilde{g} were not constant, by adding a constant we could assume that \tilde{g} does not vanish on Γ'_ϵ but vanishes at some point inside Γ'_ϵ . Since τ_1 reverses orientation (from Γ_ϵ to Γ'_ϵ), the variation of the argument of \tilde{f} along Γ_ϵ would be $2\pi k$ for some $k < 0$, a contradiction.

Remark. — The bound $|\epsilon| \leq \frac{d^2}{25}$ should not be taken seriously. In fact if γ is a circle it is easy to see that the proof above gives the result for every $\epsilon \neq 0$. In case of functions (instead of distributions) one clearly does not need the curve to be smooth.

3.3. The distributional case. — The basic fact to be used is the following : if γ is a curve in a *CR* manifold, which is not “complex tangential” (i.e. $\dot{\gamma}$ does not belong to the span of the Cauchy Riemann vector fields and their conjugates) then the restriction of any *CR* distribution to the curve γ makes sense as a distribution (and in the space of distribution varies smoothly as the curve varies smoothly). This does not require the *CR* structure to be embedded. On the contrary, it comes from a much more general theory, see [11] Proposition 5.2, page 39. To extend the proof given in 3.1 is then easy. To avoid any difficulty on Σ^\pm , one can take advantage of the fact the structure on Σ^+ is embedded, the one on Σ^- is embeddable (by τ). Then any *CR* distribution has local holomorphic extensions to the convex side.

4. Remarks.

4.1. For the next paragraph it will be useful to indicate how to reformulate our claim, to lead obviously to an “invariant” formulation :

Let $N = (0, 1)$. This is a normal vector to S at the point $p, p = 0$, pointing towards the convex side of S . Then $\{\operatorname{Re} z_2 = \alpha\} = \{\operatorname{Re}\langle (z - p) \cdot N \rangle = \alpha\}$. And the conclusion has been that *CR* functions were depending

only on $\langle z \cdot N \rangle$.

4.2. It is not hard to see that the canonical bundle of the structure constructed in 2, cannot have on Σ (or even Σ') a non-zero closed section. For definitions, and interesting related results, see [3]. Indeed such a section could be written $h(z_1, z_2)dz_1 \wedge dz_2$ on Σ^+ and $k(z_1, z_2)dz_1^\# \wedge dz_2$ on Σ^- , where h and k should be CR (for our structure) on Σ^\pm . On $\text{Re } z_2 = \alpha$ one must then have $h(z_1, z_2) = \left(1 - \frac{\epsilon}{z_1^2} \chi(z_2)\right) k(z_1, z_2)$. Let γ be as in 3, and let us use again the notations in the proof of the Lemma. For $z_2 = \alpha$, fixed, the function $z_1 \mapsto h(z_1, \alpha)$ extends holomorphically inside γ , while the function $z_1 \mapsto k(\tau_1^{-1}(z_1), \alpha)$ defined on $\tau_1(\gamma)$ extends holomorphically inside the curve $\tau_1(\gamma)$. Denote these holomorphic extensions, by \tilde{k} and \tilde{h} . By unique analytic continuation one must have :

$$\frac{z_1^2 \tilde{h}(z_1)}{z_1^2 - \epsilon} = \tilde{k} \circ \tau_1(z_1), \text{ for } z_1$$

inside γ , $|z_1| > \frac{\sqrt{\epsilon}}{2}$. The singularities of the left-hand side, at $z_1 = \pm\sqrt{\epsilon}$, are therefore removable. By the Lemma, $\frac{z_1^2 \tilde{h}}{z_1^2 - \epsilon}$ must be constant, which can happen only if \tilde{h} is identically 0. Again, the same must happen for $\text{Re } z_2 = \alpha$, z_2 close to α , on the corresponding curves. And one gets that the canonical bundle has only the zero section, on Σ .

II. EXAMPLES

1. We consider a strictly convex surface S and $\beta > 0$ as in I.2. Let $S_j = \left(\lambda_j, \frac{\beta}{2^j}\right)$ be a sequence of points in S (converging to 0). Denote by N_j the unit normal to S at S_j , pointing towards the convex side. Set $\Gamma_j = \{z \in s, \text{Re}\langle(z - s_j) \cdot N_j\rangle \leq \alpha_j\}$, where the α_j 's are positive numbers, chosen small enough in order that on Γ_j $\frac{\beta}{2^j} - \frac{\beta}{2^{j+2}} < \text{Re } z_2 < \frac{\beta}{2^j} + \frac{\beta}{2^{j+2}}$. The convex hulls of the Γ_j are disjoint. On each Γ_j make a small perturbation of the CR structure induced from \mathbf{C}^2 , following the basic construction with 0 replaced by ζ_j , so that every CR function on Γ_j depends only on $\langle z \cdot N_j \rangle$. One gets a CR structure on S , arbitrarily close to the original one, for which :

PROPOSITION 1. — *If f is any CR distribution defined on a connected neighborhood of 0 in S (for the CR structure just defined), then f must be constant.*

Proof. — Set $\lambda = \frac{\beta}{2^r} + \frac{\beta}{2^{r+2}}$ for $r \geq 1$, r large enough in order that f be defined on $S \cap \{\operatorname{Re} z_2 < \lambda\}$. Notice that our choice of λ makes sure that the hyperplane $\{\operatorname{Re} z_2 = \lambda\}$ does not intersect any Γ_j . Let ω be the open set bounded by S and the hyperplane $\{\operatorname{Re} z_2 = \lambda\}$. Set $\widehat{\Gamma}_j$ to be the convex hull of Γ_j . Due to Lupacoliu's theorem, f extends holomorphically to $\omega - \bigcup_j \widehat{\Gamma}_j$. Indeed one can show that $b\omega \cap \left[\left(\bigcup_j \widehat{\Gamma}_j \right) \cup \{\operatorname{Re} z_2 = \lambda\} \right]$ is polynomially convex, or more easily one can take care of each Γ_j , separately. And our structure is just the usual one off the Γ_j 's. Due to what we have seen before, for infinitely many indices j (therefore for at least two distinct), this holomorphic extension must depend only on $\langle z \cdot N_j \rangle$. So it must be constant. Then, as seen in I.2, f must be constant on $S \cap \{\operatorname{Re} z_2 < \lambda\}$ (by the last part of the claim). If one desires, it is not hard to prove that in fact f must be constant on the whole connected neighborhood on which it is defined, by reasoning as in I.

2. Going back to the construction given in I, for $j = 1, 2, \dots$ set $\alpha_j = \frac{\beta}{2^j}$. For each α_j , we can do a small perturbation of the standard CR structure exactly as described in I (just take $\alpha = \alpha_j$). This perturbation takes place in this region $\left\{ \frac{2\alpha_j}{3} < \operatorname{Re} z_2 < \alpha_j \right\}$. One ends with a CR structure on S , arbitrarily close to the standard one. From the claim in I (this why we introduced Σ'), we get immediately that, for this CR structure :

PROPOSITION 2. — *z_2 is a CR function, and every CR function or distribution, defined in a connected neighborhood of 0 in S , depends only on z_2 .*

3. *Remarks.* — A variant of the construction done in 1 will be presented in III. Somewhat "closer" to the construction in 2, this construction will lead more easily to a CR structure which extends to a complex structure on the concave side.

Every CR structure on a 3-dimensional manifold can be approximated by a real analytic one (since there is no integrability condition in dimension 3), and therefore a locally embeddable one. So we can obtain examples

of non-embeddable CR structure with no non-constant CR distribution arbitrarily close to any given CR structure.

From I.4.2, one sees that in our examples the canonical bundles has no non-zero section in any connected neighborhood of 0.

III. HIGHER DIMENSIONS

1. Let us make some comments on the situation. To carry out our construction in higher dimensions, with extra variables z_3, \dots, z_m , we meet the following problem. We would like to set again $z_1^\# = z_1 + \frac{\epsilon}{z_1} \chi$, where χ is a cut off function which must be 0 when $z_1 = 0$.

And we wish to keep $\bar{\partial}_b z_1^\# = 0$ on $\{\operatorname{Re} z_2 = \alpha\}$. This is clearly impossible if, as in general, χ has to depend effectively on the extra variables z_3, \dots, z_m . Things however can be worked out in case of concavity (in the extra variables). Another way to see an obstruction is the following : if for $z_1 = \alpha$ the trace of S on $\{z_2 = \alpha\}$ were a bounded domain in \mathbf{C}^{n-1} , the matching of the CR structures of Σ^\pm along $\{\operatorname{Re} z_2 = \alpha\}$ would require that the (lower dimensional) CR structures induced on $S \cap \{z_2 = \alpha\}$ coincide. Then Hartogs extension phenomenon would go precisely in the opposite direction to the Lemma given in I, by imposing some matching of the CR functions on Σ^\pm .

2. In the remaining part of this paper we will consider S an hypersurface defined in a neighborhood of 0 in \mathbf{C}^m , $0 \in S$. And we assume that S is defined by

$$\operatorname{Re} z_2 = |z_1|^2 + |z_2|^2 - \sum_{j=3}^n |z_j|^2 + \mathcal{O}(|z|^3).$$

This covers (after holomorphic change of variables) the case of hypersurfaces with non-degenerate Levi form with only one positive eigenvalue. If we fix $\alpha > 0$ (small) the key fact is that, on $S \cap \{z_2 = \alpha\}$, z_1 does not vanish. This allows to adapt *immediately* the construction done in II.2 (treating z_3, \dots, z_m as mere parameters on which χ would not depend) to give an example of a non-embeddable CR structure arbitrarily close to the standard CR structure induced from \mathbf{C}^m on S , with following properties. The functions z_2, \dots, z_m are CR functions. Every CR function f

depends only on (z_2, \dots, z_m) , not on z_1 , and therefore, being CR, satisfies $df \wedge dz_2 \wedge \dots \wedge dz_m = 0$.

To work with several sets of coordinates allows one to strengthen the conclusion. This is what is just sketched in the next paragraph.

3. Let $k \geq 2$. We want to get, for a perturbed CR structure on S , that z_{k+1}, \dots, z_m are CR functions and that, in a neighborhood of 0, any CR function can depend only on z_{k+1}, \dots, z_m (in particular is constant if $k \geq m$). Here, there is a change of notations with respect to the introduction.

All the perturbations which follow will be made small enough to keep the signature of the Levi form, and therefore the smoothness of CR functions. We pick "small" disjoint open neighborhoods V_j of the complex hyperplanes $\{z_2 = 2^{-j}\}$, for j large. Set $S_j = S \cap V_j$ and $S' = S - (\bigcup_j V_j)$. On a neighborhood of S' in S we keep the standard structure for which, in a neighborhood of 0, the Levi form has one positive eigenvalue and $(m-2)$ negative eigenvalues. We assume, as one can, that S' is connected, and locally connected, and that each S_j is connected. On each S_j we modify the structure in such a way that :

(i) Every CR function on S_j satisfies

$$df \wedge dz_2^{(j)} \wedge \dots \wedge dz_k^{(j)} \wedge dz_{k+1} \wedge \dots \wedge dz_m = 0,$$

where each $z_p^{(j)}$ is some linear combination (depending on j) of the coordinates functions z_1, \dots, z_k , very close in the C^∞ topology to z_p ($p = 2, \dots, k$).

(ii) Every CR function f on S_j , which on some non empty open subset of S_j satisfies the equation $df \wedge dz_{k+1} \wedge \dots \wedge dz_m = 0$, satisfies the same equation on S_j .

We assume the $z_2^{(j)}$ to be closed enough to z_2 so that for Ω a fixed neighborhood of 0, in which the construction will be carried, $\{z_2^{(j)} = 2^{-j}\} \cap \Omega \subset V_j$.

To achieve (i) one proceeds as already done several times. One modifies the standard structure in a neighborhood of the complex hyperplane $\{z_2^{(j)} = 2^{-j}\}$, by patching along the real hyperplane $\{\operatorname{Re} z_2^{(j)} = 2^{-j}\}$ the standard structure with the structure defined by prescribing the functions

$$z_1^\# = z_1 + \frac{\epsilon}{z_1} \chi(z_2^{(j)}, z_2^{(j)}, \dots, z_k^{(j)}, z_{k+1}, \dots, z_m$$

to be CR [one can apply Lupacoliu's theorem "slicewise" for $z_3^{(j)}, \dots, z_k^{(j)}, z_{k+1}, \dots, z_m$ fixed and small]. The function χ is a smooth cut off function with small support around 2^{-j} and $\bar{\partial}$ flat on the real line $\{\text{Re } z = 2^{-j}\}$, ϵ is a small positive number.

Let us turn to (ii). For $\delta > 0$, small enough the set $S_j - \{z \in \mathbf{C}^n, \text{Re } z_2^{(j)} = 2^{-j}, |\text{Im } z_2^{(j)}| < \delta\}$ is connected. If the support of χ has been chosen small enough (that for simplicity of the proof we will assume) our CR structure is locally embeddable at each point of this set (by $z \mapsto (\zeta_1, \dots, \zeta_k, z_{k+1}, \dots, z_m)$, for $\zeta_p = z_p$ or $z_p^{(j)}$ $p = 2, \dots, k$ and $\zeta_1 = z_1$ or $z_1^\#$). Then (ii) follows immediately from the fact that a holomorphic function defined on a connected open set cannot vanish on a nonempty open subset of an hypersurface without being identically 0, since on S_j the functions f, z_{k+1}, \dots, z_m are CR functions.

The same argument shows the following. Let \mathcal{O} be an open neighborhood of O in S , assume $\mathcal{O} \cap S'$ to be connected. Let f be a CR function defined on \mathcal{O} . For large j , $S_j \subset \mathcal{O}$. From (i) we have

$$df \wedge dz_2^{(j)} \wedge \dots \wedge dz_k^{(j)} \wedge dz_{k+1} \wedge \dots \wedge dz_m = 0, \text{ on } S_j$$

but, on a neighborhood of S' $z_2^{(j)}, \dots, z_k^{(j)}, z_{k+1}, \dots, z_m$ are CR functions and we then get that

$$df \wedge dz_2^{(j)} \wedge \dots \wedge dz_k^{(j)} \wedge dz_{k+1} \wedge \dots \wedge dz_m = 0, \text{ on } S'.$$

We do not immediately get this on S_ℓ for $\ell \neq j$, since on S_ℓ the $z_p^{(j)}$ are not CR, for our structure.

By using, for different indices j , different linear combinations $z_2^{(j)}, \dots, z_k^{(j)}$ of the functions z_1, z_2, \dots, z_k , (it is crucial that z_1 appears!) we can get that on a neighborhood of $\mathcal{O} \cap S'$: $df \wedge dz_{k+1} \wedge \dots \wedge dz_m = 0$.


Then according to (ii) $df \wedge dz_{k+1} \wedge \dots \wedge dz_m = 0$ on S and this yields the desired conclusion.

5. Remark. — We wish to present some remarks about the possibility to do our construction in such a way that the CR structure extends to one side of S to a complex structure.

The perturbation of the CR structure that we did on S_j leads to a corresponding perturbation on $V_j - \{z_1 = 0\}$. While one keeps the usual complex structure on one side of the real hyperplane $\{\text{Re } z_2^{(j)} =$

2^{-j} }, one imposes on the other side the functions $z_1^\#$ (same definition), $z_2^{(j)}, \dots, z_k^{(j)}, z_{k+1}, \dots, z_m$ to be holomorphic to define the complex structure.

On a neighborhood of the boundary of V_j , this is still (in a fixed neighborhood of 0) the standard structure. Finally, this gives an extension of the CR structure to a complex structure on the side of S towards which $(0, -1, 0, \dots, 0)$ is pointing out (but not on the other side). The following figure, for $(z_3, \dots, z_m) = 0$, may help :



For $\text{Re } z_2 < 0$, the complex structure to be considered is just the usual one.

Added in proof : Hill's conjecture has been recently proved by D. Catlin (communication at the AMS summer Institute at Santa Cruz, July 89).

BIBLIOGRAPHY

- [1] T. AKAHORI, A new approach to the local embedding theorem for *CR* structures for $n \geq 4$, *Memoirs of the AMS* no. 336, Providence, RI 1987.
- [2] D. HILL, What is the notion of a complex manifold with boundary, *Prospect in Algebraic Analysis* [M. Saito 60th birthday vol.].
- [3] H. JACOBOWITZ, The canonical bundle and realizable *CR* hypersurfaces, *Pacific J. Math.*, 127 (1987), 91-101.
- [4] H. JACOBOWITZ, F. TRÈVES, Nonrealizable *CR* structures, *Inventiones Math.*, 66 (1982), 321-249.
- [5] M. KURANISHI, Strongly pseudoconvex *CR* structures over small balls, *Ann. of Math.*, I 115 (1982), 451-500, II 116 (1982), 1-64, III 116 (1982), 249-330.
- [6] G. LUPACCIOLU A theorem on holomorphic extension for *CR* functions, *Pacific J. Math.*, 124 (1986), 177-191.
- [7] L. NIRENBERG, Lectures on linear partial differential equations, *Conference Board of Math. Sc., Regional Conference Series in mathematics* No. 17, AMS, 1973.
- [8] L. NIRENBERG, On a question of Hans Lewy, *Russian Math. Surveys*, 29, (1974), 251-262.
- [9] J.P. ROSAY, E.L. STOUT, Rado's theorem for *CR* functions, to appear in *Proc. AMS*.
- [10] M.C. SHAW, Hypoellipticity of a system of complex vector fields, *Duke Math. J.*, 50 no. 3 (1983), 713-728.
- [11] F. TRÈVES, Approximation and representation of functions and distributions annihilated by a system of complex vector fields, *Ecole polytechnique* (1981).
- [12] F. TRÈVES, Introduction to pseudodifferential and Fourier Integral operators, *Plenum* (1980).
- [13] S. WEBSTER, On the proof of Kuranishi's embedding theorem, (preprint).
- [14] D. CATLIN, A Newlander Nirenberg theorem for manifolds with boundary, *Mich. Math. J.*, 35 (1988), 233-240.

Manuscrit reçu le 17 janvier 1989.

Jean-Pierre ROSAY,
Department of Mathematics
University of Wisconsin
Madison WI 53706 (USA).