

# ANNALES DE L'INSTITUT FOURIER

NICHOLAS BUCHDAHL

## **A Nakai-Moishezon criterion for non-Kähler surfaces**

*Annales de l'institut Fourier*, tome 50, n° 5 (2000), p. 1533-1538

[http://www.numdam.org/item?id=AIF\\_2000\\_\\_50\\_5\\_1533\\_0](http://www.numdam.org/item?id=AIF_2000__50_5_1533_0)

© Annales de l'institut Fourier, 2000, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## A NAKAI-MOISHEZON CRITERION FOR NON-KÄHLER SURFACES

by Nicholas BUCHDAHL

---

### 0. Introduction.

In Corollary 15 of [B], the classical Nakai-Moishezon criterion for a compact complex surface  $X$  was generalised to yield a characterization of the set of classes in  $H_{\mathbb{R}}^{1,1}(X)$  which can be represented by a Kähler form, a result obtained independently by Lamari [L]. Under the assumption that  $b_1(X)$  is even, this result was further generalised in Theorem 16 of [B] to the case of  $\bar{\partial}\partial$ -closed modulo  $\bar{\partial}\partial$ -exact  $(1, 1)$ -forms. The purpose of this paper is to demonstrate that the assumption on  $b_1(X)$  can be dropped entirely. Namely, the following will be proved:

**THEOREM.** — *Let  $X$  be a compact complex surface equipped with a positive  $\bar{\partial}\partial$ -closed  $(1, 1)$ -form  $\omega$  and let  $\varphi$  be a smooth real  $\bar{\partial}\partial$ -closed  $(1, 1)$ -form satisfying  $\int_X \varphi \wedge \varphi > 0$ ,  $\int_X \varphi \wedge \omega > 0$  and  $\int_D \varphi > 0$  for every irreducible effective divisor  $D \subset X$  with  $D \cdot D < 0$ . Then there is a smooth function  $g$  on  $X$  such that  $\varphi + i\bar{\partial}\partial g$  is positive.*

Theorem 16 of [B] differs from this only in that it assumes  $b_1(X)$  is even and that  $\int_D \varphi > 0$  for every effective divisor  $D \subset X$ ; however, this inequality must hold for any effective divisor  $D$  with  $D \cdot D \geq 0$  by Proposition 5 of that paper.

*Acknowledgement.* — This first version of this paper was written during the second half of 1999 when the author was a visitor at l'Institut des Hautes Études Scientifiques. The author wishes to express his deep gratitude to IHÉS for its hospitality.

## 1. Proof of the theorem.

Let  $X$  be a compact complex surface. Since the theorem has already been proved in the case of even first Betti number, it will be assumed henceforth that  $b_1(X)$  is odd. The same notation as in [B] is employed throughout, so  $\Lambda^{p,q}$  denotes the sheaf of germs of smooth  $(p, q)$ -forms on  $X$ , with  $\Lambda^{p,q}(X)$  denoting the global sections. A  $\bar{\partial}\partial$ -closed positive  $(1, 1)$ -form  $\omega \in \Lambda_{\mathbb{R}}^{1,1}(X)$  is chosen once and for all, its existence guaranteed by Gauduchon's theorem [G].

For any  $f \in \Lambda^{1,1}(X)$  there is a function  $g \in \Lambda^{0,0}(X)$ , unique up to the addition of a constant, such that  $\omega \wedge (f + g'')$  is a constant multiple of  $\omega^2$  where  $g'' := i\bar{\partial}\partial g$ . Since  $b_1(X)$  is odd, the proof of Lemma 8 in [B] implies that there is a unique form  $\sigma_0 \in \Lambda_{\mathbb{R}}^{1,1}(X)$  with the properties that it is  $d$ -exact and satisfies  $\omega \wedge \sigma_0 = \omega^2$ . The harmonic representative of a closed  $(1, 1)$ -form  $f$  on  $X$  satisfying  $\omega \wedge f = c\omega^2$  for some constant  $c$  is then  $f - c\sigma_0$ . This form is anti-self-dual with respect to  $\omega$ , a manifestation of the fact that the intersection form on  $H^2(X, \mathbb{R})$  restricted to  $H_{\mathbb{R}}^{1,1}(X)$  is negative definite ([BPV], IV 2.13).

For a holomorphic line bundle  $L$  on  $X$ , there is a unique hermitian metric on  $L$  such that the corresponding hermitian connection has curvature  $f_L$  satisfying  $\omega \wedge f_L = \text{Const} \cdot \omega^2$ . If  $s \in \Gamma(X, \mathcal{O}(L))$  is non-zero and  $E$  is the associated effective divisor  $s^{-1}(0)$ , the equation of currents  $2\pi E = i f - i\bar{\partial}\partial \log |s|^2$  holds by the Poincaré-Lelong theorem ([GH]). Therefore  $\int_E \varphi = \frac{i}{2\pi} \int_X f_L \wedge \varphi$  for any smooth  $\bar{\partial}\partial$ -closed  $(1, 1)$ -form  $\varphi$ . When the divisor  $E$  is given without reference to  $L$ , the notation  $f_E$  will be used to denote  $f_L$  for  $L = \mathcal{O}(E)$ .

A *real divisor* on  $X$  is by definition a finite formal sum of the form  $D = \sum_i \nu_i D_i$  where  $D_i \subset X$  is an irreducible effective divisor on  $X$  and  $\nu_i$  is a real number;  $D$  is *effective* if  $\nu_i \geq 0$  for all  $i$ , in which case the usual notation  $D \geq 0$  is employed; similarly,  $D \geq E$  iff  $D - E \geq 0$ . As for integral divisors, the notation  $f_D$  is used to denote  $\sum_i \nu_i f_{D_i}$ .

The intersection form on  $H^2(X, \mathbb{R})$  is denoted by the dot product symbol. Thus  $E \cdot E$  is the self-intersection number of an effective divisor

$E$  in  $X$ , realised by the integral  $-\frac{1}{4\pi^2} \int_X f_E \wedge f_E$ . The notation extends by  $\mathbb{R}$ -linearity to all real divisors, and is further extended to denote the pairing between  $\bar{\partial}\partial$ -closed  $(1, 1)$ -forms:  $\varphi \cdot \psi := \int_X \varphi \wedge \psi$  for  $\bar{\partial}\partial$ -closed  $\varphi, \psi \in \Lambda_{\mathbb{R}}^{1,1}(X)$ . If  $\psi = if_D$  for some real divisor  $D$ , the notation  $\varphi \cdot D$  may also be used in place of  $\frac{1}{2\pi} \varphi \cdot if_D$ .

LEMMA 1. — *Let  $E \subset X$  be an effective integral divisor such that  $E \cdot E = 0$ . Then for any  $\varepsilon > 0$  there is a smooth function  $g$  such that  $if_E + g'' \geq -\varepsilon\omega$ .*

*Proof.* — If there is no smooth function  $g$  on  $X$  such that  $if_E + g'' + \varepsilon\omega$  is positive in a neighbourhood of  $E$ , the Hahn-Banach Theorem implies the existence of a current  $T$  and a constant  $c$  such that  $T(if_E + \varepsilon\omega + g'') \leq c$  for every smooth function  $g$  and  $T(\psi) > c$  for every smooth 2-form  $\psi$  whose  $(1, 1)$ -component is positive in a neighbourhood of  $E$ .

It follows immediately that  $T$  is a  $(1, 1)$ -current, that  $\bar{\partial}\partial T = 0$ , that  $c$  must be non-positive, that  $T(if_E + \varepsilon\omega) \leq c$ , that  $T(\psi) \geq 0$  for any smooth  $(1, 1)$ -form  $\psi$  which is positive in a neighbourhood of  $E$  and finally that the support of  $T$  must be contained in  $E$ . By Lemma 32 of [HL], it follows that  $T = \sum_i h_i E_i$  where  $h_i$  is a non-negative constant and  $E_1, E_2, \dots$  are the irreducible components of  $E$ . Since  $E \cdot E = 0$  and  $b_1(X)$  is odd,  $[E] = 0$  in  $H^2(X, \mathbb{R})$ . Hence  $E_i \cdot E = 0$  for all  $i$ , and this gives a contradiction since then  $c \geq T(if_E + \varepsilon\omega) = T(\varepsilon\omega) > c$ .

It can therefore be supposed that  $E$  is the zero set of a section  $s$  of a holomorphic line bundle  $L$  which has a hermitian connection whose curvature form  $f$  satisfies  $if > -\varepsilon\omega$  in an open neighbourhood  $U$  of  $E$ . After rescaling  $s$  if necessary, it can be assumed that  $\{x \in X \mid |s(x)| \leq 1\} \subset U$ .

Let  $\chi$  be a smooth convex increasing function on  $\mathbb{R}$  such that  $0 \leq \chi'(t) \leq 1$  for all  $t$ , with  $\chi(t) = t$  for  $t \geq 0$  and with  $\chi(t) = -1$  for  $t \leq -1$ . Then  $i\bar{\partial}\partial(\chi(\log |s|^2)) = \chi'(\log |s|^2) i\bar{\partial}\partial \log |s|^2 + \chi''(\log |s|^2) i\bar{\partial}(\log |s|^2) \wedge \partial(\log |s|^2) \leq \chi'(\log |s|^2) if$ , so  $if - i\bar{\partial}\partial(\chi(\log |s|^2)) \geq (1 - \chi'(\log |s|^2)) if \geq -\varepsilon\omega$ , as required. □

Remark. — The above proof also works in some cases when  $b_1(X)$  is even. For example, if  $E$  is irreducible (with  $E \cdot E = 0$ ), or if every effective divisor on  $X$  has non-negative self-intersection.

LEMMA 2. — *Suppose  $\psi \in \Lambda_{\mathbb{R}}^{1,1}(X)$  satisfies  $\bar{\partial}\partial\psi = 0$ ,  $\psi \cdot \psi = 0$ ,  $\psi \cdot \omega \geq 0$  and  $\psi \cdot D \geq 0$  for every effective divisor  $D \subset X$ . Then for any*

$\varepsilon > 0$  there is a smooth function  $g$  such that  $\psi + g'' \geq -\varepsilon\omega$ .

*Proof.* — By Lemma 7 of [B],  $\psi$  can be approximated arbitrarily closely in  $L^2$  norm by forms of the kind  $p - g''$  where  $p$  is smooth and positive and  $g$  is smooth. Following exactly the same argument as used in the proof of Theorem 11 of [B], a sequence of smooth functions  $g_n$  and smooth positive  $(1, 1)$ -forms  $p_n$  can be found such that  $\|\psi + g_n'' - p_n\|_{L^2(\omega)}$  is converging to 0 and  $g_n$  is converging in  $L^1$  to define an almost-positive closed  $(1, 1)$ -current  $P = g_\infty'' \geq -\psi$ . Applying the same arguments as in the proofs of Theorems 11 and 16 in [B] shows that for any given  $\varepsilon > 0$  there is a real effective divisor  $D_\varepsilon$  and a smooth function  $g_\varepsilon$  such that  $-if_{D_\varepsilon} + g_\varepsilon'' \geq -\psi - \varepsilon\omega$ . The construction of  $D_\varepsilon$  is such that it can be assumed that  $D_{\varepsilon'} \geq D_\varepsilon$  for  $\varepsilon' < \varepsilon$  and the coefficient of an irreducible component common to both  $D_\varepsilon$  and  $D_{\varepsilon'}$  is the same in both.

Now take a sequence of positive numbers  $\varepsilon$  converging monotonically to 0. Since  $\chi_\varepsilon := \varepsilon\omega + \psi - if_{D_\varepsilon} + g_\varepsilon''$  is positive,  $0 \leq \chi_\varepsilon \cdot \chi_\varepsilon = \varepsilon^2 \omega \cdot \omega + 4\pi^2 D_\varepsilon \cdot D_\varepsilon + 2\varepsilon \omega \cdot \psi - 4\pi\varepsilon \omega \cdot D_\varepsilon - 2\pi \psi \cdot D_\varepsilon$ . The hypotheses on  $\psi$  and negativity of the intersection form restricted to  $H_{\mathbb{R}}^{1,1}(X)$  therefore imply that the cohomology classes  $[D_\varepsilon] \in H^2(X, \mathbb{R})$  are uniformly bounded. After passing to a subsequence if necessary, the corresponding sequence of harmonic representatives can be assumed to converge smoothly. Moreover, the inequality  $0 \leq \omega \cdot \chi_\varepsilon = \varepsilon \omega \cdot \omega + \omega \cdot \psi - 2\pi \omega \cdot D_\varepsilon$  implies that the increasing sequence of non-negative numbers  $\{\omega \cdot D_\varepsilon\}$  is bounded above and hence converges. Therefore the sequence of forms  $\{f_{D_\varepsilon}\}$  converges smoothly to a closed  $(1, 1)$ -form  $f_{\mathcal{D}}$  satisfying  $f_{\mathcal{D}} \cdot f_{\mathcal{D}} = 0 = \psi \cdot f_{\mathcal{D}}$  and  $\omega \wedge if_{\mathcal{D}} = c\omega^2$  for some constant  $c \geq 0$ . Since  $[if_{\mathcal{D}}] = 0$  in  $H^2(X, \mathbb{R})$  it follows  $if_{\mathcal{D}} = c\sigma_0$ .

If  $c = 0$ , it follows from the fact that  $\{\omega \cdot D_\varepsilon\}$  is non-negative and increasing that  $\omega \cdot D_\varepsilon = 0$  for all  $\varepsilon$ ; in this case  $D_\varepsilon = 0$  for all  $\varepsilon$  and therefore  $\psi + g_\varepsilon'' \geq -\varepsilon\omega$  as required.

If  $c > 0$ , the identity  $\psi \cdot \sigma_0 = 0$  and Proposition 5 of [B] imply that  $\psi + g''$  is a non-negative multiple of  $\sigma_0$  for some smooth function  $g$ . If there is a non-zero integral effective divisor  $E$  on  $X$  such that  $E \cdot E = 0$ , since  $[\sigma_0] = 0$  in  $H^2(X, \mathbb{R})$  it follows that  $\sigma_0 \cdot E = 0$  and by Proposition 5 of [B] again, that  $\sigma_0$  is a positive multiple of  $if_E$ ; in this case, the desired result follows from Lemma 1. If  $X$  has algebraic dimension 1, it is well-known that  $X$  is an elliptic surface ([BPV], VI 4.1) and therefore such a divisor  $E$  exists.

If  $X$  has algebraic dimension 0, then by [BPV], IV 6.2, there are only

finitely many irreducible curves on  $X$  so that for  $\varepsilon$  sufficiently small, the real divisors  $D_\varepsilon$  are independent of  $\varepsilon$ . Hence  $f_{\mathcal{D}} = f_D$  for some genuine real effective divisor  $D$  on  $X$  satisfying  $D \cdot D = 0$ . By Lemma 4 in §3.5 of Ch. V of [Bou], the symmetric negative semi-definite intersection matrix  $M$  associated with the irreducible components of a connected component of  $D$  has a 1-dimensional kernel, and the entries in a generating vector  $\mathbf{v}$  all have the same sign. Since  $\mathbf{v}$  must be a multiple of a column of the cofactor matrix of  $M$ , after multiplying by a real constant it has positive integer entries. This implies that there is an effective non-zero integral divisor  $E$  on  $X$  with  $E \cdot E = 0$ , so the desired result follows from the previous paragraph.  $\square$

The proof of the main theorem can now be completed. Let  $\varphi \in \Lambda_{\mathbb{R}}^{1,1}(X)$  be a  $\bar{\partial}\partial$ -closed form satisfying the hypotheses of the theorem. By the proof of Theorem 14 of [B], there is a form  $u \in \Lambda^{0,1}(X)$  such that  $\tilde{\varphi} := \varphi + \partial u + \bar{\partial}\bar{u}$  is positive; (the hypothesis that  $b_1(X)$  be even in that theorem is used only in the final sentence of the proof).

By Proposition 5 of [B],  $\tilde{\varphi} \cdot \varphi$  is strictly positive. Let  $t_0$  be the smaller solution of the equation  $(\varphi - t_0\tilde{\varphi}) \cdot (\varphi - t_0\tilde{\varphi}) = 0$ , and set  $\psi := \varphi - t_0\tilde{\varphi}$ . Since  $(\varphi - t\tilde{\varphi}) \cdot (\varphi - t\tilde{\varphi}) > 0$  for  $t$  satisfying  $0 \leq t < t_0$ , the sign of  $\omega \cdot (\varphi - t\tilde{\varphi})$  cannot change for such  $t$  so  $\omega \cdot \psi \geq 0$ . Since  $(\varphi - \tilde{\varphi}) \cdot (\varphi - \tilde{\varphi}) = -2\|\bar{\partial}u\|^2 \leq 0$ , it follows that  $t_0 \leq 1$  and therefore for any effective divisor  $E \subset X$ ,  $\psi \cdot E = (1 - t_0)\varphi \cdot E \geq 0$ .

The form  $\psi$  therefore satisfies the hypotheses of Lemma 2. Applying that lemma, given  $\varepsilon > 0$  there is a smooth function  $g_\varepsilon$  such that  $\psi + g_\varepsilon'' \geq -\varepsilon\omega$ , so if  $\varepsilon$  is chosen so small that  $t_0\tilde{\varphi} - \varepsilon\omega > 0$ , it follows that  $\varphi + g_\varepsilon'' > 0$ , as required.  $\square$

*Remark.* — The methods of this paper show that if  $\varphi \in \Lambda_{\mathbb{R}}^{1,1}(X)$  satisfies the hypotheses of the theorem except for the condition that  $\int_E \varphi$  be positive for every effective  $E \subset X$  with negative self-intersection, there is an effective real divisor  $D$  on  $X$  such that  $\varphi - if_D$  is  $i\bar{\partial}\partial$ -homologous to a positive form.

## BIBLIOGRAPHY

- [BPV] W. BARTH, C. PETERS and A. VAN DE VEN, *Compact Complex Surfaces*, Berlin-Heidelberg-New York, Springer, 1984.

- [Bou] N. BOURBAKI, Groupes et Algèbres de Lie, Ch. 4,5,6, "Éléments de mathématiques" Fasc. XXXIV, Paris, Hermann, 1968.
- [B] N. P. BUCHDAHL, On compact Kähler surfaces, *Ann. Inst. Fourier*, 49-1 (1999), 287–302.
- [G] P. GAUDUCHON, Le théorème de l'excentricité nulle, *C. R. Acad. Sci. Paris*, 285 (1977), 387–390.
- [GH] P. A. GRIFFITHS and J. HARRIS, *Principles of Algebraic Geometry*, New York, Wiley, 1978.
- [HL] R. HARVEY and H. B. LAWSON, An intrinsic characterisation of Kähler manifolds, *Invent. Math.* 74 (1983), 169–198.
- [L] A. LAMARI, Le cône kählérien d'une surface, *J. Math. Pures Appl.*, 78 (1999), 249–263.

Manuscrit reçu le 15 novembre 1999,  
révisé le 16 mars 2000,  
accepté le 27 avril 2000.

Nicholas BUCHDAHL,  
University of Adelaide  
Department of Pure Mathematics  
Adelaide 5005 (Australia).  
nbuchdah@maths.adelaide.edu.au