



# ANNALES

DE

# L'INSTITUT FOURIER

Kazuya KATO & Henrik RUSSELL

**Albanese varieties with modulus and Hodge theory**

Tome 62, n° 2 (2012), p. 783-806.

[http://aif.cedram.org/item?id=AIF\\_2012\\_\\_62\\_2\\_783\\_0](http://aif.cedram.org/item?id=AIF_2012__62_2_783_0)

© Association des Annales de l'institut Fourier, 2012, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du  
Centre de diffusion des revues académiques de mathématiques  
<http://www.cedram.org/>*

# ALBANESE VARIETIES WITH MODULUS AND HODGE THEORY

by Kazuya KATO & Henrik RUSSELL (\*)

---

ABSTRACT. — Let  $X$  be a proper smooth variety over a field  $k$  of characteristic 0 and  $Y$  an effective divisor on  $X$  with multiplicity. We introduce a generalized Albanese variety  $\text{Alb}(X, Y)$  of  $X$  of modulus  $Y$ , as higher dimensional analogue of the generalized Jacobian with modulus of Rosenlicht-Serre. Our construction is algebraic. For  $k = \mathbb{C}$  we give a Hodge theoretic description.

RÉSUMÉ. — Soient  $X$  une variété propre et lisse sur un corps  $k$  de caractéristique 0 et  $Y$  un diviseur effectif avec multiplicité sur  $X$ . Nous introduisons une variété d'Albanese généralisée  $\text{Alb}(X, Y)$  de  $X$ , de module  $Y$ , comme analogue en dimension supérieure de la jacobienne généralisée avec module de Rosenlicht-Serre. Notre construction est algébrique. Si  $k = \mathbb{C}$ , nous donnons une description en termes de théorie de Hodge.

## 1. Introduction

**1.1.** Let  $X$  be a proper smooth variety over a field  $k$  of characteristic 0, and let  $\text{Alb}(X)$  be the Albanese variety of  $X$ . In the work [10], the second author constructed generalized Albanese varieties  $\text{Alb}_{\mathcal{F}}(X)$ , which are commutative connected algebraic groups over  $k$  with surjective homomorphisms  $\text{Alb}_{\mathcal{F}}(X) \rightarrow \text{Alb}(X)$  (see Section 5 for a review). If  $Y$  is an effective divisor on  $X$ , a special case of  $\text{Alb}_{\mathcal{F}}(X)$  becomes the generalized Albanese variety  $\text{Alb}(X, Y)$  of  $X$  of modulus  $Y$  (*cf.*, Section 5). This is a higher dimensional analogue of the generalized Jacobian variety with modulus of Rosenlicht-Serre. Note that the divisor  $Y$  can have multiplicity, and so the algebraic group  $\text{Alb}(X, Y)$  can have an additive part.

---

*Keywords:* generalized Albanese variety, modulus of a rational map, generalized mixed Hodge structure.

*Math. classification:* 14L10, 14C30, 14F42.

(\*) The second author was supported by the DFG.

Assume now  $k = \mathbb{C}$ . The purpose of this paper is to give Hodge theoretic presentations (Theorem 1.1) of  $\text{Alb}(X, Y)$ .

The case when  $Y$  has no multiplicity was studied in the work [3] of Barbieri-Viale and Srinivas. A Hodge theoretic presentation of a generalized Albanese variety in the case without modulus but allowing singularities on  $X$  was given in the work [6] of Esnault, Srinivas and Viehweg.

**1.2.** First we review the curve case. Let  $X$  be a proper smooth curve over  $\mathbb{C}$  and let  $Y$  be an effective divisor on  $X$ . In this case, the Albanese variety  $\text{Alb}(X, Y)$  of  $X$  relative to  $Y$  coincides with the generalized Jacobian variety  $J(X, Y)$  of  $X$  relative to  $Y$ . In the following, we will write the complex analytic space associated to  $X$  simply by  $X$ , and the sheaf of holomorphic functions on it by  $\mathcal{O}_X$ . Let  $I = \text{Ker}(\mathcal{O}_X \rightarrow \mathcal{O}_Y)$  be the ideal of  $\mathcal{O}_X$  which defines  $Y$ . The cohomology below is for the topology of the analytic space  $X$  (not for Zariski topology).

The generalized Jacobian variety  $J(X, Y)$  is the kernel of the degree map  $\text{Pic}(X, Y) \rightarrow \mathbb{Z}$  where  $\text{Pic}(X, Y) = H^1(X, \text{Ker}(\mathcal{O}_X^\times \rightarrow \mathcal{O}_Y^\times))$ . Let  $j : X - Y \rightarrow X$  be the inclusion map and let  $j_*\mathbb{Z}(1)$  be the 0-extension of the constant sheaf  $\mathbb{Z}(1)$  of  $X - Y$  to  $X$ . (For  $r \in \mathbb{Z}$ ,  $\mathbb{Z}(r)$  denotes  $\mathbb{Z}(2\pi i)^r$  as usual.) Then we have an exact sequence

$$0 \rightarrow j_*\mathbb{Z}(1) \rightarrow I \xrightarrow{\text{exp}} \text{Ker}(\mathcal{O}_X^\times \rightarrow \mathcal{O}_Y^\times) \rightarrow 0$$

and hence we have an isomorphism

$$(1.1) \quad \text{Pic}(X, Y) \cong H^2(X, [j_*\mathbb{Z}(1) \rightarrow I]).$$

Here in the complex  $[j_*\mathbb{Z}(1) \rightarrow I]$ ,  $j_*\mathbb{Z}(1)$  is put in degree 0.

We have another presentation of  $J(X, Y)$  given in (2) below. Let  $I_1$  be the ideal of  $\mathcal{O}_X$  which defines the reduced part of  $Y$  and let  $J = II_1^{-1} \subset \mathcal{O}_X$ . Note that the composition of the two inclusion maps of complexes

$$[I \xrightarrow{d} J\Omega_X^1] \rightarrow [I \xrightarrow{d} \Omega_X^1] \rightarrow [I_1 \xrightarrow{d} \Omega_X^1]$$

is a quasi-isomorphism. Hence we have an isomorphism in the derived category

$$[I \xrightarrow{d} \Omega_X^1] \cong [I_1 \xrightarrow{d} \Omega_X^1] \oplus (\Omega_X^1/J\Omega_X^1)[-1].$$

Since  $j_*\mathbb{C} \rightarrow [I_1 \xrightarrow{d} \Omega_X^1]$  is a quasi-isomorphism, we have an exact sequence

$$(1.2) \quad H^0(X, \Omega_X^1) \rightarrow H_c^1(X - Y, \mathbb{C}/\mathbb{Z}(1)) \oplus H^0(X, \Omega_X^1/J\Omega_X^1) \rightarrow J(X, Y) \rightarrow 0.$$

(Here  $H_c$  is the cohomology with compact supports.)

**1.3.** Now let  $X$  be a proper smooth variety over  $\mathbb{C}$  of dimension  $n$  and let  $Y$  be an effective divisor on  $X$ .

Again in the following theorem, cohomology groups are for the topology of the complex analytic spaces, and the notation  $\mathcal{O}$  and  $\Omega$  stand for analytic sheaves.

Let  $I$  be the ideal of  $\mathcal{O}_X$  which defines  $Y$ , let  $I_1$  be the ideal of  $\mathcal{O}_X$  which defines the reduced part of  $Y$ , and let  $J = II_1^{-1} \subset \mathcal{O}_X$ .

**THEOREM 1.1.**

(1) We have an exact sequence

$$0 \longrightarrow \text{Alb}(X, Y) \longrightarrow H^{2n}(X, \mathcal{D}_{X,Y}(n)) \xrightarrow{\text{deg}} \mathbb{Z} \longrightarrow 0,$$

where for  $r \in \mathbb{Z}$ ,  $\mathcal{D}_{X,Y}(r)$  denotes the kernel of the surjective homomorphism of complexes  $\mathcal{D}_X(r) \rightarrow \mathcal{D}_Y(r)$  with  $\mathcal{D}_X(r)$  the Deligne complex

$$[\mathbb{Z}(r) \rightarrow \mathcal{O}_X \xrightarrow{d} \Omega_X^1 \xrightarrow{d} \dots \xrightarrow{d} \Omega_X^{r-1}]$$

and  $\mathcal{D}_Y(r)$  the similar complex

$$[\mathbb{Z}(r)_Y \rightarrow \mathcal{O}_Y \xrightarrow{d} \Omega_Y^1 \xrightarrow{d} \dots \xrightarrow{d} \Omega_Y^{r-1}].$$

(2) We have an exact sequence

$$\begin{aligned} H^{n-1}(X, \Omega_X^n) \longrightarrow H_c^{2n-1}(X - Y, \mathbb{C}/\mathbb{Z}(n)) \oplus H^{n-1}(X, \Omega_X^n/J\Omega_X^n) \\ \longrightarrow \text{Alb}(X, Y) \longrightarrow 0. \end{aligned}$$

Note that the case  $n = 1$  of Theorem 1.1 (1) (resp. (2)) becomes the presentation of  $J(X, Y)$  given by (1) (resp. (2)) in No. 1.2.

*Remark 1.2.* — We give some remarks on this theorem.

(a) The case  $Y = 0$  of Theorem 1.1 (1) is nothing but the well known exact sequence

$$(1.3) \quad 0 \longrightarrow \text{Alb}(X) \longrightarrow H^{2n}(X, \mathcal{D}_X(n)) \xrightarrow{\text{deg}} \mathbb{Z} \longrightarrow 0$$

by using the Deligne cohomology  $H^{2n}(X, \mathcal{D}_X(n))$ . (Usually the Deligne cohomology  $H^m(X, \mathcal{D}_X(r))$  is denoted by  $H_D^m(X, \mathbb{Z}(r))$ .)

The case  $Y = 0$  of Theorem 1.1 (2) is nothing but the usual presentation

$$(1.4) \quad \text{Alb}(X) \cong H_{\mathbb{Z}} \setminus H_{\mathbb{C}} / F^0 H_{\mathbb{C}}$$

of the Albanese variety  $\text{Alb}(X)$  of  $X$ , where  $(H_{\mathbb{Z}}, H_{\mathbb{C}}, F^\bullet)$  is the following Hodge structure of weight  $-1$ .  $H_{\mathbb{Z}} = H^{2n-1}(X, \mathbb{Z}(n))/(\text{torsion part})$ ,  $H_{\mathbb{C}} =$

$\mathbb{C} \otimes_{\mathbb{Z}} H_{\mathbb{Z}} = H^{2n-1}(X, \Omega_X^\bullet)$ , and  $F^\bullet$  is the Hodge filtration on  $H_{\mathbb{C}}$  defined as

$$F^{-1} = H_{\mathbb{C}}, \quad F^0 = H^{n-1}(X, \Omega_X^n), \quad F^1 = 0.$$

(b) Recall that the presentations (3) and (4) of  $\text{Alb}(X)$  are related as follows. Consider the exact sequence of complexes  $0 \rightarrow \Omega_X^{\leq n-1}[-1] \rightarrow \mathcal{D}_X(n) \rightarrow \mathbb{Z}(n) \rightarrow 0$ , where  $\Omega_X^{\leq n-1}$  denotes the part of degree  $\leq n-1$  of the de Rham complex  $\Omega_X^\bullet$ , which is actually a quotient complex of  $\Omega_X^\bullet$ . By taking the cohomology associated to this exact sequence, we have an exact sequence

$$H^{2n-1}(X, \mathbb{Z}(n)) \longrightarrow H^{2n-1}(X, \Omega_X^{\leq n-1}) \longrightarrow H_D^{2n}(X, \mathbb{Z}(n)) \xrightarrow{\text{deg}} \mathbb{Z} \longrightarrow 0.$$

Since

$$\begin{aligned} H^{2n-1}(X, \Omega_X^{\leq n-1}) &\cong H^{2n-1}(X, \Omega_X^\bullet) / H^{n-1}(X, \Omega_X^n) \\ &\cong H^{2n-1}(X, \mathbb{C}) / H^{n-1}(X, \Omega_X^n), \end{aligned}$$

the exact sequence (4) is equivalent to (3).

(c) (1) and (2) of Theorem 1.1 are related similarly. Let  $S$  be the subcomplex of the de Rham complex  $\Omega_X^\bullet$  of  $X$  defined by  $S^p = \text{Ker}(\Omega_X^p \rightarrow \Omega_Y^p)$  for  $0 \leq p \leq n-1$  and  $S^n = \Omega_X^n$ . Then Theorem 1.1 (1) is equivalent to

$$\text{Alb}(X, Y) \cong H_{\mathbb{Z}} \setminus H^{2n-1}(X, S) / H^{n-1}(X, \Omega_X^n)$$

where  $H_{\mathbb{Z}} = H_c^{2n-1}(X - Y, \mathbb{Z}(n)) / (\text{torsion part})$ . As shown in § 6, we have a commutative diagram with an isomorphism in the lower row

$$\begin{array}{ccc} H^{n-1}(X, \Omega_X^n) & = & H^{n-1}(X, \Omega_X^n) \\ \downarrow & & \downarrow \\ H^{2n-1}(X, S) & \cong & H_c^{2n-1}(X - Y, \mathbb{C}) \oplus H^{n-1}(X, \Omega_X^n / J\Omega_X^n). \end{array}$$

Thus (1) and (2) of Theorem 1.1 are deduced from each other.

**1.4.** As mentioned above, Theorem 1.1 shows that  $\text{Alb}(X, Y)$  is expressed as  $H_{\mathbb{Z}} \setminus H_V / F^0$  where:

$$\begin{aligned} H_{\mathbb{Z}} &= H_c^{2n-1}(X - Y, \mathbb{Z}(n)) / (\text{torsion part}), \\ H_V &= H_{\mathbb{C}} \oplus H^{n-1}(X, \Omega_X^n / J\Omega_X^n) \cong H^{2n-1}(X, S) \end{aligned}$$

$$(H_{\mathbb{C}} = \mathbb{C} \otimes H_{\mathbb{Z}} \text{ and } S \text{ is as in 1.5 (d)),$$

$F^\bullet$  is the decreasing filtration on  $H_V$  given by

$$F^{-1} = H_V, \quad F^0 = H^{n-1}(X, \Omega_X^n), \quad F^1 = 0.$$

Note that  $H_V$  can be different from  $H_{\mathbb{C}}$  here, and so  $(H_{\mathbb{Z}}, H_V, F^\bullet)$  here need not be a Hodge structure. It is some kind of “mixed Hodge structure with additive part”. This object  $(H_{\mathbb{Z}}, H_V, F^\bullet)$  with a weight filtration, which we will denote by  $H^{2n-1}(X, Y_-)(n)$  in Section 6, belongs to a category  $\mathcal{H}$  introduced in Section 2 which contains the category of mixed Hodge structures but is larger than that. In the proof of Theorem 1.1, it is essential to consider such an object. This category  $\mathcal{H}$  is related to the category of “enriched Hodge structures” of Bloch-Srinivas [4] and to the category of “formal Hodge structures” of Barbieri-Viale [1]. However, the relations between these three categories are not trivial, see 4.6 and [2, 4.2]. Our definition of  $\mathcal{H}$  aims to stick close to the classical language of Hodge structures and to express duality in a simplest possible way. In the proof of Theorem 1.1, we use a Hodge theoretic description of the category of “1-motives with additive parts” over  $\mathbb{C}$  in terms of  $\mathcal{H}$ . This description is similar to the result of Barbieri-Viale in [1].

**1.5.** The theory of generalized Albanese varieties in characteristic  $p > 0$  is given in [11], basing on duality theory of “1-motives with unipotent parts”.

In characteristic  $p > 0$ , *syntomic cohomology* is an analogue of *Deligne cohomology*. We expect that we can have presentations of the  $p$ -adic completion of  $\text{Alb}(X, Y)(k)$  ( $k$  is the base field), which is similar to Theorem 1.1, by using crystalline cohomology theory and syntomic cohomology theory.

We are thankful to Professor H el ene Esnault for advice.

## 2. Mixed Hodge structures with additive parts

**2.1.** For a proper smooth variety  $X$  over  $\mathbb{C}$  of dimension  $n$  and for an effective divisor  $Y$  on  $X$ , we will have in Section 6 certain structures  $H^1(X, Y_+)$  and  $H^{2n-1}(X, Y_-)$  which are kinds of “mixed Hodge structures with additive parts”. (These structures for the case when  $X$  is a curve are explained in Example 2.1 below.) The authors imagine that there is a nice definition of the category of “mixed Hodge structures with additive parts”, which contains these  $H^1(X, Y_+)$  and  $H^{2n-1}(X, Y_-)$  as objects, but can not define it. Instead, we define a category  $\mathcal{H}$  containing these objects, which may be a very simple approximation of such a nice category.

**2.2.** The category  $\mathcal{H}$ . An object of  $\mathcal{H}$  is by definition a tuple  $H = (H_{\mathbb{Z}}, H_V, W_{\bullet}H_{\mathbb{Q}}, W_{\bullet}H_V, F^{\bullet}H_V, a, b)$ , where  $H_{\mathbb{Z}}$  is a finitely generated  $\mathbb{Z}$ -module,  $H_V$  is a finite dimensional  $\mathbb{C}$ -vector space,  $W_{\bullet}H_{\mathbb{Q}}$  is an increasing

filtration on  $H_{\mathbb{Q}} := \mathbb{Q} \otimes H_{\mathbb{Z}}$  (called weight filtration),  $W_{\bullet}H_V$  is an increasing filtration on  $H_V$  (called weight filtration),  $F^{\bullet}$  is a decreasing filtration on  $H_V$  (called Hodge filtration),  $a$  is a  $\mathbb{C}$ -linear map  $H_{\mathbb{C}} := \mathbb{C} \otimes H_{\mathbb{Z}} \rightarrow H_V$  which sends  $W_w H_{\mathbb{C}} := \mathbb{C} \otimes_{\mathbb{Q}} W_w H_{\mathbb{Q}}$  into  $W_w H_V$  for any  $w \in \mathbb{Z}$ , and  $b$  is a  $\mathbb{C}$ -linear map  $H_V \rightarrow H_{\mathbb{C}}$  which sends  $W_w H_V$  into  $W_w H_{\mathbb{C}}$  for any  $w \in \mathbb{Z}$  such that  $b \circ a$  is the identity map of  $H_{\mathbb{C}}$ . We sometimes denote an object  $H$  of  $\mathcal{H}$  simply by  $(H_{\mathbb{Z}}, H_V)$ .

A morphism  $f : H \rightarrow H'$  in  $\mathcal{H}$  is a pair of homomorphisms  $(f_{\mathbb{Z}}, f_V)$ , where  $f_{\mathbb{Z}} : H_{\mathbb{Z}} \rightarrow H'_{\mathbb{Z}}$  is compatible with the weight filtrations and  $f_V : H_V \rightarrow H'_V$  is compatible with weight filtrations and Hodge filtrations, which is compatible with the maps  $a, b$  and  $a', b'$ .

The category of mixed Hodge structures is naturally embedded into  $\mathcal{H}$  as a full subcategory, by putting  $H_V = H_{\mathbb{C}}$ .

Similarly as for mixed Hodge structures we can give  $\underline{\text{Hom}}(H, H')$  the structure of an object of  $\mathcal{H}$  for  $H, H' \in \text{Ob}(\mathcal{H})$ . We call  $\underline{\text{Hom}}(H, \mathbb{Z})$  the object dual to  $H$ . The full subcategory of  $\mathcal{H}$  consisting of all objects  $H$  such that  $H_{\mathbb{Z}}$  are torsion free is clearly self-dual.

We will say that a sequence  $H' \rightarrow H \rightarrow H''$  in  $\mathcal{H}$  is exact, if and only if the following sequences are all exact:

$$\begin{aligned} H'_{\mathbb{Z}} \rightarrow H_{\mathbb{Z}} \rightarrow H''_{\mathbb{Z}}, & & H'_V \rightarrow H_V \rightarrow H''_V, \\ W_w H'_{\mathbb{Q}} \rightarrow W_w H_{\mathbb{Q}} \rightarrow W_w H''_{\mathbb{Q}}, & & W_w H'_V \rightarrow W_w H_V \rightarrow W_w H''_V, \\ & & F^p H'_V \rightarrow F^p H_V \rightarrow F^p H''_V, \end{aligned}$$

for all  $w, p \in \mathbb{Z}$ .

See No. 4.6 for the relation of this category  $\mathcal{H}$  to the category of enriched Hodge structures of Bloch-Srinivas [4] and to the category of formal Hodge structures of Barbieri-Viale [1].

*Example 2.1.* — Let  $X$  be a proper smooth curve over  $\mathbb{C}$  and let  $Y$  be an effective divisor on  $X$ . Let  $I$  be the ideal of  $\mathcal{O}_X$  which defines  $Y$ , let  $I_1$  be the ideal of  $\mathcal{O}_X$  which defines the reduced part of  $Y$ , and let  $J = II_1^{-1} \subset \mathcal{O}_X$ .

We define objects  $H^1(X, Y_+)$  and  $H^1(X, Y_-)$  of  $\mathcal{H}$ .

First, we define  $H = H^1(X, Y_+)$ . Let

$$H_{\mathbb{Z}} = H^1(X - Y, \mathbb{Z}), \quad H_V = H^1(X, [\mathcal{O}_X \xrightarrow{d} I^{-1}\Omega^1_X]).$$

The map  $a : H_{\mathbb{C}} \rightarrow H_V$  is

$$H^1(X - Y, \mathbb{C}) \cong H^1(X, [\mathcal{O}_X \rightarrow I_1^{-1}\Omega^1_X]) \longrightarrow H^1(X, [\mathcal{O}_X \rightarrow I^{-1}\Omega^1_X]).$$

The map  $b : H_V \rightarrow H_{\mathbb{C}}$  is the composition

$$\begin{aligned}
 H^1(X, [\mathcal{O}_X \rightarrow I^{-1}\Omega_X^1]) &\longrightarrow H^1(X, [J^{-1} \rightarrow I^{-1}\Omega_X^1]) \\
 &\xleftarrow{\cong} H^1(X, [\mathcal{O}_X \rightarrow I_1^{-1}\Omega_X]) \cong H^1(X - Y, \mathbb{C}).
 \end{aligned}$$

The weight filtrations and the Hodge filtration are given by

$$\begin{aligned}
 W_2H_{\mathbb{Q}} &= H_{\mathbb{Q}}, & W_1H_{\mathbb{Q}} &= H^1(X, \mathbb{Q}), & W_0H_{\mathbb{Q}} &= 0, \\
 W_2H_V &= H_V, & W_1H_V &= H^1(X, \mathbb{C}), & W_0H_V &= 0,
 \end{aligned}$$

where  $H^1(X, \mathbb{C})$  is embedded in  $H_V$  via  $a$ , and

$$F^0H_V = H_V, \quad F^1H_V = H^1(X, \mathbb{C}), \quad F^2H_{\mathbb{C}} = 0.$$

Next, we define  $H = H^1(X, Y_-)$ . Let

$$H_{\mathbb{Z}} = H_c^1(X - Y, \mathbb{Z}), \quad H_V = H^1(X, [I \xrightarrow{d} \Omega_X^1]).$$

The map  $a : H_{\mathbb{C}} \rightarrow H_V$  is the composition

$$\begin{aligned}
 H_c^1(X - Y, \mathbb{C}) &\cong H^1(X, [I_1 \rightarrow \Omega_X^1]) \\
 &\xleftarrow{\cong} H^1(X, [I \rightarrow J\Omega_X^1]) \longrightarrow H^1(X, [I \rightarrow \Omega_X^1]).
 \end{aligned}$$

The map  $b : H_V \rightarrow H_{\mathbb{C}}$  is

$$H^1(X, [I \rightarrow \Omega_X^1]) \longrightarrow H^1(X, [I_1 \rightarrow \Omega_X^1]) \cong H_c^1(X - Y, \mathbb{C}).$$

The weight filtrations and the Hodge filtration are given by

$$\begin{aligned}
 W_1H_{\mathbb{Q}} &= H_{\mathbb{Q}}, & W_0H_{\mathbb{Q}} &= \text{Ker}(H_{\mathbb{Q}} \rightarrow H^1(X, \mathbb{Q})), & W_{-1}H_{\mathbb{Q}} &= 0, \\
 W_1H_V &= H_V, & W_0H_V &= \text{Ker}(H_V \rightarrow H^1(X, \mathbb{C})), & W_{-1}H_V &= 0,
 \end{aligned}$$

where  $H^1(X, \mathbb{C})$  is regarded as quotient of  $H_V$  via  $b$ , and

$$F^0H_V = H_V, \quad F^1H_V = \text{Ker}(H_V \rightarrow H^1(X, \mathcal{O}_X)), \quad F^2H_{\mathbb{C}} = 0.$$

Then we have exact sequences in  $\mathcal{H}$

$$\begin{aligned}
 0 &\longrightarrow H^1(X) \longrightarrow H^1(X, Y_+) \longrightarrow H^0(Y)(-1) \longrightarrow \mathbb{Z}(-1) \longrightarrow 0, \\
 0 &\longrightarrow \mathbb{Z} \longrightarrow H^0(Y) \longrightarrow H^1(X, Y_-) \longrightarrow H^1(X) \longrightarrow 0.
 \end{aligned}$$

Here for  $r \in \mathbb{Z}$ ,  $\mathbb{Z}(r)$  is the usual Hodge structure  $\mathbb{Z}(r)$  regarded as an object of  $\mathcal{H}$ .  $H^1(X)$  is also the usual Hodge structure of weight 1 associated to the first cohomology of  $X$ , regarded as an object of  $\mathcal{H}$ . Finally the object  $H^0(Y)$  of  $\mathcal{H}$  is defined as below, and  $H^0(Y)(-1)$  is the  $-1$  Tate twist.

The definition of  $H = H^0(Y)$  is as follows.  $H_{\mathbb{Z}} = H^0(Y, \mathbb{Z}) = \bigoplus_{y \in Y} \mathbb{Z}$ .  $H_V = H^0(Y, \mathcal{O}_Y)$ .  $a$  is the canonical map  $H^0(Y, \mathbb{C}) \rightarrow H^0(Y, \mathcal{O}_Y)$ .  $b$  is the canonical map  $H^0(Y, \mathcal{O}_Y) \rightarrow H^0(Y, \mathbb{C})$  given by  $\mathcal{O}_Y \rightarrow \mathbb{C}$  which is



$\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{Y,y}/m_y = \mathbb{C}$  at each  $y \in Y$  ( $m_y$  denotes the maximal ideal of  $\mathcal{O}_{Y,y}$ ). The weight filtration and the Hodge filtration are given by

$$\begin{aligned} W_0H &= H, & W_{-1}H &= 0, \\ F^0H_V &= H_V, & F^1H_V &= 0. \end{aligned}$$

Note that  $H_{\mathbb{C}} \rightarrow H_V$  can be like  $\mathbb{C} \rightarrow \mathbb{C}[T]/(T^n)$ , and need not be an isomorphism.

The evident self-duality  $\underline{\text{Hom}}(\_, \mathbb{Z})$  for torsion free objects in  $\mathcal{H}$  induces

$$H^1(X, Y_-) \cong \underline{\text{Hom}}(H^1(X, Y_+), \mathbb{Z})(-1).$$

### 3. 1-motives with additive parts

In [9], Laumon formulated the notion of a “1-motive with additive part” over a field of characteristic 0. We give a short review assuming that the base field is algebraically closed for simplicity.

Fix an algebraically closed field  $k$  of characteristic 0.

**3.1.** Let  $\mathcal{A}b/k$  be the category of sheaves of abelian groups on the fppf-site of the category of affine schemes over  $k$ . Let  $\mathcal{C}^{[-1,0]}(\mathcal{A}b/k)$  be the abelian category of complexes in  $\mathcal{A}b/k$  concentrated in degrees  $-1$  and  $0$ .

A 1-motive with additive part over  $k$  is an object of  $\mathcal{C}^{[-1,0]}(\mathcal{A}b/k)$  of the form  $[\mathcal{F} \rightarrow G]$ , where  $G$  is a commutative connected algebraic group over  $k$  and  $\mathcal{F} \cong \mathbb{Z}^t \oplus (\widehat{\mathbb{G}}_a)^s$  for some  $t$  and  $s$ . (cf., [9, Def. (5.1.1)].) Here  $\mathbb{Z}$  is regarded as a constant sheaf and  $\widehat{\mathbb{G}}_a$  denotes the formal completion of the additive group  $\mathbb{G}_a$  at 0. Recall that for any commutative ring  $R$ ,  $\widehat{\mathbb{G}}_a(R)$  is the subgroup of the additive group  $R$  consisting of all nilpotent elements. We have  $\mathcal{F} = \mathcal{F}_{\text{ét}} \oplus \mathcal{F}_{\text{inf}}$ , where  $\mathcal{F}_{\text{ét}}$  is the étale part of  $\mathcal{F}$  which corresponds to  $\mathbb{Z}^t$  in the above isomorphism and  $\mathcal{F}_{\text{inf}}$  is the infinitesimal part of  $\mathcal{F}$  which corresponds to  $(\widehat{\mathbb{G}}_a)^s$ .

We denote the category of 1-motives with additive parts over  $k$  by  $\mathcal{M}_1$ .

**3.2.** The category  $\mathcal{M}_1$  admits a notion of duality (called “Cartier duality”). Let  $[\mathcal{F} \rightarrow G]$  be a 1-motive with additive part over  $k$ . Then we have the “Cartier dual”  $[\mathcal{F}' \rightarrow G']$  of  $[\mathcal{F} \rightarrow G]$  which is an object of  $\mathcal{M}_1$  obtained as follows. Let  $0 \rightarrow L \rightarrow G \rightarrow A \rightarrow 0$  be the canonical decomposition of  $G$  as an extension of an abelian variety  $A$  by a commutative connected affine algebraic group  $L$ . Note that  $L \cong (\mathbb{G}_m)^t \oplus (\mathbb{G}_a)^s$  for some  $t$  and  $s$ . We have

$$\mathcal{F}' = \underline{\text{Hom}}_{\mathcal{A}b/k}(L, \mathbb{G}_m), \quad G' = \underline{\text{Ext}}^1_{\mathcal{C}^{[-1,0]}(\mathcal{A}b/k)}([\mathcal{F} \rightarrow A], \mathbb{G}_m)$$

and the homomorphism  $\mathcal{F}' \rightarrow G'$  is the connecting homomorphism

$$\underline{\mathrm{Hom}}_{\mathcal{A}b/k}(L, \mathbb{G}_m) \longrightarrow \underline{\mathrm{Ext}}^1_{\mathcal{C}^{[-1,0]}(\mathcal{A}b/k)}([\mathcal{F} \rightarrow A], \mathbb{G}_m)$$

associated to the short exact sequence  $0 \rightarrow L \rightarrow [\mathcal{F} \rightarrow G] \rightarrow [\mathcal{F} \rightarrow A] \rightarrow 0$  in  $\mathcal{C}^{[-1,0]}(\mathcal{A}b/k)$ . Since

$$\underline{\mathrm{Hom}}_{\mathcal{A}b/k}(\mathbb{G}_m, \mathbb{G}_m) \cong \mathbb{Z}, \quad \underline{\mathrm{Hom}}_{\mathcal{A}b/k}(\mathbb{G}_a, \mathbb{G}_m) \cong \widehat{\mathbb{G}}_a,$$

we have  $\mathcal{F}' \simeq \mathbb{Z}^t \oplus (\widehat{\mathbb{G}}_a)^s$  for some  $t$  and  $s$ . We have an exact sequence

$$0 \longrightarrow \underline{\mathrm{Hom}}_{\mathcal{A}b/k}(\mathcal{F}, \mathbb{G}_m) \longrightarrow \underline{\mathrm{Ext}}^1_{\mathcal{C}^{[-1,0]}(\mathcal{A}b/k)}([\mathcal{F} \rightarrow A], \mathbb{G}_m) \longrightarrow \underline{\mathrm{Ext}}^1_{\mathcal{A}b/k}(A, \mathbb{G}_m) \longrightarrow 0,$$

$\underline{\mathrm{Ext}}^1_{\mathcal{A}b/k}(A, \mathbb{G}_m)$  is the dual abelian variety of  $A$ , and since

$$\underline{\mathrm{Hom}}_{\mathcal{A}b/k}(\mathbb{Z}, \mathbb{G}_m) \cong \mathbb{G}_m, \quad \underline{\mathrm{Hom}}_{\mathcal{A}b/k}(\widehat{\mathbb{G}}_a, \mathbb{G}_m) \cong \mathbb{G}_a,$$

$\underline{\mathrm{Hom}}_{\mathcal{A}b/k}(\mathcal{F}, \mathbb{G}_m) \cong (\mathbb{G}_m)^t \oplus (\mathbb{G}_a)^s$  for some  $t$  and  $s$ . Hence  $G'$  is a commutative connected algebraic group over  $k$ . Thus  $[\mathcal{F}' \rightarrow G']$  is a 1-motive with additive part. The Cartier dual of  $[\mathcal{F}' \rightarrow G']$  is canonically isomorphic to  $[\mathcal{F} \rightarrow G]$ .

See [9, Section 5] for details or [10, Section 1] for another review.

**3.3.** Let  $\mathcal{M}_{1,\{-1,-2\}}$  be the full subcategory of  $\mathcal{M}_1$  consisting of all objects  $[\mathcal{F} \rightarrow G]$  such that  $\mathcal{F} = 0$ .

Let  $\mathcal{M}_{1,\{0,-1\}}$  be the full subcategory of  $\mathcal{M}_1$  consisting of all objects  $[\mathcal{F} \rightarrow G]$  such that  $G$  is an abelian variety.

Then the self-duality of  $\mathcal{M}_1$  in No. 3.2 induces an anti-equivalence between the categories  $\mathcal{M}_{1,\{-1,-2\}}$  and  $\mathcal{M}_{1,\{0,-1\}}$ .

### 4. Equivalences of categories

In [1], Barbieri-Viale constructed a Hodge theoretic category and proved that in the case when the base field is  $\mathbb{C}$ , the category  $\mathcal{M}_1$  is equivalent to his Hodge theoretic category. Here we reformulate his equivalence in the style which is convenient for us, by using the category  $\mathcal{H}$  from Section 2.

**4.1.** The category  $\mathcal{H}_1$ . An object of  $\mathcal{H}_1$  is an object  $H$  of  $\mathcal{H}$  endowed with a splitting of the weight filtration on  $\mathrm{Ker}(H_V \rightarrow H_{\mathbb{C}})$  satisfying the following conditions (i)–(iv).

- (i)  $H_{\mathbb{Z}}$  is torsion free,  $F^{-1}H_V = H_V, F^1H_V = 0, W_0H = H, W_{-3}H = 0$ .

(ii)  $\text{gr}_{-1}^W H$  is a polarizable Hodge structure of weight  $-1$ . That is,  $\text{gr}_{-1}^W H_{\mathbb{C}} = \text{gr}_{-1}^W H_V$  and  $\text{gr}_{-1}^W H_{\mathbb{Z}}$  with the Hodge filtration on  $\text{gr}_{-1}^W H_{\mathbb{C}}$  is a polarizable Hodge structure of weight  $-1$ .

(iii)  $F^0 \text{gr}_0^W H_V = \text{gr}_0^W H_V$ .

(iv)  $F^0 W_{-2} H_V = 0$ .

Morphisms of  $\mathcal{H}_1$  are the evident ones.

The category  $\mathcal{H}_1$  is self-dual by the functor  $\underline{\text{Hom}}(\ , \mathbb{Z})(1)$ .

**4.2.** For a subset  $\Delta$  of  $\{0, -1, -2\}$ , let  $\mathcal{H}_{1, \Delta}$  be the full subcategory of  $\mathcal{H}_1$  consisting of all objects  $H$  such that  $\text{gr}_w^W H = 0$  unless  $w \in \Delta$ .

The categories  $\mathcal{H}_{1, \{-1, -2\}}$  and  $\mathcal{H}_{1, \{0, -1\}}$  are important for us. These categories are in fact defined as full subcategories of  $\mathcal{H}$  without reference to the splitting of the weight filtration on  $\text{Ker}(H_V \rightarrow H_{\mathbb{C}})$ , for the weight filtrations on  $\text{Ker}(H_V \rightarrow H_{\mathbb{C}})$  of objects of these categories are pure.

Thus  $\mathcal{H}_{1, \{-1, -2\}}$  is the full subcategory of  $\mathcal{H}$  consisting of all objects  $H$  satisfying the following conditions (i)–(iii).

(i)  $H_{\mathbb{Z}}$  is torsion free,  $F^{-1} H_V = H_V$ ,  $F^1 H_V = 0$ ,  $W_{-1} H = H$ ,  $W_{-3} H = 0$ .

(ii)  $\text{gr}_{-1}^W H$  is a polarizable Hodge structure of weight  $-1$ .

(iii)  $F^0 W_{-2} H_V = 0$ .

For example, the Tate twist  $H^1(X, Y_-)(1)$  of the object  $H^1(X, Y_-)$  of  $\mathcal{H}$  in Example 2.1 belongs to  $\mathcal{H}_{1, \{-1, -2\}}$ .

Similarly,  $\mathcal{H}_{1, \{0, -1\}}$  is the full subcategory of  $\mathcal{H}$  consisting of all objects  $H$  satisfying the following conditions (i)–(iii).

(i)  $H_{\mathbb{Z}}$  is torsion free,  $F^{-1} H_V = H_V$ ,  $F^1 H_V = 0$ ,  $W_0 H = H$ ,  $W_{-2} H = 0$ .

(ii)  $\text{gr}_{-1}^W H$  is a polarizable Hodge structure of weight  $-1$ .

(iii)  $F^0 \text{gr}_0^W H_V = \text{gr}_0^W H_V$ .

For example, the Tate twist  $H^1(X, Y_+)(1)$  of the object  $H^1(X, Y_+)$  of  $\mathcal{H}$  in Example 2.1 belongs to  $\mathcal{H}_{1, \{0, -1\}}$ .

The self-duality  $\underline{\text{Hom}}(\ , \mathbb{Z})(1)$  of  $\mathcal{H}_1$  induces an anti-equivalence between the categories  $\mathcal{H}_{1, \{-1, -2\}}$  and  $\mathcal{H}_{1, \{0, -1\}}$ .

**THEOREM 4.1.** — (*This is an analogue of the equivalence of categories proved by Barbieri-Viale in [1].*) We have an equivalence of categories  $\mathcal{H}_1 \simeq \mathcal{M}_1$  which is compatible with dualities, and which induces the equivalences

$$\mathcal{H}_{1, \{-1, 0\}} \simeq \mathcal{M}_{1, \{-1, 0\}}, \quad \mathcal{H}_{1, \{-2, -1\}} \simeq \mathcal{M}_{1, \{-2, -1\}}.$$

The equivalence  $\mathcal{H}_1 \simeq \mathcal{M}_1$  is described in No.s 4.3 and 4.4 below.

**4.3.** First we define the functor  $\mathcal{H}_1 \rightarrow \mathcal{M}_1$ .

Let  $H$  be an object of  $\mathcal{H}_1$ . The corresponding object  $[\mathcal{F} \rightarrow G]$  of  $\mathcal{M}_1$  is as follows.

$$\begin{aligned}
 G &= W_{-1}H_{\mathbb{Z}} \setminus W_{-1}H_V / F^0W_{-1}H_V, \\
 \mathcal{F}_{\text{ét}} &= \text{gr}_0^W(H_{\mathbb{Z}}), \\
 \mathcal{F}_{\text{inf}} &= \text{the formal completion of } \text{Ker}(\text{gr}_0^W(H_V) \rightarrow \text{gr}_0^W(H_{\mathbb{C}})).
 \end{aligned}$$

Here  $\mathcal{F}_{\text{ét}}$  is the étale part of  $\mathcal{F}$  and  $\mathcal{F}_{\text{inf}}$  is the infinitesimal part of  $\mathcal{F}$ . The homomorphism  $\mathcal{F} = \mathcal{F}_{\text{ét}} \oplus \mathcal{F}_{\text{inf}} \rightarrow G$  is given as follows.

The part  $\mathcal{F}_{\text{ét}} \rightarrow G$ : Let  $x \in \mathcal{F}_{\text{ét}} = \text{gr}_0^W H_{\mathbb{Z}}$ . Since the sequence  $0 \rightarrow W_{-1}H_{\mathbb{Z}} \rightarrow H_{\mathbb{Z}} \rightarrow \text{gr}_0^W H_{\mathbb{Z}} \rightarrow 0$  is exact, we can lift  $x$  to an element  $y$  of  $H_{\mathbb{Z}}$  and this lifting is unique modulo  $W_{-1}H_{\mathbb{Z}}$ . Since the sequence  $0 \rightarrow F^0W_{-1}H_V \rightarrow F^0H_V \rightarrow F^0 \text{gr}_0^W H_V \rightarrow 0$  is exact, we can lift  $x$  to an element  $z$  of  $F^0H_V$  and this lifting is unique modulo  $F^0W_{-1}H_V$ . Note that  $y - z \in W_{-1}H_V$ . We have a well-defined homomorphism

$$\mathcal{F}_{\text{ét}} = \text{gr}_0^W H_{\mathbb{Z}} \longrightarrow W_{-1}H_{\mathbb{Z}} \setminus W_{-1}H_V / F^0W_{-1}H_V = G ; \quad x \longmapsto y - z.$$

The part  $\mathcal{F}_{\text{inf}} \rightarrow G$ : Identify  $\text{Hom}(\mathcal{F}_{\text{inf}}, G)$  with  $\text{Hom}_{\mathbb{C}}(\text{Lie}(\mathcal{F}_{\text{inf}}), \text{Lie}(G))$ . We give the corresponding homomorphism  $\text{Lie}(\mathcal{F}_{\text{inf}}) = \text{Ker}(\text{gr}_0^W(H_V) \rightarrow \text{gr}_0^W(H_{\mathbb{C}})) \rightarrow \text{Lie}(G) = W_{-1}H_V / F^0W_{-1}H_V$ . Let  $x \in \text{Ker}(\text{gr}_0^W(H_V) \rightarrow \text{gr}_0^W(H_{\mathbb{C}}))$ . The given splitting of the weight filtration on  $\text{Ker}(H_V \rightarrow H_{\mathbb{C}})$  sends  $x$  to an element  $y$  of  $\text{Ker}(H_V \rightarrow H_{\mathbb{C}})$ . Since the sequence  $0 \rightarrow F^0W_{-1}H_V \rightarrow F^0H_V \rightarrow F^0 \text{gr}_0^W H_V \rightarrow 0$  is exact, we can lift  $x$  to an element  $z$  of  $F^0H_V$  and this lifting is unique modulo  $F^0W_{-1}H_V$ . Note that  $y - z \in W_{-1}H_V$ . We have a well-defined homomorphism

$$\text{Ker}(\text{gr}_0^W H_V \rightarrow \text{gr}_0^W H_{\mathbb{C}}) \longrightarrow W_{-1}H_V / F^0W_{-1}H_V = \text{Lie}(G) ; \quad x \longmapsto y - z.$$

**4.4.** We give the functor  $\mathcal{M}_1 \rightarrow \mathcal{H}_1$ .

Let  $[\mathcal{F} \rightarrow G]$  be an object of  $\mathcal{M}_1$ . The corresponding object  $H$  of  $\mathcal{H}_1$  is as follows. Let  $0 \rightarrow L \rightarrow G \rightarrow A \rightarrow 0$  be the exact sequence of commutative algebraic groups where  $A$  is an abelian variety and  $L$  is affine. Let  $\mathcal{F}_{\text{ét}}$  be the étale part of  $\mathcal{F}$  and let  $\mathcal{F}_{\text{inf}}$  be the infinitesimal part of  $\mathcal{F}$ .

First,  $H_{\mathbb{Z}}$  is the fiber product of  $\mathcal{F}_{\text{ét}} \rightarrow G \leftarrow \text{Lie}(G)$ , where  $\text{Lie}(G) \rightarrow G$  is the exponential map, so we have a commutative diagram of exact sequences

$$\begin{array}{ccccccc}
 0 & \rightarrow & H_1(G, \mathbb{Z}) & \rightarrow & H_{\mathbb{Z}} & \rightarrow & \mathcal{F}_{\text{ét}} \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \rightarrow & H_1(G, \mathbb{Z}) & \rightarrow & \text{Lie}(G) & \rightarrow & G \rightarrow 0.
 \end{array}$$

The weight filtration on  $H_{\mathbb{Z}}$  is given as follows.

$$\begin{aligned} W_0 H_{\mathbb{Z}} &= H_{\mathbb{Z}}, \\ W_{-1} H_{\mathbb{Z}} &= H_1(G, \mathbb{Z}), \\ W_{-2} H_{\mathbb{Z}} &= H_1(L, \mathbb{Z}) = \text{Ker} (H_1(G, \mathbb{Z}) \rightarrow H_1(A, \mathbb{Z})), \\ W_{-3} H_{\mathbb{Z}} &= 0. \end{aligned}$$

Next,

$$H_V = H_{\mathbb{C}} \oplus \text{Lie}(L_a) \oplus \text{Lie}(\mathcal{F}_{\text{inf}})$$

where  $L_a$  is the additive part of  $L$ . The weight filtration on  $H_V$  is as follows.

$$\begin{aligned} W_0 H_V &= H_V, \\ W_{-1} H_V &= H_1(G, \mathbb{C}) \oplus \text{Lie}(L_a), \\ W_{-2} H_V &= H_1(L, \mathbb{C}) \oplus \text{Lie}(L_a), \\ W_{-3} H_V &= 0. \end{aligned}$$

The splitting of the weight filtration on  $\text{Ker}(H_V \rightarrow H_{\mathbb{C}}) = \text{Lie}(L_a) \oplus \text{Lie}(\mathcal{F}_{\text{inf}})$  is by definition this direct decomposition.

The Hodge filtration on  $H_V$  is given as follows.

$$\begin{aligned} F^{-1} H_V &= H_V, \\ F^1 H_V &= 0, \\ F^0 H_V &= \text{Ker} (H_V \rightarrow \text{Lie}(G)) \end{aligned}$$

where  $H_V \rightarrow \text{Lie}(G)$  is defined as follows. The part  $H_{\mathbb{C}} \rightarrow \text{Lie}(G)$  of it is the  $\mathbb{C}$ -linear map induced by the canonical map  $H_{\mathbb{Z}} \rightarrow \text{Lie}(G)$ . The part  $\text{Lie}(L_a) \rightarrow \text{Lie}(G)$  of it is the inclusion map. The part  $\text{Lie}(\mathcal{F}_{\text{inf}}) \rightarrow \text{Lie}(G)$  of it is the homomorphism induced by  $\mathcal{F}_{\text{inf}} \rightarrow G$ . We have hence  $H_V/F^0 H_V \cong \text{Lie}(G)$ .

It is easy to see that this functor  $\mathcal{M}_1 \rightarrow \mathcal{H}_1$  is quasi-inverse to the functor  $\mathcal{H}_1 \rightarrow \mathcal{M}_1$  in No. 4.3.

**4.5.** The induced functor  $\mathcal{H}_{1, \{-1, -2\}} \xrightarrow{\cong} \mathcal{M}_{1, \{-1, -2\}}$  is especially simple. It is given by

$$H \longmapsto [0 \rightarrow H_{\mathbb{Z}} \setminus H_V / F^0 H_V].$$

**4.6.** For those who are familiar with formal Hodge structures from [1] we explain the relation between  $\mathcal{H}_1$  and the category  $\text{FHS}_1^{\text{fr}}$  of torsion free formal Hodge structures of level  $\leq 1$ , see [1, Def. 1.1.2]. (This No. is not used in the rest of the paper.)

The categories  $\mathcal{H}_1$  and  $\text{FHS}_1^{\text{fr}}$  are equivalent. The functor  $\mathcal{H}_1 \rightarrow \text{FHS}_1^{\text{fr}}$  is given by  $(H_{\mathbb{Z}}, H_V) \mapsto (\mathcal{F}, V)$ , where  $(\mathcal{F}, V)$  is the following object of  $\text{FHS}_1^{\text{fr}}$ .

$$\begin{aligned} \mathcal{F} &= \mathcal{F}_{\text{ét}} \oplus \mathcal{F}_{\text{inf}}, \\ \mathcal{F}_{\text{ét}} &= H_{\mathbb{Z}}, \\ \mathcal{F}_{\text{inf}} &= \text{formal completion of } \text{Ker} \left( \text{gr}_0^W(H_V) \rightarrow \text{gr}_0^W(H_{\mathbb{C}}) \right), \end{aligned}$$

$$\begin{aligned} V &= W_{-1}H_V / W_{-1}F^0H_V \\ &\supseteq V^1 = W_{-2}H_V \\ &\supseteq V^0 = \text{Ker}(W_{-2}H_V \rightarrow W_{-2}H_{\mathbb{C}}), \end{aligned}$$

$v: \mathcal{F} \rightarrow V$  is def. by  $\begin{cases} v|_{\mathcal{F}_{\text{ét}}} = a|_{H_{\mathbb{Z}}} \pmod{F^0H_V} \text{ (we have } V = H_V/F^0H_V), \\ v|_{\mathcal{F}_{\text{inf}}} \text{ is the map } \mathcal{F}_{\text{inf}} \subset \text{Lie}(\mathcal{F}_{\text{inf}}) \rightarrow \text{Lie}(G) \text{ as in No. 4.3,} \end{cases}$

$H_{\mathbb{C}}/F^0H_{\mathbb{C}} \xrightarrow{\cong} V/V^0$  is the map induced by  $a$ .

The functor  $\text{FHS}_1^{\text{fr}} \rightarrow \mathcal{H}_1$  is given by  $(\mathcal{F}, V) \mapsto (H_{\mathbb{Z}}, H_V)$ , where  $(H_{\mathbb{Z}}, H_V)$  is the following object of  $\mathcal{H}_1$ .

$$\begin{aligned} H_{\mathbb{Z}} &= \mathcal{F}_{\text{ét}}, \\ H_V &= H_{\mathbb{C}} \oplus \text{Lie}(\mathcal{F}_{\text{inf}}) \oplus V^0, \\ W_0H_V &= H_V, \\ F^{-1}H_V &= H_V, & W_{-1}H_V &= W_{-1}H_{\mathbb{C}} \oplus V^0, \\ F^0H_V &= \text{Ker}(H_V \rightarrow V), & W_{-2}H_V &= W_{-2}H_{\mathbb{C}} \oplus V^0, \\ F^1H_V &= 0, & W_{-3}H_V &= 0, \end{aligned}$$

where  $H_V \rightarrow V$  is the map given by  $(v|_{\mathcal{F}_{\text{ét}}} \otimes \mathbb{C}, \text{Lie}(v|_{\mathcal{F}_{\text{inf}}}), V^0 \hookrightarrow V)$ .

These functors are quasi-inverse to each other and yield an equivalence of categories  $\mathcal{H}_1 \simeq \text{FHS}_1^{\text{fr}}$ . The relation between  $\text{FHS}_1$  and the category  $\text{EHS}_1$  of enriched Hodge structures of level  $\leq 1$  from [4] is given in [2, 4.2] by explicit functors. Composition yields an explicit functor  $\text{EHS}_1^{\text{fr}} \rightarrow \mathcal{H}_1$  (left to the reader). The category  $\text{EHS}_1^{\text{fr}}$  of torsion free enriched Hodge structures of level  $\leq 1$  is equivalent to a subcategory of  $\text{FHS}_1^{\text{fr}}$  resp.  $\mathcal{H}_1$ , see [2, Prop. 4.2.3].

### 5. Generalized Albanese varieties

Let  $k$  be an algebraically closed field of characteristic 0 and let  $X$  be a proper smooth algebraic variety over  $k$  of dimension  $n$ . We review generalized Albanese varieties  $\text{Alb}_{\mathcal{F}}(X)$  defined in [10]<sup>(1)</sup>. For an effective divisor  $Y$  on  $X$ , the generalized Albanese variety  $\text{Alb}(X, Y)$  of modulus  $Y$  is a special case of  $\text{Alb}_{\mathcal{F}}(X)$ .

The Albanese variety  $\text{Alb}(X)$  is defined by a universal mapping property for morphisms from  $X$  to abelian varieties. Similarly, the generalized Albanese variety  $\text{Alb}(X, Y)$  of modulus  $Y$  is characterized by a universal property for morphisms from  $X - Y$  into commutative algebraic groups with “modulus”  $\leq Y$ . See Proposition 5.1.

**5.1.** Let  $\underline{\text{Div}}_X$  be the sheaf of abelian groups on  $\mathcal{A}b/k$  defined as follows. For any commutative ring  $R$  over  $k$ ,  $\underline{\text{Div}}_X(R)$  is the group of all Cartier divisors on  $X \otimes_k R$  generated locally on  $\text{Spec}(R)$  by effective Cartier divisors which are flat over  $R$ . Let  $\underline{\text{Pic}}_X$  be the Picard functor, and let  $\underline{\text{Pic}}_X^0 \subset \underline{\text{Pic}}_X$  be the Picard variety of  $X$ . We have the class map  $\underline{\text{Div}}_X \rightarrow \underline{\text{Pic}}_X$ . Let  $\underline{\text{Div}}_X^0 \subset \underline{\text{Div}}_X$  be the inverse image of  $\underline{\text{Pic}}_X^0$ .

**5.2.** Let  $\Lambda$  be the set of all subgroup sheaves  $\mathcal{F}$  of  $\underline{\text{Div}}_X^0$  such that  $\mathcal{F} \cong \mathbb{Z}^t \oplus (\widehat{\mathbb{G}}_a)^s$  for some  $t$  and  $s$ . For  $\mathcal{F} \in \Lambda$ , we have an object  $[\mathcal{F} \rightarrow \underline{\text{Pic}}_X^0]$  of  $\mathcal{M}_{1, \{0, -1\}}$ . The generalized Albanese variety  $\text{Alb}_{\mathcal{F}}(X)$  is defined in [10] to be the Cartier dual of  $[\mathcal{F} \rightarrow \underline{\text{Pic}}_X^0]$ . It is an object of  $\mathcal{M}_{1, \{-1, -2\}}$  and hence is a commutative connected algebraic group over  $k$ .

If  $\mathcal{F}, \mathcal{F}' \in \Lambda$  and  $\mathcal{F} \subset \mathcal{F}'$ , we have a canonical surjective homomorphism  $\text{Alb}_{\mathcal{F}'}(X) \rightarrow \text{Alb}_{\mathcal{F}}(X)$ . In the case  $\mathcal{F} = 0$ ,  $\text{Alb}_{\mathcal{F}}(X) = \text{Alb}(X)$ .

**5.3.** Let  $Y$  be an effective divisor of  $X$ . Then the generalized Albanese variety with modulus  $Y$  is defined as  $\text{Alb}_{\mathcal{F}}(X)$  where  $\mathcal{F} = \mathcal{F}_{X, Y} \in \Lambda$  is defined as follows. The étale part  $\mathcal{F}_{\text{ét}}$  of  $\mathcal{F}$  is the subgroup of  $\underline{\text{Div}}_X^0(k)$  consisting of all divisors whose support is contained in the support of  $Y$ . The infinitesimal part  $\mathcal{F}_{\text{inf}}$  of  $\mathcal{F}$  is as follows. Let  $I$  be the ideal of  $\mathcal{O}_X$  (though the notation  $\mathcal{O}_X$  is often used in this paper for the sheaf of analytic functions,  $\mathcal{O}_X$  here stands for the usual algebraic object on the Zariski site) defining  $Y$ , let  $I_1$  be the ideal of  $\mathcal{O}_X$  which defines the reduced part of  $Y$ , and let  $J = II_1^{-1} \subset \mathcal{O}_X$ . Then  $\mathcal{F}_{\text{inf}}$  is the formal completion  $\widehat{\mathbb{G}}_a \otimes_k H^0(X, J^{-1}/\mathcal{O}_X)$  of the finite dimensional  $k$ -vector space  $H^0(X, J^{-1}/\mathcal{O}_X)$ ,

<sup>(1)</sup> In [10],  $X$  was assumed to be projective. This assumption was used only for singular  $X$ , which is not our concern here. The construction of the  $\text{Alb}_{\mathcal{F}}(X)$  is valid in the same way for proper  $X$ .

which is embedded in  $\underline{\text{Div}}_X^0$  by the exponential map

$$\exp : \widehat{\mathbb{G}}_a \otimes_k H^0(X, J^{-1}/\mathcal{O}_X) \longrightarrow \underline{\text{Div}}_X^0.$$

If  $Y'$  is an effective divisor on  $X$  such that  $Y' \geq Y$ , then  $\mathcal{F}_{X,Y'} \supset \mathcal{F}_{X,Y}$  and hence we have a canonical surjective homomorphism  $\text{Alb}(X, Y') \rightarrow \text{Alb}(X, Y)$ . In the case  $Y = 0$ ,  $\text{Alb}(X, Y) = \text{Alb}(X)$ .

In the case when  $X$  is a curve,  $\text{Alb}(X, Y)$  coincides with the generalized Jacobian variety  $J(X, Y)$  of  $X$  with modulus  $Y$  as is explained in [10, Exm. 2.34].

**5.4.** As in [10], for  $\mathcal{F} \in \Lambda$  we have a rational map

$$\alpha_{\mathcal{F}} : X \longrightarrow \text{Alb}_{\mathcal{F}}(X)$$

which is canonically defined up to translation by a  $k$ -rational point of  $\text{Alb}_{\mathcal{F}}(X)$ . If  $\mathcal{F}' \in \Lambda$  and  $\mathcal{F} \subset \mathcal{F}'$ , then  $\alpha_{\mathcal{F}}$  and  $\alpha_{\mathcal{F}'}$  are compatible via the canonical surjection  $\text{Alb}_{\mathcal{F}'}(X) \rightarrow \text{Alb}_{\mathcal{F}}(X)$ .

For an effective divisor  $Y$  on  $X$ , we denote the rational map  $\alpha_{\mathcal{F}_{X,Y}}$  simply by  $\alpha_{X,Y}$ . In Proposition 5.1 (2) below, we give a universal property of  $\alpha_{X,Y} : X \rightarrow \text{Alb}(X, Y)$  concerning rational maps from  $X$  to commutative algebraic groups. This property follows from a general universal property of  $\alpha_{\mathcal{F}} : X \rightarrow \text{Alb}_{\mathcal{F}}(X)$  obtained in [10], as is shown in No. 5.6 below.

**5.5.** Let  $G$  be a commutative connected algebraic group over  $k$  and let  $\varphi : X \rightarrow G$  be a rational map. We define an effective divisor  $\text{mod}(\varphi)$  on  $X$  which we call the modulus of  $\varphi$ .

We treat  $X$  as a scheme. This divisor  $\text{mod}(\varphi)$  is written in the form  $\sum_v \text{mod}_v(\varphi) v$ , where  $v$  ranges over all points of  $X$  of codimension one and  $\text{mod}_v(\varphi)$  is a non-negative integer defined as follows.

Let  $0 \rightarrow L \rightarrow G \rightarrow A \rightarrow 0$  be the canonical decomposition of  $G$  and take an isomorphism

$$(1) \quad L_a \cong (\mathbb{G}_a)^s$$

where  $L_a$  is the additive part of  $L$ .

Let  $K$  be the function field of  $X$ , and regard  $\varphi$  as an element of  $G(K)$ . Since the local ring  $\mathcal{O}_{X,v}$  of  $X$  at  $v$  is a discrete valuation ring and since  $A$  is proper, we have  $A(\mathcal{O}_{X,v}) = A(K)$ . By the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \rightarrow & L(\mathcal{O}_{X,v}) & \rightarrow & G(\mathcal{O}_{X,v}) & \rightarrow & A(\mathcal{O}_{X,v}) & \rightarrow & 0 \\ & & \cap & & \cap & & \parallel & & \\ 0 & \rightarrow & L(K) & \rightarrow & G(K) & \rightarrow & A(K) & \rightarrow & 0, \end{array}$$

we have  $G(K) = L(K)G(\mathcal{O}_{X,v})$ . Write  $\varphi \in G(K)$  as



(2)  $\varphi = lg$  with  $l \in L(K)$  and  $g \in G(O_{X,v})$ .

Let  $(l_j)_{1 \leq j \leq s}$  be the image of  $l$  in  $(\mathbb{G}_a)^s(K)$ .

If  $\varphi$  belongs to  $G(O_{X,v})$ , we define  $\text{mod}_v(\varphi) = 0$ . Assume that  $\varphi$  does not belong to  $G(O_{X,v})$ . Then we define

$$\text{mod}_v(\varphi) = 1 + \max(\{-\text{ord}_v(l_j) \mid 1 \leq j \leq s\} \cup \{0\}).$$

This integer  $\text{mod}_v(\varphi)$  is independent of the choice of the isomorphism (1) and of the choice of the presentation (2) of  $\varphi$ .

For example, if  $G = \mathbb{G}_m$ ,  $\text{mod}_v(\varphi)$  is 0 if the element  $\varphi$  of  $G(K) = K^\times$  belongs to  $\mathcal{O}_{X,v}^\times$ , and is 1 otherwise. If  $G = \mathbb{G}_a$ ,  $\text{mod}_v(\varphi)$  is 0 if the element  $\varphi$  of  $G(K) = K$  belongs to  $\mathcal{O}_{X,v}$ , and is  $m + 1$  if  $\varphi$  has a pole of order  $m \geq 1$  at  $v$ .

PROPOSITION 5.1. — *Let  $G$  be a commutative connected algebraic group over  $k$  and let  $\varphi : X \rightarrow G$  be a rational map.*

- (1) *For a dense open set  $U$  of  $X$ ,  $\varphi$  induces a morphism  $U \rightarrow G$  (not only a rational map) if and only if the support of  $\text{mod}(\varphi)$  does not meet  $U$ .*
- (2) *Let  $Y$  be an effective divisor on  $X$ . Then the following two conditions (i) and (ii) are equivalent.*
  - (i) *There is a homomorphism  $h : \text{Alb}(X, Y) \rightarrow G$  such that  $\varphi$  coincides with  $h \circ \alpha_{X,Y}$  modulo a translation by  $G(k)$ .*
  - (ii)  *$\text{mod}(\varphi) \leq Y$ .*

Furthermore, if these equivalent conditions are satisfied, such homomorphism  $h$  is unique.

It is easy to prove (1). The proof of (2) is given in No. 5.7 below after we review results on  $\text{Alb}_{\mathcal{F}}(X)$  from [10].

**5.6.** We review a general universal property of  $\text{Alb}_{\mathcal{F}}(X)$  proved in [10] concerning rational maps from  $X$  into commutative algebraic groups.

Let  $\varphi : X \rightarrow G$  be a rational map into a commutative connected algebraic group  $G$ , and let  $L$  be the canonical connected affine subgroup such that the quotient  $G/L$  is an abelian variety. One observes that  $\varphi$  induces a natural transformation  $\tau_\varphi : L^\vee \rightarrow \text{Div}_X^0$  (see [10, Section 2.2]), where  $L^\vee = \underline{\text{Hom}}_{\mathcal{A}b/k}(L, \mathbb{G}_m)$  is the Cartier dual of  $L$ . It is shown in [10, Section 2.3] that if  $\mathcal{F} \in \Lambda$ , there is a rational map  $\alpha_{\mathcal{F}} : X \rightarrow \text{Alb}_{\mathcal{F}}(X)$  for which the corresponding homomorphism  $\tau_{\alpha_{\mathcal{F}}} : \mathcal{F} \rightarrow \text{Div}_X^0$  coincides with the inclusion map, and such rational map  $\alpha_{\mathcal{F}}$  is unique up to translation by a  $k$ -rational point of  $\text{Alb}_{\mathcal{F}}(X)$ . For a rational map  $\varphi : X \rightarrow G$  into a commutative connected algebraic group  $G$  and for  $\mathcal{F} \in \Lambda$ , there is a homomorphism

$h : \text{Alb}_{\mathcal{F}}(X) \rightarrow G$  such that  $f$  coincides with  $h \circ \alpha_{\mathcal{F}}$  up to translation by an element of  $G(k)$  if and only if the image of the homomorphism  $\tau_{\varphi} : L^{\vee} \rightarrow \underline{\text{Div}}_X^0$  is contained in  $\mathcal{F}$ . Furthermore, if such  $h$  exists, it is unique.

Moreover, any rational map  $\varphi : X \rightarrow G$  into a commutative connected algebraic group  $G$  coincides with  $h \circ \alpha_{\mathcal{F}}$  up to translation by an element of  $G(k)$  for some  $\mathcal{F} \in \Lambda$  and for some homomorphism  $h : \text{Alb}_{\mathcal{F}}(X) \rightarrow G$ . This is because there is always some  $\mathcal{F} \in \Lambda$  which contains the image of  $L^{\vee} \rightarrow \underline{\text{Div}}_X^0$ .

**5.7.** We prove Proposition 5.1. By No. 5.6 we find that condition (i) of Proposition 5.1 (2) is equivalent to

(i') The image of  $\tau_{\varphi}$  is contained in  $\mathcal{F}_{X,Y}$ .

Write

$$Y = \sum_v e_v v$$

where  $v$  ranges over all points of  $X$  of codimension one and  $e_v \in \mathbb{N}$ . Condition (ii) of Proposition 5.1 (2) is expressed as

(ii')  $\text{mod}_v(\varphi) \leq e_v$  for all points  $v$  of codimension one in  $X$ .

Fix an isomorphism  $L \cong (\mathbb{G}_m)^t \times (\mathbb{G}_a)^s$ . For each point  $v$  of  $X$  of codimension one, take a presentation  $\varphi = lg$  as in (2) in No. 5.5, let  $(l'_{v,j})_{1 \leq j \leq t}$  be the image of  $l$  in  $(\mathbb{G}_m)^t(K) = (K^{\times})^t$ , and as in No. 5.5, let  $(l_{v,j})_{1 \leq j \leq s}$  be the image of  $l$  in  $(\mathbb{G}_a)^s(K) = K^s$ . Note that

(a)  $\varphi \in G(\mathcal{O}_{X,v})$  if and only if  $l'_{v,j} \in \mathcal{O}_{X,v}^{\times}$  for  $1 \leq j \leq t$  and  $l_{v,j} \in \mathcal{O}_{X,v}$  for  $1 \leq j \leq s$ .

By construction of the transformation  $\tau_{\varphi}$  in [10, Section 2.2], we have the following (b) and (c).

(b) The étale part of  $\tau_{\varphi}$

$$\tau_{\varphi,\text{ét}} : \mathbb{Z}^t \longrightarrow \underline{\text{Div}}_X^0(k)$$

sends the  $j$ -th base of  $\mathbb{Z}^t$  ( $1 \leq j \leq t$ ) to the divisor  $\sum_v \text{ord}_v(l'_{v,j}) v$ .

(c) The infinitesimal part of  $\tau_{\varphi}$

$$\tau_{\varphi,\text{inf}} : (\widehat{\mathbb{G}}_a)^s \longrightarrow \underline{\text{Div}}_X^0$$

has the form

$$(a_j)_{1 \leq j \leq s} \longmapsto \exp \left( \sum_{j=1}^s a_j f_j \right)$$

for some  $f_j \in \Gamma(X, K/\mathcal{O}_X) = \text{Lie}(\underline{\text{Div}}_X^0)$  ( $1 \leq j \leq s$ ) such that for any point  $v$  of  $X$  of codimension one, the stalk of  $f_j$  at  $v$  coincides with  $l_{v,j} \text{ mod } \mathcal{O}_{X,v}$ .

Condition (i') is equivalent to the condition that the following (i'<sub>ét</sub>) and (i'<sub>inf</sub>) are satisfied.

(i'<sub>ét</sub>) The image of  $\tau_{\varphi, \text{ét}}$  is contained in the étale part of  $\mathcal{F}_{X,Y}$ .

(i'<sub>inf</sub>) The image of  $\tau_{\varphi, \text{inf}}$  is contained in the infinitesimal part of  $\mathcal{F}_{X,Y}$ .

By the above (b), (i'<sub>ét</sub>) is equivalent to the condition that the following (i'<sub>ét,v</sub>) is satisfied for any point  $v$  of  $X$  of codimension one.

(i'<sub>ét,v</sub>) If  $e_v = 0$ , then  $l'_{v,j} \in \mathcal{O}_{X,v}^\times$  for  $1 \leq j \leq t$ .

On the other hand, by the above (c), (i'<sub>inf</sub>) is equivalent to

$$f_j \in \Gamma(X, J^{-1}/\mathcal{O}_X) \text{ for } 1 \leq j \leq s,$$

and hence equivalent to the condition that the following (i'<sub>inf,v</sub>) is satisfied for any point  $v$  of  $X$  of codimension one.

(i'<sub>inf,v</sub>) If  $e_v = 0$ , then  $l_{v,j} \in \mathcal{O}_{X,v}$  for  $1 \leq j \leq s$ .

If  $e_v \geq 1$ , then  $\text{ord}_v(l_{v,j}) \geq 1 - e_v$  for  $1 \leq j \leq s$ .

By (a) above, for each  $v$ , (i'<sub>ét,v</sub>) and (i'<sub>inf,v</sub>) are satisfied if and only if  $\text{mod}_v(\varphi) \leq e_v$ . □

**COROLLARY 5.2.** — *For any  $\mathcal{F} \in \Lambda$ , there exists an effective divisor  $Y$  such that  $\mathcal{F} \subset \mathcal{F}_{X,Y}$ .*

*Proof.* — Let  $Y = \text{mod}(\alpha_{\mathcal{F}})$  be the modulus of the rational map  $\alpha_{\mathcal{F}} : X \rightarrow \text{Alb}_{\mathcal{F}}(X)$  associated with  $\mathcal{F} \in \Lambda$ . Then  $\mathcal{F} = \text{Image}(\tau_{\alpha_{\mathcal{F}}}) \subset \mathcal{F}_{X,Y}$ . □

### 6. Proof of Theorem 1.1

We prove Theorem 1.1. Let  $X$  be a proper smooth algebraic variety over  $\mathbb{C}$  of dimension  $n$ , and let  $Y$  be an effective divisor on  $X$ . Let  $I$  be the ideal of  $\mathcal{O}_X$  which defines  $Y$ , let  $I_1$  be the ideal of  $\mathcal{O}_X$  which defines the reduced part of  $Y$ , and let  $J = II_1^{-1} \subset \mathcal{O}_X$ .

**6.1.** Let  $H^1(X, Y_+)(1)$  be the object of  $\mathcal{H}_{1, \{0, -1\}}$  corresponding to the object  $[\mathcal{F}_{X,Y} \rightarrow \text{Pic}^0(X)]$  of  $\mathcal{M}_{1, \{0, -1\}}$  in the equivalence of categories of Theorem 4.1. Let  $H^{2n-1}(X, Y_-)(n)$  be the object of  $\mathcal{H}_{1, \{-1, -2\}}$  corresponding to the object  $\text{Alb}(X, Y)$  of  $\mathcal{M}_{1, \{-1, -2\}}$ .

Since the equivalence of categories in Theorem 4.1 is compatible with dualities, we have

$$(6.1) \quad H^{2n-1}(X, Y_-)(n) \cong \underline{\text{Hom}}(H^1(X, Y_+)(1), \mathbb{Z})(1).$$

We prove Theorem 1.1 in the following way. First in No. 6.3, we give an explicit description of  $H^1(X, Y_+)(1)$ . From this, by (6.1), we can obtain an

explicit description of  $H^{2n-1}(X, Y_-)(n)$  as in No. 6.4. Since  $\text{Alb}(X, Y)$  corresponds to  $H^{2n-1}(X, Y_-)(n)$  in the equivalence of categories  $\mathcal{H}_{1, \{-1, -2\}} \simeq \mathcal{M}_{1, \{-1, -2\}}$ , we can obtain from No. 6.4 the explicit descriptions of  $\text{Alb}(X, Y)$  as stated in Theorem 1.1.

We define objects  $H^1(X, Y_+)$  and  $H^{2n-1}(X, Y_-)$  of  $\mathcal{H}$  as follows:  $H^1(X, Y_+)$  is the Tate twist  $(H^1(X, Y_+)(1))(-1)$  of  $H^1(X, Y_+)(1)$ , and  $H^{2n-1}(X, Y_-)$  is the Tate twist  $(H^{2n-1}(X, Y_-)(n))(-n)$  of  $H^{2n-1}(X, Y_+)(n)$ . These are natural generalizations of the objects of  $\mathcal{H}$  for the curve case considered in Example 2.1.

**6.2.** We define canonical  $\mathbb{C}$ -linear maps

$$(6.2) \quad H^1(X - Y, \mathbb{C}) \longrightarrow H^1(X, \mathcal{O}_X),$$

$$(6.3) \quad H^{n-1}(X, \Omega_X^n) \longrightarrow H_c^{2n-1}(X - Y, \mathbb{C})$$

First assume that  $Y$  is with normal crossings. Then by [5], we have canonical isomorphisms

$$\begin{aligned} H^m(X - Y, \mathbb{C}) &\cong H^m(X, \Omega_X^\bullet(\log(Y))), \\ H_c^m(X - Y, \mathbb{C}) &\cong H^m(X, \Omega_X^\bullet(-\log(Y))) \end{aligned}$$

for  $m \in \mathbb{Z}$ , where  $\Omega_X^p(\log(Y))$  is the sheaf of differential  $p$ -forms with log poles along  $Y$ , and  $\Omega_X^p(-\log(Y)) = I_1 \Omega_X^p(\log(Y))$ . Since  $\mathcal{O}_X = \Omega_X^0(\log(Y))$  and  $\Omega_X^n = \Omega_X^n(-\log(Y))$ , we have canonical maps of complexes  $\Omega_X^\bullet(\log(Y)) \rightarrow \mathcal{O}_X$  and  $\Omega_X^\bullet[-n] \rightarrow \Omega_X^\bullet(-\log(Y))$ . These maps induce the maps (6.2) and (6.3) in the case  $Y$  is with normal crossings, respectively.

In general, take a birational morphism  $X' \rightarrow X$  of proper smooth algebraic varieties over  $\mathbb{C}$  such that the inverse image  $Y'$  of  $Y$  on  $X'$  is with normal crossings. Then we have maps

$$H^{n-1}(X, \Omega_X^n) \longrightarrow H^{n-1}(X', \Omega_{X'}^n) \longrightarrow H_c^{2n-1}(X' - Y', \mathbb{C}) = H_c^{2n-1}(X - Y, \mathbb{C})$$

where the second arrow is the map (6.3) for  $X'$ , and the composition  $H^{n-1}(X, \Omega_X^n) \rightarrow H_c^{2n-1}(X - Y, \mathbb{C})$  is independent of the choice of  $X' \rightarrow X$ . The  $\mathbb{C}$ -linear dual of (6.3) with respect to the Poincaré duality and Serre duality gives the map (6.2). The map (6.2) is also obtained as the composition

$$H^1(X - Y, \mathbb{C}) = H^1(X' - Y', \mathbb{C}) \longrightarrow H^1(X', \mathcal{O}_{X'}) \xleftarrow{\simeq} H^1(X, \mathcal{O}_X).$$

**6.3.** Let  $H = H^1(X, Y_+)(1)$ , the object of  $\mathcal{H}_{1, \{0, -1\}}$  corresponding to the object  $[\mathcal{F}_{X, Y} \rightarrow \text{Pic}^0(X)]$  of  $\mathcal{M}_{1, \{0, -1\}}$ . We describe  $H$ . By [3, Thm. 4.7] which treats the case when  $Y$  has no multiplicity, we can identify  $H_{\mathbb{Z}}$  with  $H^1(X - Y, \mathbb{Z}(1))$  and identify the map  $H_{\mathbb{C}} \rightarrow \text{Lie}(\text{Pic}^0(X)) = H^1(X, \mathcal{O}_X)$

with the map (6.2) in No. 6.2. We have  $H_V = H_{\mathbb{C}} \oplus H^0(X, J^{-1}/\mathcal{O}_X)$ , the maps  $a : H_{\mathbb{C}} \rightarrow H_V$  and  $b : H_V \rightarrow H_{\mathbb{C}}$  are the evident ones, the weight filtration is given by  $W_0H = H$ ,  $W_{-2}H = 0$ ,

$$\begin{aligned} W_{-1}H_{\mathbb{Q}} &= H^1(X, \mathbb{Q}(1)), \\ W_{-1}H_V &= H^1(X, \mathbb{C}), \end{aligned}$$

and the Hodge filtration is given by  $F^{-1}H_V = H_V$ ,  $F^1H_V = 0$ , and

$$F^0H_V = \text{Ker} (H^1(X - Y, \mathbb{C}) \oplus H^0(J^{-1}/\mathcal{O}_X) \rightarrow H^1(X, \mathcal{O}_X))$$

where the map  $H^0(J^{-1}/\mathcal{O}_X) \rightarrow H^1(X, \mathcal{O}_X)$  is the connecting map of the exact sequence  $0 \rightarrow \mathcal{O}_X \rightarrow J^{-1} \rightarrow J^{-1}/\mathcal{O}_X \rightarrow 0$ .

**6.4.** Let  $H = H^{2n-1}(X, Y_{-})(n)$ , the object of  $\mathcal{H}_{1, \{-1, -2\}}$  corresponding to the object  $\text{Alb}(X, Y)$  of  $\mathcal{M}_{1, \{-1, -2\}}$ . By (6.1) in No. 6.1, we obtain the following description of  $H$  from the description of  $H^1(X, Y_{+})(1)$  in No. 6.3.

$$\begin{aligned} H_{\mathbb{Z}} &= H_c^{2n-1}(X - Y, \mathbb{Z})/(\text{torsion}), \\ H_V &= H_{\mathbb{C}} \oplus H^{n-1}(X, \Omega_X^n/J\Omega_X^n), \end{aligned}$$

the maps  $a : H_{\mathbb{C}} \rightarrow H_V$  and  $b : H_V \rightarrow H_{\mathbb{C}}$  are the evident ones, the weight filtration is given by  $W_{-1}H = H$ ,  $W_{-3}H = 0$ ,

$$\begin{aligned} W_{-2}H_{\mathbb{Q}} &= \text{Ker} (H_{\mathbb{Q}} \rightarrow H^{2n-1}(X, \mathbb{Q}(n))), \\ W_{-2}H_V &= \text{Ker} (H_V \rightarrow H^{2n-1}(X, \mathbb{C})), \end{aligned}$$

and the Hodge filtration is given by  $F^{-1}H_V = H_V$ ,  $F^1H_V = 0$ , and

$$F^0H_V = \text{Image}(H^{n-1}(X, \Omega_X^n) \rightarrow H_c^{2n-1}(X - Y, \mathbb{C}) \oplus H^{n-1}(X, \Omega_X^n/J\Omega_X^n))$$

where the map  $H^{n-1}(X, \Omega_X^n) \rightarrow H_c^{2n-1}(X - Y, \mathbb{C})$  is (6.3) in No. 6.2 and the map  $H^{n-1}(X, \Omega_X^n) \rightarrow H^{n-1}(X, \Omega_X^n/J\Omega_X^n)$  is the evident one.

**6.5.** We prove Theorem 1.1 (2). Let  $H = H^{2n-1}(X, Y_{-})(n)$ . Then

$$\text{Alb}(X, Y) = H_{\mathbb{Z}} \setminus H_V / F^0H_V$$

by No. 4.5. Hence the description of  $H^{2n-1}(X, Y_{-})(n)$  in No. 6.4 proves Theorem 1.1 (2).

**6.6.** As a preparation for the proof of Theorem 1.1 (1), we review a kind of Serre-duality obtained in the appendix by Deligne of the book [8].

Let  $S$  be a proper scheme over a field  $k$ , let  $C$  be a closed subscheme of  $S$ , let  $U = S - C$ , and let  $I_C$  be the ideal of  $\mathcal{O}_S$  which defines  $C$ . Assume  $U$  is

smooth over  $k$  and purely of dimension  $n$ . Let  $\mathcal{F}$  be a coherent  $\mathcal{O}_S$ -module. Then for any  $p \in \mathbb{Z}$ , we have a canonical isomorphism

$$H^p(U, R \operatorname{Hom}_{\mathcal{O}_U}(\mathcal{F}|_U, \Omega_U^n)) \cong \varinjlim_m \operatorname{Hom}_k(H^{n-p}(X, I_C^m \mathcal{F}), k).$$

In the case when  $C$  is empty and  $\mathcal{F}$  is locally free, this is the usual Serre duality

$$H^p(X, \operatorname{Hom}_{\mathcal{O}_X}(\mathcal{F}, \Omega_X^n)) \cong \operatorname{Hom}_k(H^{n-p}(X, \mathcal{F}), k).$$

**6.7.** We start the proof of Theorem 1.1 (1).

Let  $C_Y$  be the subcomplex of  $\Omega_X^\bullet$  defined as

$$C_Y^p = \ker(\Omega_X^p \rightarrow \Omega_Y^p) \text{ for } 0 \leq p \leq n - 1, \quad C_Y^n = J\Omega_X^n.$$

**PROPOSITION 6.1.** — *For  $p = 2n, 2n - 1$ , the maps  $H_c^p(X - Y, \mathbb{C}) \rightarrow H^p(X, C_Y)$  induced by the homomorphism  $j: \mathbb{C} \rightarrow C_Y$  are isomorphisms.*

**6.8.** We prove Proposition 6.1 in the case  $Y = Y_1$ . We have an exact sequence of complexes

$$0 \longrightarrow C_{Y_1} \longrightarrow \Omega_X^\bullet \longrightarrow \Omega_{Y_1}^{\leq n-1} \longrightarrow 0.$$

Since the support of  $\Omega_{Y_1}^{\leq n-1}$  is of dimension  $\leq n - 1$  and since  $\Omega_{Y_1}^{\leq n-1}$  has only terms of degree  $\leq n - 1$ , we have  $H^p(X, \Omega_{Y_1}^{\leq n-1}) = 0$  for  $p \geq 2n - 1$ . Hence

$$H^{2n}(X, C_{Y_1}) \cong H^{2n}(X, \Omega_X^\bullet) \cong H^{2n}(X, \mathbb{C}) \cong H_c^{2n}(X - Y, \mathbb{C}).$$

The above exact sequence of complexes induces the lower row of the commutative diagram with exact rows

$$\begin{array}{ccccccc} H^{2n-2}(X, \mathbb{C}) & \rightarrow & H^{2n-2}(Y_1, \mathbb{C}) & \rightarrow & H_c^{2n-1}(X \setminus Y_1, \mathbb{C}) & \rightarrow & H^{2n-1}(X, \mathbb{C}) \rightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^{2n-2}(X, \Omega_X^\bullet) & \rightarrow & H^{2n-2}(Y_1, \Omega_{Y_1}^{\leq n-1}) & \rightarrow & H^{2n-1}(X, C_{Y_1}) & \rightarrow & H^{2n-1}(X, \Omega_X^\bullet) \rightarrow 0. \end{array}$$

The vertical arrows except possibly the map  $H_c^{2n-1}(X - Y_1, \mathbb{C}) \rightarrow H^{2n-1}(X, C_{Y_1})$  are isomorphisms. Hence the last map is also an isomorphism.

**LEMMA 6.2.** — *Let  $Y'$  and  $Y''$  be effective divisors on  $X$  whose supports coincide with  $Y_1$  and assume  $Y' \geq Y''$ . Then the canonical map  $H^{2n-1}(X, C_{Y'}) \rightarrow H^{2n-1}(X, C_{Y''})$  is surjective and the canonical map  $H^{2n}(X, C_{Y'}) \rightarrow H^{2n}(X, C_{Y''})$  is an isomorphism.*

*Proof.* — Let  $N = C_{Y''}/C_{Y'}$ . We have

$$N^p = \operatorname{Ker}(\Omega_{Y'}^p \rightarrow \Omega_{Y''}^p) \text{ for } 0 \leq p \leq n - 1, \quad N^n = J''\Omega_X^n / J'\Omega_X^n.$$

Here,  $J' = I'I_1^{-1}$ ,  $J'' = I''I_1^{-1}$  with  $I'$  (resp.  $I''$ ) the ideal of  $\mathcal{O}_X$  which defines  $Y'$  (resp.  $Y''$ ). Since the support of  $N$  is of dimension  $\leq n-1$  and  $N$  has only terms of degree  $\leq n$ , we have  $H^{2n}(X, N) = 0$ . Hence it is sufficient to prove  $H^{2n-1}(X, N) = 0$ .

Let  $\Sigma$  be the set of all singular points of  $Y_1$ . Then  $\Sigma$  is of dimension  $\leq n-2$ . Let  $\Omega_X^\bullet(\log(Y_1))$  be the de Rham complex on  $X - \Sigma$  with log poles along  $Y_1 - \Sigma$ . Then, as is easily seen, the restriction of  $C_Y$  to  $X - \Sigma$  coincides with  $I\Omega_X^\bullet(\log(Y_1))$ . Let  $I_\Sigma$  be the ideal of  $\mathcal{O}_X$  defining  $\Sigma$  (here  $\Sigma$  is endowed with the reduced structure). For  $k \geq 0$ , let  $N_k$  be the sub-complex of  $N$  defined by  $N_k^p = I_\Sigma^{\max(k-p, 0)}N^p$ . In particular,  $N_0 = N$ . Then if  $k \geq j \geq 0$ , since the support of  $N_j/N_k$  is of dimension  $\leq n-2$  and  $N_j/N_k$  has only terms of degree  $\leq n$ , we have  $H^{2n-1}(X, N_j/N_k) = 0$ . Hence  $H^{2n-1}(X, N_k) \rightarrow H^{2n-1}(X, N_j)$  is surjective. Applying No. 6.6 for  $S = X$  and  $C = \Sigma$  yields that  $\varprojlim_k H^{2n-1}(X, N_k)$  is the dual vector space of  $H^0(X - \Sigma, [(J')^{-1}/(J'')^{-1} \xrightarrow{d} (J')^{-1}\Omega_X(\log Y_1)/(J'')^{-1}\Omega_X(\log Y_1)])$ . Since  $d : (J')^{-1}/(J'')^{-1} \rightarrow (J')^{-1}\Omega_X(\log Y_1)/(J'')^{-1}\Omega_X(\log Y_1)$  is injective, the last cohomology group is 0. Hence  $H^{2n-1}(X, N_k) = 0$  for all  $k \geq 0$ . In particular,  $H^{2n-1}(X, N) = 0$ . □

**6.9.** We prove Proposition 6.1 in general. By Lemma 6.2, the map  $\varprojlim_{Y'} H^{2n-1}(X, C_{Y'}) \rightarrow H^{2n-1}(X, C_Y)$  is surjective, where  $Y'$  ranges over all effective divisors on  $X$  whose supports coincide with  $Y_1$ . By No. 6.6, which we apply by taking  $S = X$  and  $C = Y$ , we have that  $\varprojlim_{Y'} H^{2n-1}(X, C_{Y'})$  is the dual vector space of  $H^1((X - Y)_{\text{zar}}, \Omega_{X-Y, \text{alg}}^\bullet)$  where “zar” means Zariski topology and “alg” means the algebraic version. But  $H^1((X - Y)_{\text{zar}}, \Omega_{X-Y, \text{alg}}^\bullet) \simeq H^1(X - Y, \mathbb{C})$  by Grothendieck’s Theorem [7, Thm. 1’]. This proves  $\varprojlim_{Y'} H^{2n-1}(X, C_{Y'}) \cong H_c^{2n-1}(X - Y, \mathbb{C})$ . Hence the map  $H_c^{2n-1}(X - Y, \mathbb{C}) \rightarrow H^{2n-1}(X, C_Y)$  is surjective. Since the composition  $H_c^{2n-1}(X - Y, \mathbb{C}) \rightarrow H^{2n-1}(X, C_Y) \rightarrow H^{2n-1}(X, C_{Y_1}) \cong H_c^{2n-1}(X - Y, \mathbb{C})$  is the identity map, the map  $H_c^{2n-1}(X - Y, \mathbb{C}) \rightarrow H^{2n-1}(X, C_Y)$  is an isomorphism.

**6.10.** We prove (1) of Theorem 1.1. Let  $S_Y = \text{Ker}(\Omega_X^\bullet \rightarrow \Omega_Y^{\leq n-1})$ . Then  $C_Y \subset S_Y \subset C_{Y_1}$ . We have an exact sequence of complexes

$$0 \rightarrow C_Y \rightarrow S_Y \rightarrow \Omega_X^n/J\Omega_X^n[-n] \rightarrow 0.$$

Hence we have an exact sequence

$$H^{2n-1}(X, C_Y) \rightarrow H^{2n-1}(X, S_Y) \rightarrow H^{n-1}(X, \Omega_X^n/J\Omega_X^n) \rightarrow H^{2n}(X, C_Y) \\ \rightarrow H^{2n}(X, S_Y).$$

Note that for  $p = 2n, 2n - 1$ , the compositions

$$H^p(X, C_Y) \longrightarrow H^p(X, S_Y) \longrightarrow H^p(X, C_{Y_1})$$

are isomorphisms by Proposition 6.1. Hence by Proposition 6.1, we have an isomorphism

$$H^{2n-1}(X, S_Y) \cong H_c^{2n-1}(X - Y, \mathbb{C}) \oplus H^{n-1}(X, \Omega_X^n/J\Omega_X^n)$$

which is compatible with the maps from  $H^{n-1}(X, \Omega_X^n)$ . Hence (1) of Theorem 1.1 follows from (2) of Theorem 1.1.

## BIBLIOGRAPHY

- [1] L. BARBIERI-VIALE, “Formal Hodge theory”, *Math. Res. Lett.* **14** (2007), no. 3, p. 385-394.
- [2] L. BARBIERI-VIALE & A. BERTAPELLE, “Sharp de Rham realization”, *Adv. Math.* **222** (2009), no. 4, p. 1308-1338.
- [3] L. BARBIERI-VIALE & V. SRINIVAS, “Albanese and Picard 1-motives”, *Mém. Soc. Math. Fr. (N.S.)* (2001), no. 87, p. vi+104.
- [4] S. BLOCH & V. SRINIVAS, “Enriched Hodge structures”, in *Algebra, arithmetic and geometry, Part I, II (Mumbai, 2000)*, Tata Inst. Fund. Res. Stud. Math., vol. 16, Tata Inst. Fund. Res., Bombay, 2002, p. 171-184.
- [5] P. DELIGNE, “Théorie de Hodge. II et III”, *Inst. Hautes Études Sci. Publ. Math.* (1971 et 1974), no. 40 et 44, p. 5-78 et 5-77.
- [6] H. ESNAULT, V. SRINIVAS & E. VIEHWEG, “The universal regular quotient of the Chow group of points on projective varieties”, *Invent. Math.* **135** (1999), no. 3, p. 595-664.
- [7] A. GROTHENDIECK, “On the de Rham cohomology of algebraic varieties”, *Inst. Hautes Études Sci. Publ. Math.* (1966), no. 29, p. 95-103.
- [8] R. HARTSHORNE, *Residues and duality*, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20, Springer-Verlag, Berlin, 1966, vii+423 pages.
- [9] G. LAUMON, “Transformation de Fourier généralisée”, 1996, Preprint arXiv:alg-geom/9603004.
- [10] H. RUSSELL, “Generalized Albanese and its dual”, *J. Math. Kyoto Univ.* **48** (2008), no. 4, p. 907-949.



- [11] ———, “Albanese varieties with modulus over a perfect field”, 2010, Preprint arXiv:0902.2533v2.

Manuscrit reçu le 26 avril 2011,  
accepté le 8 février 2011.

Kazuya KATO  
University of Chicago  
Department of Mathematics  
Chicago, IL 60637 (USA)  
kkato@math.uchicago.edu

Henrik RUSSELL  
Universität Duisburg-Essen  
FB6 Mathematik, Campus Essen  
45117 Essen (Germany)  
henrik.russell@uni-due.de