



ANNALES

DE

L'INSTITUT FOURIER

Daniel SKODLERACK

The centralizer of a classical group and Bruhat-Tits buildings

Tome 63, n° 2 (2013), p. 515-546.

http://aif.cedram.org/item?id=AIF_2013__63_2_515_0

© Association des Annales de l'institut Fourier, 2013, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

THE CENTRALIZER OF A CLASSICAL GROUP AND BRUHAT-TITS BUILDINGS

by Daniel SKODLERACK

ABSTRACT. — Let G be a unitary group defined over a non-Archimedean local field of odd residue characteristic and let H be the centralizer of a semisimple rational Lie algebra element of G . We prove that the Bruhat-Tits building $\mathfrak{B}^1(H)$ of H can be affinely and G -equivariantly embedded in the Bruhat-Tits building $\mathfrak{B}^1(G)$ of G so that the Moy-Prasad filtrations are preserved. The latter property forces uniqueness in the following way. Let j and j' be maps from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ which preserve the Moy-Prasad filtrations. We prove that if there is no split torus in the center of the connected component of H then j and j' are equal, and in general if both maps are affine and satisfy a mild equivariance condition they differ up to a translation of $\mathfrak{B}^1(H)$.

RÉSUMÉ. — Soit G un groupe unitaire défini sur un corps local non-Archimédien de caractéristique résiduelle impaire et soit H le centralisateur d'un élément rationnel semi-simple de l'algèbre de Lie de G . Nous démontrons qu'il existe une application affine injective H -équivariante de l'immeuble de Bruhat-Tits $\mathfrak{B}^1(H)$ de H vers l'immeuble de Bruhat-Tits $\mathfrak{B}^1(G)$ de G qui préserve les filtrations de Moy-Prasad. La dernière propriété implique l'unicité comme suit : soient j et j' des applications de $\mathfrak{B}^1(H)$ vers $\mathfrak{B}^1(G)$ qui préservent les filtrations de Moy-Prasad. Nous démontrons que j et j' sont égales s'il n'y a pas de tore déployé dans le centre de la composante connexe de H . En général, les deux diffèrent par une translation de $\mathfrak{B}^1(H)$ si elles sont affines et vérifient une autre condition faible.

1. Introduction

The subject of this article is a functoriality question for maps between Bruhat-Tits buildings which is connected with the representation theory of classical groups. Embeddings of buildings of reductive groups have previously been studied for example by Landvogt [11] and Prasad and Yu [15]. The aim of this article is to show to what extent the property of preserving

Keywords: Building, classical group over a local field, centralizer.

Math. classification: 11E57, 11E95, 14L35, 20E42, 20G25.

the Moy-Prasad filtrations determines the choice of an embedding uniquely. It completes recent works of Broussous, Lemaire and Stevens [2], [4], which have applications in representation theory.

More precisely Bushnell's and Kutzko's strategy in their theory of types for the classification of irreducible smooth representations of $\mathrm{GL}_n(k)$ in [9] is applied to other classical groups defined over a non-Archimedean local field k . In [17], [18], [19], [20], [3] and [21] Sécherre together with Broussous and Stevens gave the classification for $\mathrm{GL}_n(D)$, where D is a central finite division algebra over k . Further in [22] Stevens constructed types for unitary groups. He applied a result of his paper with Broussous [4], *i.e.* he used an embedding of Bruhat-Tits buildings for a certain subgroup of a unitary group to apply an induction. The important property of this map is the compatibility with the Lie algebra filtrations (**CLF**) which by [12] correspond to the Moy-Prasad filtrations [13]. In [4] the quaternion algebra case is missing and the authors proposed a uniqueness and generalization conjecture to the reader.

Let k be a non-Archimedean local field with valuation ν and of residue characteristic $\neq 2$, and let D be a division algebra central and finite-dimensional over k equipped with an involution ρ , whose set of fixed points in k is denoted by k_0 . To the classical group $\mathbf{G} := \mathbf{U}(h)$, *i.e.* the unitary group of an ϵ -hermitian form h on a finite-dimensional right D -vector space V , is attached the Bruhat-Tits building $\mathfrak{B}^1(\mathbf{G}, k_0)$ which can be described in terms of lattice functions. To every point of $\mathfrak{B}^1(\mathbf{G}, k_0)$ there is attached a Lie algebra filtration \mathfrak{g}_x which is exactly the Moy-Prasad filtration. More precisely if x is a point of $\mathfrak{B}^1(\mathbf{G}, k_0)$ seen as a lattice function it can be interpreted as a point of $\mathfrak{B}(\mathbf{GL}_D(V), k)$ which has a Lie algebra filtration $\tilde{\mathfrak{g}}_x$ in $\mathrm{End}_D(V)$. If one identifies $\mathrm{Lie}(\mathbf{G})(k_0)$ with the set of skew-symmetric elements of $\mathrm{End}_D(V)$ with respect to the adjoint involution of h , the filtration \mathfrak{g}_x of x coincides with

$$t \mapsto \tilde{\mathfrak{g}}_x(t) \cap \mathrm{Lie}(\mathbf{G})(k_0).$$

Now we take an element $\beta \in \mathrm{Lie}(\mathbf{G})(k_0)$ whose k -algebra $k[\beta]$ is a product of field extensions E_i of k . In this introduction let us assume $k[\beta]$ to be separable over k .

Its centralizer in \mathbf{G} is an algebraic group \mathbf{H} defined over k_0 which is a product of restrictions of scalars to k_0 of classical groups \mathbf{H}_i which are either general linear groups or unitary groups. In the manner of [4] we prove the existence of an injective affine $\mathbf{H}(k_0)$ -equivariant and toral map j_β from $\mathfrak{B}^1(\mathbf{H}, k_0)$ to $\mathfrak{B}^1(\mathbf{G}, k_0)$ using lattice function models. In addition

j_β has the CLF-property, *i.e.*

$$\mathrm{Lie}(\mathbf{H})(k_0) \cap \mathfrak{g}_{j_\beta(y)}$$

is the Lie algebra filtration \mathfrak{h}_x of x . This article solves the following problem: To what extent does the CLF-property determine j_β ?

In order to give an answer to this question we consider two cases.

- (a) If only unitary groups appear among the \mathbf{H}_i , none of which is k_0 -isomorphic to the isotropic orthogonal group of rank one, then j_β is uniquely determined by CLF.
- (b) If there are no restrictions on the \mathbf{H}_i then j_β is unique up to a translation of $\mathfrak{B}^1(\mathbf{H}, k_0)$ in being affine, $Z(\mathbf{H}^0(k_0))$ -equivariant and having the CLF-property.

For the proofs we use the decomposition of $k[\beta]$ as a product of fields to restrict to the cases where $k[\beta]$ is a field or a product of two fields interchanged by the adjoint involution of h . Let us call these cases atomic cases. The map j_β is now induced by j_E^{-1} constructed in [2]. In (a) $k[\beta]$ is a field in the atomic case. If β is non-zero we follow the strategy of [4]. The statement follows essentially from a uniqueness statement for j_E^{-1} given in [4, 10.3]. If β is zero we prove that for all unitary groups except for the isotropic orthogonal group of rank one the Moy-Prasad filtration determines the point completely. For (b) to restrict to the atomic case we need further a rigidity proposition for Euclidean buildings, stated in 10.2. We use (a) to finish the proof of (b).

The whole strategy and the proofs do not require β to be separable. In that case we define the building of $Z_{\mathbf{G}(k_0)}(\beta)$ in view of [4] using lattice functions and we work mainly with the rational points instead of the algebraic groups.

The article is structured in the following way. After preliminary notation in §2 the model of the Bruhat-Tits building of \mathbf{G} over k_0 in terms of lattice functions is explained in §3. In §4 we introduce the Moy-Prasad filtration for our purposes. We give the building of the centralizer in §5 and introduce the notion of CLF in §6 followed by the existence theorem in §7. The uniqueness theorems in §9, where no GL_m is involved, and §11, for the general case, are prepared in the preceding sections. Lastly in §12 we show that the constructed map respects apartments.

I thank very much P. Broussous, Prof. S. Stevens and Prof. E.-W. Zink for useful hints and fruitful communications. I want to express my gratitude to the German Research Foundation, who supported this work within the framework of BMS and SFB 878.

2. Notation

We are given a division algebra D of finite index d and central over a non-Archimedean local field k of odd residue characteristic. The valuation on k and its unique extension to D are denoted by ν . We assume that the image of $\nu|_{k^\times}$ is \mathbb{Z} . Further let ρ be an involution on D , i.e. a skew field isomorphism from D to D^{op} of order one or two, in particular ρ is an isometry. The fixed field of ρ in k is denoted by k_0 .

Remark 2.1 ([16], 10.2.2). — The existence of ρ implies $d \leq 2$. If $k \neq k_0$, then $d = 1$.

We fix an element $\epsilon \in \{1, -1\}$ and a non-degenerate ϵ -hermitian form h on an m -dimensional right D -vector space V , i.e. a \mathbb{Z} -bi-linear map h on $V \times V$ with values in D such that

$$h(v_1\lambda_1, v_2\lambda_2) = \epsilon\rho(\lambda_1)\rho(h(v_2, v_1))\lambda_2$$

for all $\lambda_1, \lambda_2 \in D$ and $v_1, v_2 \in V$. Further $\rho|_k$ extends to the adjoint involution σ of h on $\text{End}_D(V)$. For a skew field D' with discrete valuation the symbols $\mathfrak{o}_{D'}$ and $\mathfrak{p}_{D'}$ denote the valuation ring and its maximal ideal respectively. We fix an algebraic closure \bar{k} of k .

By a Bruhat-Tits building we always understand the extended one [6, 4.2.16.]. The set

$$G := \mathbf{U}(\sigma) := \{f \in \text{End}_D(V) \mid \sigma(f)f = \text{id}_V\}$$

is the set of k_0 -rational points of an algebraic group $\mathbf{U}(\sigma)$ defined over k_0 . We also write $\mathbf{U}(h)$ for $\mathbf{U}(\sigma)$. We denote $\mathbf{U}(h)$ by \mathbf{G} and $\text{Lie}(\mathbf{G})(k_0)$ by \mathfrak{g} and identify the latter with the set $\text{Skew}(\text{End}_D(V), \sigma)$ of skew-symmetric elements of $\text{End}_D(V)$ with respect to σ . The Bruhat-Tits building $\mathfrak{B}^1(\mathbf{G}, k_0)$ is also denoted by $\mathfrak{B}^1(G)$. We repeat the strategy in [4] to describe the building as a set of self-dual lattice functions based on the description using norms, see [8].

3. Norms and lattice functions

The description of the Bruhat-Tits building in terms of norms and lattice functions needs some basic properties which are collected in this section. Proofs and more details for norms can mainly be found in [7], [8]. For lattice functions we refer to [4] and [2].

DEFINITION 3.1 ([7], 1.1). — A D -norm on V is a map $\alpha: V \rightarrow \mathbb{R} \cup \{\infty\}$ such that

- $\alpha(v) = \infty$ if and only if $v = 0$,
- $\alpha(v\lambda) = \nu(\lambda) + \alpha(v)$,
- $\alpha(v + w) \geq \inf\{\alpha(v), \alpha(w)\}$,

for all $v, w \in V$ and $\lambda \in D$. Fix a D -norm α on V . The dual $\alpha^\#$ of α with respect to h is defined to be the D -norm

$$v \in V \mapsto \inf\{\nu(h(v, w)) - \alpha(w) \mid w \in V, w \neq 0\}.$$

One calls α self-dual with respect to h if it is equal to his dual. The set of D -norms and self-dual D -norms of V are denoted by $\text{Norm}_D^1(V)$ and $\text{Norm}_h^1(V)$ respectively. Given a further right D -vector space V' and a norm $\alpha' \in \text{Norm}_D^1(V')$ the direct sum of α and α' is defined by

$$(\alpha \oplus \alpha')(v + v') := \inf\{\alpha(v), \alpha'(v')\}, \quad v \in V, v' \in V'.$$

We consider a set \mathcal{R} of D -subspaces of V whose direct sum is V . We call \mathcal{R} a frame if all elements of \mathcal{R} are one dimensional. An element $\alpha \in \text{Norm}_D^1(V)$ is split by \mathcal{R} if α is the direct sum of the $\alpha|_W$, $W \in \mathcal{R}$. If in addition \mathcal{R} is a frame and $(w_i)_i$ is a D -basis of V consisting of elements of $\bigcup_{W \in \mathcal{R}} W$ one calls $(w_i)_i$ a splitting basis of α .

Remark 3.2 ([7], 1.26). — Every pair of D -norms on V has a common splitting basis.

Given two norms $\alpha, \gamma \in \text{Norm}_D^1(V)$ with common splitting basis (v_i) and a real number $\lambda \in [0, 1]$ the convex combination of α with γ with λ is defined to be the D -norm on V split by (v_i) whose value at v_i is

$$\lambda\alpha(v_i) + (1 - \lambda)\gamma(v_i).$$

This definition does not depend on the choice of the splitting basis and we get an affine structure on $\text{Norm}_D^1(V)$. The $\text{Aut}_D(V)$ -action on $\text{Norm}_D^1(V)$, more precisely

$$(g \cdot \alpha)(v) := \alpha(g^{-1}v),$$

restricts to a $U(\sigma)$ -action on $\text{Norm}_h^1(V)$.

The family of balls of a D -norm α leads to the idea of a lattice function:

$$\Lambda_\alpha(t) := \{v \in V \mid \alpha(v) \geq t\}, \quad t \in \mathbb{R}.$$

Before we give the definition we want to remark that by an o_D -lattice in V we mean a finitely generated o_D -submodule of V which contains a D -basis of V , i.e. we omit the word “full”.

DEFINITION 3.3 ([2], I.2.1). — A family $(\Lambda(r))_{r \in \mathbb{R}}$ of o_D -lattices in V is said to be an o_D -lattice function in V if

- (1) $\Lambda(r) \supseteq \Lambda(s)$,
- (2) $\Lambda(r) = \bigcap_{t < r} \Lambda(t)$ and
- (3) $\Lambda(r)\mathfrak{p}_D = \Lambda(r + \frac{1}{d})$,

for all $r, s \in \mathbb{R}$ with $r < s$. The set of o_D -lattice functions in V is denoted by $\text{Latt}_{o_D}^1(V)$. For the right limit of Λ at s we write $\Lambda(s+)$, i.e.

$$\Lambda(s+) := \bigcup_{t > s} \Lambda(t).$$

For $t \in \mathbb{R}$ we define $[t]$ to be smallest integer not smaller than t .

PROPOSITION 3.4 ([2], I.2.4). — The map

$$\alpha \mapsto \Lambda_\alpha$$

is a bijection from $\text{Norm}_D^1(V)$ to $\text{Latt}_{o_D}^1(V)$. Its inverse is given by

$$\alpha_\Lambda(v) := \sup\{t \in \mathbb{R} \mid v \in \Lambda(t)\}.$$

All notions for norms carry over to lattice functions in the following way. The dual of a lattice function Λ is the lattice function which corresponds to $\alpha_\Lambda^\#$. A lattice function is called *split* by a given basis if the corresponding norm is split by this basis. The $\text{Aut}_D(V)$ -action on $\text{Norm}_D^1(V)$ defines an $\text{Aut}_D(V)$ -action on $\text{Latt}_{o_D}^1(V)$ via push forward.

PROPOSITION 3.5 ([4], 3.3, [2], I.2.4). — Let Λ be an o_D -lattice function in V . Let g be an element of $\text{Aut}_D(V)$ and let \mathcal{R} be a set of D -subspaces of V whose direct sum is V .

- (1) The function Λ is split by \mathcal{R} if and only if, for all real numbers t , the lattice $\Lambda(t)$ is the direct sum of the o_D -modules $W \cap \Lambda(t)$, $W \in \mathcal{R}$.
- (2) If $(v_i)_i$ is a splitting basis of Λ , then there are real numbers a_i , $1 \leq i \leq m$, such that, for all t , we have

$$\Lambda(t) = \bigoplus_i v_i \mathfrak{p}_D^{\lceil (t-a_i)d \rceil}.$$

We call $(a_i)_i$ the coordinate tuple of Λ with respect to (v_i) . The map which assigns the coordinate tuple to a lattice function split by (v_i) is an affine bijection onto \mathbb{R}^m .

- (3) We have $(g \cdot \Lambda)(t) = g(\Lambda(t))$.
- (4) The dual $\Lambda^\#$ of Λ with respect to h is the o_D -lattice function whose value at $t \in \mathbb{R}$ is

$$\{v \in V \mid h(v, \Lambda((-t)+)) \subseteq \mathfrak{p}_D\}.$$

Proof. — All statements except for 4 can be found in part I of [2]. Point 4 has been proven in [4, 3.3] for the case $D = k$, but the proof is valid for the general case if one replaces o_F by o_D . □

The set of self-dual o_D -lattice functions is denoted by $\text{Latt}_h^1(V)$. We recall that we have fixed an ϵ -hermitian form h on V .

THEOREM 3.6 ([8], 2.12). — *There is a unique affine and G -equivariant bijection from $\mathfrak{B}^1(G)$ to $\text{Norm}_h^1(V)$.*

Propositions 3.4 and 3.5 imply the existence of a unique affine and G -equivariant bijection from $\mathfrak{B}^1(G)$ to $\text{Latt}_h^1(V)$. It defines on $\text{Latt}_h^1(V)$ a system of apartments which correspond to the Witt-decompositions of V .

DEFINITION 3.7. — *A set of one dimensional h -isotropic D -subspaces W_l of V , $l \in L$, is said to be a Witt decomposition of V if:*

- (1) *for every $l \in L$ there is exactly one index $l' \in L$ such that $h(W_l, W_{l'})$ is non-zero,*
- (2) *the sum of the W_l is direct, and*
- (3) *the orthogonal complement W of the sum of the W_l is anisotropic.*

A lattice function is said to be split by a Witt decomposition $\{W_l \mid l \in L\}$ if it is split by $\{W_l \mid l \in L\} \cup \{W\}$. A D -basis of V is adapted to $\{W_l \mid l \in L\}$ if all basis elements lie in the union of W and all W_l . Further we assume $0 \notin L$ and denote W by W_0 .

Witt decompositions always exists, and even more, for every element of $\text{Latt}_h^1(V)$ there is a splitting Witt decomposition. A proof for the latter in terms of norms can be found in [8, 2.13].

Remark 3.8 ([8], 2.9).

- (1) The system of apartments of $\text{Latt}_h^1(V)$ is the system of sets

$$\text{Latt}_{h,\mathcal{S}}^1(V) := \{\Lambda \in \text{Latt}_h^1(V) \mid \Lambda \text{ is split by } \mathcal{S}\},$$

where \mathcal{S} runs over the set of Witt decompositions of V with respect to h .

- (2) A self dual D -norm α split by a Witt decomposition $\{W_l \mid l \in L\}$ satisfies

$$\alpha(w_0) = \frac{1}{2} \nu(h(w_0, w_0)),$$

for all $w_0 \in W_0$ (see [8, 2.9]).

PROPOSITION 3.9. — *If α is a self dual D -norm on V and $\{W_l \mid l \in L\}$ is a Witt decomposition splitting α , then every orthogonal D -basis of W_0 splits $\alpha|_{W_0}$.*

Proof. — Let w' and w'' be elements of W_0 which are orthogonal to each other, then we have

$$\begin{aligned} \alpha(w') &= \alpha\left(\frac{1}{2}(w' + w'') + \frac{1}{2}(w' - w'')\right) \\ &\geq \inf\left\{\alpha\left(\frac{1}{2}(w' + w'')\right), \alpha\left(\frac{1}{2}(w' - w'')\right)\right\} \\ &= \alpha(w' + w''), \end{aligned}$$

and the last equality follows from

$$h\left(\frac{1}{2}(w' + w''), \frac{1}{2}(w' + w'')\right) = h\left(\frac{1}{2}(w' - w''), \frac{1}{2}(w' - w'')\right)$$

and $\nu(2) = 0$. □

4. The Moy-Prasad filtrations

An o_D -lattice function Λ gives rise to an o_k -lattice function in $A := \text{End}_D(V)$

$$\tilde{g}_\Lambda(t) := \{a \in A \mid a(\Lambda(s)) \subseteq \Lambda(s + t), \forall s \in \mathbb{R}\}$$

called the *square lattice function* of Λ in A . It defines a Lie algebra filtration of Λ in \mathfrak{g} by intersection

$$\mathfrak{g}_\Lambda(t) := \tilde{g}_\Lambda(t) \cap \mathfrak{g}.$$

In [13] Moy-Prasad attached to every point x of $\mathfrak{B}^1(G', k')$ a filtration $(\mathfrak{a}_{x,t})_{t \in \mathbb{R}}$ of $\text{Lie}(G')(k')$ for a reductive group G' defined over a non-Archimedean local field k' . The next theorem states that in our special situation of unitary groups we do not need the quite involved description.

THEOREM 4.1 ([12]). — *The Lie algebra filtration of a point of $\mathfrak{B}^1(G)$ is exactly its Moy-Prasad filtration.*

In this article the square lattice functions are the key for the rigidity of Lie algebra filtrations because of the following proposition.

PROPOSITION 4.2 ([4], 3.5). — *Let Λ be an o_D -lattice function in V .*

- (1) *The o_k -lattice function $(\sigma(\tilde{g}_\Lambda(t)))_{t \in \mathbb{R}}$ is the square lattice of $\Lambda^\#$ and we denote it by \tilde{g}_Λ^σ . Square lattice functions fixed under σ are said to be self-dual.*
- (2) *The map*

$$\Lambda \in \text{Latt}_h^1(V) \mapsto \tilde{g}_\Lambda$$

is injective and onto to the set $\text{Latt}_\sigma^2(A)$ of self-dual square lattice functions.

Proof. — The proof in [4, 3.5] is valid if one replaces F by D . □

5. The centralizer

In the following the symbol $Z_*(?)$ denotes the centralizer of $?$ in $*$, and $Z(?)$ denotes the center of $?$. Let us fix an element $\beta \in \text{Skew}(\text{End}_D(V), \sigma)$, such that the k -algebra $k[\beta]$ is a product of fields E_i , $i \in I$, with identity element 1_i . We call β *separable* if $E_i|1_i k$ is separable for all i . The centralizers $Z_G(\beta)$ and $Z_{\mathbf{G}}(\beta)$ are denoted by H and \mathbf{H} . If β is separable then the algebraic group \mathbf{H} is reductive and defined over k_0 and its set of k_0 -rational points is H (see appendix A).

Notation 5.1. — There is an action of σ on I via $\sigma(1_i) =: 1_{\sigma(i)}$. We denote by I_0 the fixed point set of the action of σ on I , and we divide the set $I \setminus I_0$ into two disjoint parts, i.e. we choose a positive part I_+ and a negative part I_- , such that

$$\sigma(I_+) = I_-.$$

We define $-i := \sigma(i)$ and put $V_i := 1_i V$. For $i \in I_0$ we denote by $(E_i)_0$ the set of fixed points of σ in E_i , and for $i \in I_+$ we put $(E_i)_0 := E_i$.

The group H is the product of sets of rational points of classical groups over the $(E_i)_0$. More precisely, for $i \in I$, the E_i -algebra $\text{End}_{E_i \otimes_k D}(V_i)$ is E_i -algebra isomorphic to $\text{End}_{D'_i}(V'_i)$, for some finite-dimensional vector space V'_i over some skew field D'_i central and of finite index over E_i . We define

$$\mathbf{H}_i := \begin{cases} \mathbf{U}(\sigma|_{\text{End}_{D'_i}(V'_i)}), & i \in I_0 \\ \mathbf{GL}_{D'_i}(V'_i), & i \in I_+ \end{cases}$$

and $H_i := \mathbf{H}_i((E_i)_0)$. There is a canonical group isomorphism from H to $\prod_{i \in I_+ \cup I_0} H_i$ which motivates the following definition for the Bruhat-Tits building of H :

$$\mathfrak{B}^1(H) := \prod_{i \in I_+ \cup I_0} \mathfrak{B}^1(\mathbf{H}_i, (E_i)_0).$$

Remark 5.2. — If β is separable the above isomorphism extends to a k_0 -isomorphism

$$(5.1) \quad \mathbf{H} \cong \prod_{i \in I_0 \cup I_+} \text{Res}_{(E_i)_0|k_0}(\mathbf{H}_i).$$

Thus there is a isomorphism of affine buildings from $\mathfrak{B}^1(\mathbf{H}, k_0)$ to $\mathfrak{B}^1(H)$.

The Lie algebra of H is isomorphic to

$$\bigoplus_{i \in I_0} \text{Skew}(\text{End}_{E_i \otimes_k D}(V_i), \sigma) \oplus \bigoplus_{i \in I_+} \text{End}_{E_i \otimes_k D}(V_i)$$

and we denote the i th factor by \mathfrak{g}_i . We define the Lie algebra filtration of $x = (x_i)_{i \in I_0 \cup I_+}$ to be the direct sum of the Lie algebra filtrations of the x_i , i.e.

$$\mathfrak{h}_x(t) := \bigoplus_{i \in I_0 \cup I_+} (\mathfrak{g}_i)_{x_i}(t).$$

where we take the square lattice function for $i \in I_+$ and identify $\text{End}_{E_i \otimes_k D}(V_i)$ with $\text{End}_{D'_i}(V_i)$.

6. Compatibility with the Lie algebra filtrations

DEFINITION 6.1.

- (1) A set with Lie algebra filtrations (LF-set) is a pair (X, L) such that X is a set, L is a Lie algebra, and for each $x \in X$, there is Lie algebra filtration $((L_x)(t))_{t \in \mathbb{R}}$, i.e. a decreasing sequence of subsets of L .
- (2) Given two LF-sets (X, L) and (X', L') and a map $\phi: L \rightarrow L'$
 - (a) a map $f: X \rightarrow X'$ is compatible with the Lie algebra filtrations (CLF) if

$$\phi(L_x(t)) = \text{im}(\phi) \cap L'_{f(x)}(t)$$

holds, for all $x \in X$ and all real numbers t .

- (b) $g: X' \rightarrow X$ is compatible with the Lie algebra filtrations (CLF) if

$$\phi(L_{g(x')}(t)) = \text{im}(\phi) \cap L'_{x'}(t)$$

holds, for all $x' \in X'$ and all real numbers t .

In this article we consider buildings together with the set of rational points of a Lie algebra of an algebraic group. Mostly ϕ is a canonical inclusion. An example of a CLF-map is given in [2]. The theorems in this section are valid without the assumption on the residue characteristic and the existence of an involution on D . Firstly, we introduce two important LF-sets after the following theorem.

DEFINITION 6.2.

- (1) Two o_D -lattice functions Λ and Λ' on V are equivalent if there is a real number t such that

$$\Lambda(s) = \Lambda'(s - t) =: (\Lambda' + t)(s),$$

for all $s \in \mathbb{R}$. The set of equivalence classes of elements of $\text{Latt}_{o_D}^1(V)$ is denoted by $\text{Latt}_{o_D}(V)$. A map f on $\text{Latt}_{o_D}^1(V)$ of the form

$$f(\Lambda) = \Lambda + t$$

is called a translation. For a map g , such that $f \circ g$ exists, we define

$$g + t := f \circ g.$$

(2) An action of $\text{Aut}_D(V)$ on $\text{Latt}_{o_D} V$ is given by

$$a.[\Lambda] := [a.\Lambda], \text{ where } (a.\Lambda)(t) := a(\Lambda(t)).$$

(3) The affine structure on $\text{Latt}_{o_D}^1(V)$ induces an affine structure on $\text{Latt}_{o_D}(V)$.

Let us consider the non-extended building $\mathfrak{B}(\text{Aut}_D(V))$ and the extended building $\mathfrak{B}^1(\text{Aut}_D(V))$.

THEOREM 6.3 ([2], I.2.4, [7], 2.13, [7], 2.11(iii)). — *The extended building of $\text{Aut}_D(V)$ is in affine and $\text{Aut}_D(V)$ -equivariant bijection with the set $\text{Latt}_{o_D}^1(V)$. For two such $\text{Aut}_D(V)$ -equivariant affine bijections f and g , there is a real number t , such that*

$$f = g + t,$$

and the map f induces via

$$[x] \mapsto [f(x)]$$

a unique affine and $\text{Aut}_D(V)$ -equivariant bijection from $\mathfrak{B}(\text{Aut}_D(V))$ to $\text{Latt}_{o_D}(V)$.

Remark 6.4. — For $x \in \mathfrak{B}^1(\text{Aut}_D(V))$, we attach to x and to $[x]$ the square lattice function of a corresponding o_D -lattice function. By the above theorem both LF-sets $(\mathfrak{B}^1(\text{Aut}_D(V)), \text{End}_D(V))$ and $(\mathfrak{B}(\text{Aut}_D(V)), \text{End}_D(V))$ are well-defined, i.e. do not depend on the choices made. In $(\mathfrak{B}(\text{Aut}_D(V)), \text{End}_D(V))$, a point is uniquely determined by its Lie algebra filtration (see [2, I.4.5]), whereas in $(\mathfrak{B}^1(\text{Aut}_D(V)), \text{End}_D(V))$ two points with the same Lie algebra filtration are translates of each other.

THEOREM 6.5. — [2, II.1.1] *Let $E|k$ be a field extension in $\text{End}_D(V)$, $\tilde{G} := \text{Aut}_D(V)$ and $\tilde{H} := Z_{\tilde{G}}(E)$. We consider the non-extended buildings $\mathfrak{B}(\tilde{G})$ and $\mathfrak{B}(\tilde{H})$ as LF-sets as constructed in the above remark. Then there is a unique CLF-map j_E from $\mathfrak{B}(\tilde{G})^{E^\times}$ to $\mathfrak{B}(\tilde{H})$. The map has the following properties.*

- (1) *It is bijective.*
- (2) *It is \tilde{H} -equivariant.*
- (3) *It is affine.*

The map $j^E := (j_E)^{-1}$ is the unique map satisfying 2. and 3..

In the above setting the CLF-property of j^E implies uniqueness.

THEOREM 6.6 ([4], 10.3). — Under the assumptions of Theorem 6.5, suppose that the points $y \in \mathfrak{B}(\tilde{G})$ and $x \in \mathfrak{B}(\tilde{H})$ satisfy

$$\tilde{\mathfrak{g}}_y \cap \text{End}_{E \otimes_k D}(V) \supseteq \tilde{\mathfrak{h}}_x.$$

Then $j^E(x) = y$.

Proof. — In [4, 10.3] this theorem was proven for the case where $D = k$ and E is generated by one element. The proof did not use the second assumption, and it carries over to $D \neq k$ without changes. \square

The map j^E is induced from a map between the extended buildings. We fix an E -algebra isomorphism from $\text{End}_{E \otimes_k D}(V)$ to $\text{End}_{D'}(V')$ where D' is a central skew field over E and V' is a right D' -vector space.

THEOREM 6.7 ([2], II.3.1, II.4, [7], 2.11(iii)). — There is a bijective affine \tilde{H} -equivariant CLF-map

$$\tilde{j}^E : \text{Latt}_{o_{D'}}^1(V') \rightarrow \text{Latt}_{o_D}^1(V)^{E^\times}$$

such that

$$[\tilde{j}^E(\Lambda)] = j^E([\Lambda])$$

for all $\Lambda \in \text{Latt}_{o_{D'}}^1(V')$. The image is the set of o_D -lattice functions which are in addition o_E -lattice functions. For every other bijective affine \tilde{H} -equivariant map j from $\text{Latt}_{o_{D'}}^1(V')$ to $\text{Latt}_{o_D}^1(V)^{E^\times}$ the composition $j^{-1} \circ \tilde{j}^E$ is a translation of $\text{Latt}_{o_{D'}}^1(V')$.

Proof. — The existence of \tilde{j}^E is stated in Lemma [2, II.3.1]. The affinity is proven in [2, II.4] for j_E , but the proof actually shows that \tilde{j}^E is affine. The \tilde{H} -equivariance follows from the formula given in [2, II.3.1]. The fact that the image of j^E is the set of E^\times -fixed points implies that $\text{im}(\tilde{j}^E)$ only consists of o_E -lattice functions and contains a representative for every element of $\text{im}(j^E)$. Being E^\times -equivariant and convex, the image of \tilde{j}^E must contain every o_E -lattice function of $\text{Latt}_{o_D}^1(V)$. The last assertion follows directly from [7, 2.11 (iii)]. \square

7. CLF-map from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$

We are now returning to the situation of section 5. Before we state the first theorem we give a definition in analogy with the set of fixed points in the building.

DEFINITION 7.1. — An o_E - o_D -lattice function of V is an o_D -lattice function in V which splits under (V_i) such that, for every i , the function

$$t \mapsto \Lambda(t) \cap V_i$$

is an o_{E_i} -lattice function in V_i . We denote the set of o_E - o_D -lattice functions by $\text{Latt}_{o_E, o_D}^1(V)$.

The next theorem has been proven for $D = k$ in [4].

THEOREM 7.2. — There is an injective, affine and H -equivariant CLF-map

$$j: \mathfrak{B}^1(H) \rightarrow \mathfrak{B}^1(G)$$

whose image in terms of lattice functions is the set of self-dual o_E - o_D -lattice functions in V .

Proof. — The product decomposition $H = \prod_{i \in I_+ \cup I_0} Z_{U(h_i |_{(V_i + V_{-i})^2})}(\beta_i + \beta_{-i})$ leads us to three steps:

- (1) the case where E is a field;
- (2) the case where I_+ has cardinality one;
- (3) the general case.

Step 1: We use the map j^E from Theorem 6.5 and the following diagram.

$$\begin{array}{ccc} \text{Latt}_{o_{D'}}(V') & \xrightarrow{j^E} & \text{Latt}_{o_D}(V) \\ \uparrow & & \uparrow \\ \mathfrak{B}^1(H) & & \mathfrak{B}^1(G) \end{array}$$

We have to prove that, for $x \in \mathfrak{B}^1(H)$, the square lattice function of $j^E(x)$ is self-dual. The latter follows from Theorem 6.6 because the square lattice function of x is self-dual and j^E is a CLF-map. The assertion about the image of $j^E|_{\mathfrak{B}^1(H)}$ follows from 6.5 and the CLF-property.

Step 2: To prove the lemma we denote the unique element in I_+ by i and we define

$$\Lambda^{\#-i}(t) := \{v \in V_{-i} \mid h(v, \Lambda((-t)+)) \subseteq \mathfrak{p}_D\}$$

for $\Lambda \in \text{Latt}_{o_D}^1(V_i)$. The map

$$(7.1) \quad g \in \text{Aut}_D(V_i) \mapsto (g, 0) + \sigma((g^{-1}, 0)) \in \text{Aut}_D(V_i) \oplus \text{Aut}_D(V_{-i})$$

defines a k -embedding from $\mathbf{GL}_D(V_i)$ to \mathbf{G} mapping $Z_{\mathbf{GL}_D(V_i)}(\beta_i)$ onto H . An injective, affine and $\text{Aut}_D(V_i)$ -equivariant CLF-map from $\mathfrak{B}^1(\mathbf{GL}_D(V_i), k)$ to $\mathfrak{B}^1(G)$ in terms of lattice functions is given by

$$(7.2) \quad \Lambda \in \text{Latt}_{o_D}^1(V_i) \mapsto (\Lambda \oplus \Lambda^{\#-i}) \in \text{Latt}_h^1(V).$$

Now we apply Theorem 6.7 to finish step 2.

Step 3: We take the direct sum of the maps constructed in the steps before. □

8. Factorization

In order to analyze the set of CLF-maps from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ this section reduces the problem to the case Where $I_0 \cup I_+$ has cardinality one.

LEMMA 8.1. — *There is at most one index $i \in I$ such that $\beta_i = 0$ and if such an index exists, it has to be in I_0 .*

Proof. — Assume that β_i and β_j are zero for two different indices i and j . Taking a polynomial P with coefficients in k such that $1_i = P(\beta)$ we obtain firstly

$$1_i = 1_i 1_i = 1_i P(\beta) = 1_i P(0)$$

and secondly

$$0 = 1_i 1_j = P(\beta) 1_j = P(0) 1_j$$

which is a contradiction. The second assertion follows from $-\beta_{-i} = \sigma(\beta_i)$ and the uniqueness. □

Let us recall that the embedding of \mathfrak{h} into \mathfrak{g} is realized by mapping an element $(a_i)_{i \in I_0 \cup I_+}$ of $(\prod_{i \in I_0} \text{Skew}(\text{End}_{E_i \otimes_k D}(V_i), \sigma)) \times (\prod_{i \in I_+} \text{End}_{E_i \otimes_k D}(V_i))$ to $(\sum_{i \in I_0} a_i) + (\sum_{i \in I_+} (a_i - \sigma(a_i)))$.

ASSUMPTION 8.2. — *For the rest of the section we fix elements $y \in \mathfrak{B}^1(G)$ and $x \in \mathfrak{B}^1(H)$ such that*

$$\mathfrak{g}_y \cap \mathfrak{h} = \mathfrak{h}_x.$$

The element x is given by a tuple $(x_i)_{i \in I_0 \cup I_+}$. The set $\text{Lie}(\mathbf{H}_i)((E_i)_0)$ is denoted by \mathfrak{h}_i . For $i \in I_0$, we write \mathfrak{h}_{x_i} and $\tilde{\mathfrak{h}}_{x_i}$ for the Lie algebra filtration and the square lattice function of x_i , respectively.

LEMMA 8.3. — *The lattice function corresponding to y splits under $\{V_i \mid i \in I\}$.*

Proof. — The assertion is equivalent to the fact that all idempotents 1_i are elements of $\tilde{\mathfrak{g}}_y(0)$.

Case 1: We first consider an index $i \in I_+$. Then 1_i is an element of $\mathfrak{h}_{x_i}(0)$ and thus $1_i - 1_{-i}$ is an element of $\mathfrak{g}_y(0)$ by 8.2. Therefore

$$1_i + 1_{-i} = (1_i - 1_{-i})^2 \in \tilde{\mathfrak{g}}_y(0).$$

Hence 1_i and 1_{-i} are elements of $\tilde{\mathfrak{g}}_y(0)$, since 2 is invertible in o_k .

Case 2: We take an index $i \in I_0$ and we assume that β_i is not zero. Since β_i is skew-symmetric and central, we have, for all $t \in \mathbb{R}$,

$$\beta_i \tilde{\mathfrak{h}}_{x_i}(t) = \tilde{\mathfrak{h}}_{x_i}(t + \nu(\beta_i))$$

and

$$\beta_i \mathfrak{h}_{x_i}(t) = \tilde{\mathfrak{h}}_{x_i}(t + \nu(\beta_i)) \cap \{w \in \tilde{\mathfrak{h}}_i \mid \sigma|_{\tilde{\mathfrak{h}}_i}(w) = w\}.$$

By the invertibility of 2 in o_k , every element of $\tilde{\mathfrak{h}}(t)$ is a sum of a skew-symmetric and a symmetric element of $\tilde{\mathfrak{h}}_{x_i}(t)$, which implies

$$\begin{aligned} \tilde{\mathfrak{h}}_{x_i}(0) &= \mathfrak{h}_{x_i}(0) + \beta_i \mathfrak{h}_{x_i}(-\nu(\beta_i)) \\ &\subseteq \mathfrak{g}_y(0) + \mathfrak{g}_y(\nu(\beta_i))\mathfrak{g}_y(-\nu(\beta_i)) \\ &\subseteq \tilde{\mathfrak{g}}_y(0). \end{aligned}$$

Thus the i th idempotent 1_i is an element of $\tilde{\mathfrak{g}}_y(0)$.

Case 3: If there is an index i_0 such that $\beta_{i_0} = 0$, it is unique by Lemma 8.1 and the two cases above imply

$$1_{i_0} = 1 - \sum_{i \neq i_0} 1_i \in \tilde{\mathfrak{g}}_y(0).$$

□

The idea of the proof of case 2 is taken from [4, 11.2]. We define $\mathbf{G}_i := \mathbf{U}(h|_{(V_i+V_{-i})^2})$, for $i \in I_0 \cup I_+$.

COROLLARY 8.4. — *For non-negative indices there are elements $y_i \in \mathfrak{B}^1(\mathbf{G}_i, k_0)$, such that the direct sum of the lattice functions of the y_i is the lattice function of y .*

Analogous to the definitions for G we use \mathfrak{g}_i and \mathfrak{g}_{y_i} for $\text{Lie}(\mathbf{G}_i)(k_0)$ and the Lie algebra filtration of y_i , respectively.

LEMMA 8.5. — *For all $i \in I_0 \cup I_+$ we have*

$$\mathfrak{h}_{x_i} = \mathfrak{h}_i \cap \mathfrak{g}_{y_i}.$$

Proof. — For $t \in \mathbb{R}$ we have

$$\begin{aligned} \bigoplus_i \mathfrak{h}_{x_i}(t) &= \mathfrak{h}(t) = \mathfrak{g}_y(t) \cap \mathfrak{h} = \left(\mathfrak{g}_y(t) \cap \left(\bigoplus_i \mathfrak{g}_i \right) \right) \cap \mathfrak{h} \\ &= \left(\bigoplus_i \mathfrak{g}_{y_i}(t) \right) \cap \left(\bigoplus_i \mathfrak{h}_i \right) = \bigoplus_i (\mathfrak{g}_{y_i}(t) \cap \mathfrak{h}_i), \end{aligned}$$

where i runs over $I_0 \cup I_+$, and we obtain the assertion. □

The last two lemmas lead to a factorization of any CLF-map. More precisely we prove the following proposition.

PROPOSITION 8.6. — *Let ψ_I denote the canonical map from $\mathfrak{B}^1(\prod_i \mathbf{G}_i, k_0)$ to $\mathfrak{B}^1(\mathbf{G}, k_0)$ which maps a tuple of self-dual lattice functions to its sum. Every CLF-map j from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ factors under ψ_I , i.e. there is a unique map*

$$\tau: \mathfrak{B}^1(H) \rightarrow \mathfrak{B}^1\left(\prod_i \mathbf{G}_i, k_0\right) \text{ such that } j = \psi_I \circ \tau.$$

The map τ is

- (1) a CLF-map,
- (2) affine if j is affine, and
- (3) H -equivariant if j is H -equivariant.

Proof. — The image of j is contained in the image of the injective, affine and $\prod_{i \in I_0 \cup I_+} \mathbf{G}_i(k_0)$ -equivariant map ψ_I by Corollary 8.4. The CLF-property of τ is a consequence of Lemma 8.5. This proves the proposition. □

9. The case where I_+ is empty

We denote the following algebraic group

$$\left\{ \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \begin{pmatrix} 0 & \lambda \\ \lambda^{-1} & 0 \end{pmatrix} \mid \lambda \in \bar{k}^\times \right\}$$

by \mathbf{O}_2^{is} . It is a form of \mathbf{O}_2 over the prime field of k .

Remark 9.1.

- (1) The group $\mathbf{U}(h)$ is k_0 -isomorphic to \mathbf{O}_2^{is} if and only if $D = k = k_0$, V is two-dimensional over k and isotropic with respect to h and σ is orthogonal.
- (2) The connected component of \mathbf{O}_2^{is} is k -isomorphic to \mathbf{G}_m , implying that $\mathfrak{B}^1(\mathbf{O}_2^{is}, k)$ is affinely isomorphic to \mathbb{R} .
- (3) All points of $\mathfrak{B}^1(\mathbf{O}_2^{is}, k)$ have the same Lie algebra filtration. Especially all of its affine endomorphisms are CLF-maps.

The remark forces us to exclude \mathbf{O}_2^{is} from the factors.

THEOREM 9.2. — *There exists only one CLF-map from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ if I_+ is empty and no group \mathbf{H}_i is $(E_i)_0$ -isomorphic to \mathbf{O}_2^{is} .*

For the proof we need the following operations on square matrices.

DEFINITION 9.3. — For a square matrix $B = (b_{i,j}) \in M_r(D)$ the matrix \tilde{B} is defined to be $(b_{r+1-j,r+1-i})_{i,j}$, i.e. \tilde{B} is obtained from B by a reflection along the antidiagonal. We define further

$$B^\rho := (\rho(b_{i,j}))_{i,j}.$$

Proof. — Applying Lemma 8.5 we can assume that E is a field.

Case 1: β is not zero. Compare with [4, 11.2]. We fix an arbitrary CLF-map j from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ and an arbitrary element x of $\mathfrak{B}^1(H)$. By the same argument as in case 2 of Lemma 8.3 we obtain

$$\tilde{\mathfrak{h}}_x(t) \subseteq \tilde{\mathfrak{g}}_{j(x)}(t)$$

for all real numbers t . Theorem 6.6 implies the uniqueness.

Case 2: β is zero. If σ is of the second type there is a skew-symmetric non-zero element β' in k and we can replace β by β' and apply case 1. Thus we only need to consider the case where σ is of the first kind. We fix a point $y \in \mathfrak{B}^1(G)$ and fix an apartment containing y . This apartment is determined by a Witt decomposition. We choose an adapted basis $(w_i)_{1 \leq j \leq m}$ such that the Gram matrix $\text{Gram}_{(w_i)}(h)$ of the ϵ -hermitian form h has the form

$$\begin{pmatrix} 0 & M & 0 \\ \epsilon M & 0 & 0 \\ 0 & 0 & N \end{pmatrix}$$

with $M := \text{antidiag}(1, \dots, 1)$ and a diagonal regular matrix N . From 3.9 we deduce that the self-dual \mathcal{o}_D -lattice function Λ corresponding to y is split by this basis. It is thus described by its intersections with the lines $w_i D$, i.e. there are real numbers a_i such that

$$\Lambda(t) = \bigoplus_i w_i \mathfrak{p}_D^{\lceil (t-a_i)d \rceil}.$$

Thus the square lattice function of y in t is

$$\tilde{\mathfrak{g}}_y(t) = \bigoplus_{i,j} \mathfrak{p}_D^{\lceil (t+a_j-a_i)d \rceil} E_{i,j}$$

where $E_{i,j}$ denotes the matrix with a 1 in the intersection of the i th row and the j th column and zeros everywhere else. See for example [2, I.4.5].

It is enough to show that $\tilde{\mathfrak{g}}_y$ is determined by the Lie algebra filtration \mathfrak{g}_y . Indeed, a class of \mathcal{o}_D -lattice functions contains at most one self-dual lattice function. Thus the self-dual square lattice function of a point of $\mathfrak{B}^1(G)$ determines the point uniquely.

The adjoint involution of h

$$B \mapsto B^\sigma = \text{Gram}_{(w_i)}(h)^{-1} (B^\rho)^T \text{Gram}_{(w_i)}(h)$$

on $M_m(D)$ has under $(w_i)_{1 \leq j \leq m}$ the form

$$\begin{pmatrix} B_{1,1} & B_{1,2} & B_{1,3} \\ B_{2,1} & B_{2,2} & B_{2,3} \\ B_{3,1} & B_{3,2} & B_{3,3} \end{pmatrix} \mapsto \begin{pmatrix} \tilde{C}_{2,2} & \epsilon \tilde{C}_{1,2} & \epsilon MC_{3,2}^T N \\ \epsilon \tilde{C}_{2,1} & \tilde{C}_{1,1} & MC_{3,1}^T N \\ \epsilon N^{-1} C_{2,3}^T M & N^{-1} C_{1,3}^T M & N^{-1} C_{3,3}^T N \end{pmatrix}$$

The matrices $B_{1,1}, B_{1,2}, B_{2,1}$ and $B_{2,2}$ are $r \times r$ -matrices and $C := B^\rho$ where r is the Witt index of h . By the above calculation we obtain that $E_{i,j}^\sigma$ is $+E_{i,j}, -E_{i,j}$ or $\lambda E_{u,l}$ with $(i, j) \neq (u, l)$ for some $\lambda \in D^\times$. From the self-duality of $\tilde{\mathfrak{g}}_y$ and since 2 is invertible in o_k we get:

$$\mathfrak{g}_y(t) \cap k(E_{i,j} - E_{i,j}^\sigma) = \mathfrak{p}_k^{\lceil t+a_j-a_i \rceil} (E_{i,j} - E_{i,j}^\sigma).$$

For the calculation see Lemma 9.4 below. Thus we can obtain the exponent $a_j - a_i$ from the knowledge of the Lie algebra filtration if $E_{i,j}$ is not fixed by σ . We now consider two cases.

Case 2.1: We assume that there is an anisotropic part in the Witt decomposition, i.e. N occurs. The matrix $E_{i,m}$ is fixed by σ if and only if i equals m . Thus from the knowledge of the Lie algebra filtration we know all differences $a_i - a_m$ for all indexes i different from m , and thus by subtractions we know the differences $a_i - a_j$ for all i and j .

Case 2.2: Now we assume that there is no anisotropic part in the Witt decomposition. If ϵ is -1 , no $E_{i,j}$ is fixed and we can obtain the differences $a_i - a_j$ for all i and j . As a consequence, we only have to consider the case where h is hermitian and $D = k$ (see [8, 1.14]). Here the matrix $E_{i,j}$ is fixed by σ if and only if $i + j = m + 1$. Thus we can determine all differences $a_i - a_j$ where $i + j \neq m + 1$. If m is at least 4 for an index i there is an index $k \neq i$ with $i + k \neq m + 1$ and we can obtain $a_i - a_{m+1-i}$ if we add $a_k - a_{m+1-i}$ to $a_i - a_k$. If m equals 2, then the group \mathbf{G} is k -isomorphic to \mathbf{O}_2^{is} which is excluded by the assumption of the theorem. \square

The idea of taking the root system of $\mathbf{GL}_D(V)$ with respect to the Witt basis was given by P. Broussous. To complete the proof we need the following lemma.

LEMMA 9.4. — For all $t \in \mathbb{R}$ we have

$$\mathfrak{p}_D^{\lceil td \rceil} \cap k = \mathfrak{p}_k^{\lceil t \rceil}.$$

Proof. — For an element x of k , we have $x \in \mathfrak{p}_D^{\lceil td \rceil}$ if and only if $\nu(x) \geq \frac{\lceil td \rceil}{d}$. There are integers l and s such that $1 \leq s \leq d$ and

$$\lceil td \rceil = ld + s$$

Thus $\lceil t \rceil = l + 1$ and we get that $\nu(x) \geq \frac{\lceil td \rceil}{d}$ if and only if $\nu(x) \geq \lceil t \rceil$. The “only if” follows from $\nu(x) \in \mathbb{Z}$. \square

Remark 9.5.

- (1) In particular the proof of Theorem 9.2 shows that, if \mathbf{H}_i is $(E_i)_0$ isomorphic to \mathbf{O}_2^{is} , then β_i has to be zero and σ is of the first kind.
- (2) For positive indices \mathbf{H}_i is not isomorphic to \mathbf{O}_2 because the latter is not connected.
- (3) Let us assume that β is separable. The above remarks and 9.1 imply, for $i \in I_0 \cup I_+$, that \mathbf{O}_2^{is} is k_0 -isomorphic to $\text{Res}_{(E_i)_0|k_0}(\mathbf{H}_i)$ if and only if \mathbf{H}_i is $(E_i)_0$ -isomorphic to \mathbf{O}_2^{is} .

10. Rigidity of Euclidean buildings

To have an approach to a uniqueness statement if there are no restrictions on I we show that Euclidean buildings are rigid for functionals.

DEFINITION 10.1.

- (1) A set with affine structure is a pair $(S, *)$ consisting of a non-empty set S and a map

$$*: [0, 1] \times S \times S \rightarrow S,$$

which we denote

$$ts_1 + (1 - t)s_2 := *(t, s_1, s_2).$$

- (2) An affine functional f on a set S with affine structure is an affine map from S to \mathbb{R} , i.e.

$$f(tx + (1 - t)y) = tf(x) + (1 - t)f(y),$$

for all $t \in [0, 1]$ and $x, y \in S$.

PROPOSITION 10.2. — *Let Ω be a thick Euclidean building and $|\Omega|$ be its geometric realization. Then every affine functional a on $|\Omega|$ is constant.*

For the definition of a thick Euclidean building and its geometric realization see [5, VI.3].

Proof. — Let C_1, C_2 and C_3 be three pairwise different adjacent chambers having a common co-dimension 1 face S . We denote by P_i the unique vertex of C_i which is not a vertex of S . The line segment $[P_1, P_2]$ meets $[P_1, P_3]$ and $[P_2, P_3]$ in a point $Q \in |\bar{S}|$. This can be seen as follows. We are working in three different apartments simultaneously. If Δ_{ij} denotes an apartment containing C_i and C_j , for different i and j , the affine isomorphism from $|\Delta_{12}|$ to $|\Delta_{13}|$ fixing $|\Delta_{12} \cap \Delta_{13}|$ sends $[P_1, P_2]$ to $[P_1, P_3]$ and thus the unique intersection point in $[P_1, P_2] \cap |\bar{C}_1| \cap |\bar{C}_2|$ lies on $[P_1, P_3]$, and

similarly on $[P_3, P_2]$. Without loss of generality assume that $a(Q)$ vanishes. If $a(P_1)$ is negative then $a(P_2)$ and $a(P_3)$ are positive by the affineness of a . Thus $a(Q)$ is positive since it lies on $[P_2, P_3]$. A contradiction. Using galleries we obtain that a is constant on vertices of the same type. An apartment is affinely generated by its vertices of a fixed type. Thus a is constant on every apartment and therefore on $|\Omega|$, since any two apartments are connected by a gallery. \square

PROPOSITION 10.3. — *If \mathbf{G} is not k_0 -isomorphic to \mathbf{O}_2^{is} , then every affine functional on $\mathfrak{B}^1(G)$ is constant.*

Proof. — Not being k_0 -isomorphic to \mathbf{O}_2^{is} , the unitary group $\mathbf{U}(h)$ has no k_0 -rational characters on the connected component of the identity implying that the non-extended and the extended buildings are equal (see B.3). If \mathbf{G} is totally isotropic, then $\mathfrak{B}^1(G)$ is a point and otherwise it is the geometric realization of a thick Euclidean building. Now we apply Proposition 10.2. \square

PROPOSITION 10.4.

- (1) A k^\times -invariant affine functional on $\mathfrak{B}^1(\mathbf{GL}_D(V), k)$, i.e. on $\text{Latt}_{o_D}^1(V)$, is constant.
- (2) Every k^\times -invariant affine functional on $\mathfrak{B}^1(\mathbf{O}_2^{is}, k)$ is constant.

Proof.

- (1) Every fiber of a k^\times -invariant affine functional on $\mathfrak{B}^1(\mathbf{GL}_D(V), k)$ is a union of classes of o_D -lattice functions. It therefore factorizes to an affine functional on $\mathfrak{B}(\mathbf{GL}_D(V), k)$. Now we apply Proposition 10.2.
- (2) This follows from part 1, because $\mathfrak{B}^1(\mathbf{O}_2^{is}, k)$ is isomorphic to $\mathfrak{B}^1(\mathbf{G}_m, k)$ via a k^\times -equivariant affine bijection. \square

11. The general case

We introduce the notion of a translation in order to understand the class of affine CLF-maps which satisfy a mild equivariance condition.

DEFINITION 11.1.

- (1) A translation of $\text{Latt}_{o_D}^1(V)$ is a map from $\text{Latt}_{o_D}^1(V)$ to itself of the form

$$\Lambda \mapsto \Lambda + s$$

where s is a real number.

- (2) A translation of $\text{Latt}_h^1(V)$ is the identity if \mathbf{G} is not k_0 -isomorphic to \mathbf{O}_2^{is} .
- (3) A translation of $\mathfrak{B}^1(\mathbf{H}, k_0)$ is a product of translations of the factors.

Remark 11.2. — A translation of $\mathfrak{B}^1(\mathbf{O}_2^{is}, k_0)$ is included in the definition because $(\mathbf{O}_2^{is})^0$ is k_0 -isomorphic to \mathbf{G}_m , i.e. there is a natural bijection from $\mathfrak{B}^1(\mathbf{O}_2^{is}, k_0)$ to $\text{Latt}_{k_0}^1(k_0)$.

Let us fix a map j from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ constructed as in the proof of Theorem 7.2. In this section we are going to prove:

THEOREM 11.3. — *If ϕ is an affine and $Z(\prod_i \mathbf{H}_i^0((E_i)_0))$ -equivariant CLF-map from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ then $j^{-1} \circ \phi$ is a translation of $\mathfrak{B}^1(H)$. In terms of lattice functions, the image of ϕ is the set of self-dual o_E - o_D -lattice functions in V and ϕ is $\prod_i \mathbf{H}_i^0((E_i)_0)$ -equivariant.*

The composition of j^{-1} with ϕ is possible by the following fact.

PROPOSITION 11.4. — *The image of a CLF-map from $\mathfrak{B}^1(H)$ to $\mathfrak{B}^1(G)$ is a subset of the set of o_E - o_D -lattice functions.*

Proof. — By Lemma 8.5 we can assume that

$$I_0 \cup I_+ = \{i\}.$$

Case 1: $\beta = 0$. The field E is k and there is nothing to prove.

Case 2: $i \in I_0$ and $\beta \neq 0$. There is only one CLF-map by Theorem 9.2 and it fulfills the assertion by Theorem 7.2.

Case 3: $i \in I_+$. We choose two arbitrary points $y \in \mathfrak{B}^1(G)$ and $x \in \mathfrak{B}^1(H)$ such that $\mathfrak{g}_y \cap \mathfrak{h} = \mathfrak{h}_x$. The lattice function Λ of y splits under (V_i, V_{-i}) by corollary 8.4. By self-duality it is sufficient to prove that $\Lambda \cap V_i$ is an o_{E_i} -lattice function. The building

$$\mathfrak{B}^1(H) = \mathfrak{B}^1(\mathbf{GL}_{E_i \otimes_k D}(V_i), E_i)$$

is identified with the set of lattice functions over a skew field whose center is E_i . Thus we get

- $(a - a^\sigma) \in \mathfrak{g}_y(0)$ for all $a \in o_{E_i}^\times$,
- $\pi_{E_i} - \pi_{E_i}^\sigma \in \mathfrak{g}_y(\frac{1}{e})$, and
- $\pi_{E_i}^{-1} - (\pi_{E_i}^{-1})^\sigma \in \mathfrak{g}_y(-\frac{1}{e})$,

where e is the ramification index of $E_i|k$ and π_{E_i} is a prime element of E_i . We conclude that $1_i \Lambda$ is an o_{E_i} -lattice function □

Proof of Theorem 11.3. — Proposition 11.4 enables us to define

$$\tau := j^{-1} \circ \phi$$

because the image of j is the set of all o_E - o_D -lattice functions in V .

If we know that τ is a translation, then all assertions of Theorem 11.3 hold: A translation is a bijection and we conclude that ϕ and j have the same image. The $\prod_i \mathbf{H}_i^0((E_i)_0)$ -equivariance of ϕ follows because j and τ are $\prod_i \mathbf{H}_i^0((E_i)_0)$ -equivariant.

We denote the coordinates of τ by $\tau_i, i \in I_0 \cup I_+$. We need two steps to show that τ is a translation. Let us fix $i \in I_0 \cup I_+$.

Step 1: We prove that the coordinate τ_i only depends on x_i .

Case 1.1: $i \in I_0$, such that \mathbf{O}_2^{is} is not $(E_i)_0$ -isomorphic to \mathbf{H}_i . By Theorem 9.2 we have $\tau_i(x) = x_i$, for all $x \in \mathfrak{B}^1(H)$.

Case 1.2: We assume that we have an index $i \in I$ such that \mathbf{H}_i is $(E_i)_0$ -isomorphic to \mathbf{O}_2^{is} , whose building is affinely isomorphic to \mathbb{R} by Remark 11.2. If we fix an index $t \in (I \cup I_+) \setminus \{i\}$ and coordinates x_l for $l \in (I \cup I_+) \setminus \{t\}$, then the map

$$x_t \mapsto \tau_i(x)$$

is constant by Proposition 10.3 or 10.4. Thus τ_i does not depend on x_t .

Case 1.3: $i \in I_+$. Let us fix $x \in \mathfrak{B}^1(H)$. The lattice functions $\Lambda_{\tau_i(x)}$ and Λ_{x_i} are equivalent by the CLF-property of τ . We define $a_i(x)$ to be the real number such that

$$\Lambda_{\tau_i(x)} = \Lambda_{x_i} + a_i(x).$$

The map a_i is affine, since τ_i is. By an analogous argument as in case 1.2 we have that a_i does not depend on the t th coordinate, for $t \in (I_+ \cup I_0) \setminus \{i\}$.

Step 1 allows us to define a map $\tilde{\tau}_i$ from $\mathfrak{B}^1(\mathbf{H}_i, (E_i)_0)$ to itself by

$$\tilde{\tau}_i(x_i) := \tau_i(x), \quad x \in \mathfrak{B}^1(H).$$

Step 2: Here we show that $\tilde{\tau}_i$ is a translation. We firstly consider an index $i \in I_0$ such that \mathbf{H}_i is $(E_i)_0$ -isomorphic to \mathbf{O}_2^{is} . In this case we have $k = k_0$ and $E_i = 1_i k$. We identify $\mathfrak{B}^1(\mathbf{O}_2^{is}, k)$ with \mathbb{R} . The $\mathbf{SO}_2^{is}(k)$ -equivariance of τ_i gives

$$\tilde{\tau}_i(x_i + 1) = \tilde{\tau}_i(x_i) + 1.$$

The affineness property implies that $\tilde{\tau}_i$ is a translation. For $i \in I_+$ the map a_i in case 3 of step 1 is an affine functional and the k^\times -equivariance of τ_i implies the k^\times -invariance of a_i , because one gets in terms of lattice functions

$$\begin{aligned} \Lambda\pi_k + a_i(\Lambda\pi_k) &= \tilde{\tau}_i(\Lambda\pi_k) = (\tilde{\tau}_i(\Lambda))\pi_k \\ &= \Lambda + a_i(\Lambda) - 1 = \Lambda\pi_k + a_i(\Lambda), \end{aligned}$$

where π_k is a prime element of k . Thus a_i is constant by Proposition 10.4. □

12. Torality

In this section we want to prove that the map constructed in section 7 respects apartments and is toral if β is separable.

DEFINITION 12.1. — *A map*

$$f: \mathfrak{B}^1(G_1, k_0) \rightarrow \mathfrak{B}^1(G_2, k_0)$$

between two extended buildings of reductive groups defined over k_0 is called toral if, for each maximal k_0 -split torus S of G_1 , there is a maximal k_0 -split torus T of G_2 containing S such that f maps the apartment corresponding to S into the apartment corresponding to T . An analogous definition applies to maps between non-extended buildings.

Maximal k_0 -split tori of \mathbf{G} can be characterized in terms of Witt decompositions. Given a Witt decomposition $\{W_l \mid l \in L\}$ of V , there is exactly one maximal k_0 -split torus T of \mathbf{G} which satisfies

$$Z_{\mathbf{G}}(T)(k_0) = \{g \in \mathbf{G}(k_0) \mid g(W_l) \subseteq W_l, l \in L, (g - \text{id}_V)(W_0) \subseteq \{0\}\}.$$

We recall, that W_0 is the orthogonal complement of the sum of all W_l . Every maximal k_0 -split torus arises in this way, because they are conjugate to each other by elements of $\mathbf{G}(k_0)$. The D -vector spaces $W_l, l \in L$, are exactly the irreducible $T(k_0)$ -invariant D -subspaces of the orthogonal complement of

$$\{v \in V \mid t(v) = v, \text{ for all } t \in T(k_0)\}.$$

Analogously one shows that the set of maximal k -split tori of $\mathbf{GL}_D(V)$ is one-to-one correspondence with the set of all decompositions of V into one-dimensional D -subspaces.

LEMMA 12.2. — *The map ψ_I from $\mathfrak{B}^1(\prod_i \mathbf{G}_i, k_0)$ to $\mathfrak{B}^1(\mathbf{G}, k_0)$ defined in Proposition 8.6 by*

$$\psi_I((\Lambda_i)_{i \in I_0 \cup I_+}) := \bigoplus_{i \in I_0 \cup I_+} \Lambda_i$$

is toral.

Proof. — For $i \in I_0 \cup I_+$, let S_i be a maximal k_0 -split torus of \mathbf{G}_i and $\{W_l^i \mid l \in L_i\}$ be the corresponding Witt decomposition of $V_i + V_{-i}$. Further let Δ_i be the apartment of S_i in $\mathfrak{B}^1(\mathbf{G}_i, k_0)$. Let α_i be an element of Δ_i seen as a self dual D -norm on $V_i + V_{-i}$. By [8, 2.9] the norm

$$\alpha := \bigoplus_i \alpha_i$$

has the form

$$\alpha\left(\sum_i w_0^i\right) = \frac{1}{2} \inf_i (\nu(h(w_0^i, w_0^i)))$$

on $V_a := \sum_i W_0^i$, i.e. this form does not depend on $(\alpha_i)_i$. Let us take a splitting Witt decomposition $\{W_l^a \mid l \in L_a\}$ of V_a for the restriction of α to V_a . Then the torus $\prod_i S_i$ is a subtorus of the maximal k_0 -split torus T , which corresponds to the Witt decomposition

$$\bigcup_{i \in I_0 \cup I_+ \cup \{a\}} \{W_l^i \mid l \in L_i\}$$

of V , and ψ_I maps $\prod_i \Delta_i$ into the apartment of T . □

PROPOSITION 12.3. — *The map j constructed in the proof of Theorem 7.2 maps apartments into apartments. Furthermore, j is toral if β is separable.*

Without loss of generality we may assume that $I_0 \cup I_+ = \{i\}$ by Lemma 12.2.

Proof $i \in I_+$. — The map \tilde{j}^{E_i} of Theorem 6.7 from $\mathfrak{B}^1(H)$ to

$$\mathfrak{B}^1(\mathbf{GL}_D(V_i), k) = \mathfrak{B}^1(\text{Res}_{k|k_0}(\mathbf{GL}_D(V_i)), k_0)$$

maps apartments into apartments and is toral if β_i is separable by [2, 5.1]. We prove that the canonical map ϕ , see (7.2), from $\mathfrak{B}^1(\text{Res}_{k|k_0}(\mathbf{GL}_D(V_i)), k_0)$ to $\mathfrak{B}^1(\mathbf{G}, k_0)$ defined by

$$\Lambda \in \text{Latt}_{o_D}^1(V_i) \mapsto (\Lambda \oplus \Lambda^{\#-i}) \in \text{Latt}_h^1(V)$$

is toral. A maximal k -split torus S of $\mathbf{GL}_D(V_i)$ corresponds to a decomposition of V_i in one-dimensional D -subspaces, i.e. there is a decomposition $V_i = \bigoplus_l V_{i,l}$ such that

$$S(k) = Z(\{g \in \text{GL}_D(V_i) \mid g(V_{i,l}) \subseteq V_{i,l} \text{ for all } l\}).$$

Let $V_{-i,j}$ be the subspace of V_{-i} dual to $V_{i,j}$, i.e.

$$V_{-i,j} := \{v \in V_{-i} \mid h(v, V_{i,k}) = \{0\}, k \neq j\},$$

and let T be the torus given by the decomposition

$$V = \bigoplus_l (V_{i,l} \oplus V_{-i,l}).$$

Under the map (7.1), from $\text{Aut}_D(V_i)$ to $\text{Aut}_D(V_i) \oplus \text{Aut}_D(V_{-i})$ defined by

$$g \in \text{Aut}_D(V_i) \mapsto (g, 0) + \sigma((g^{-1}, 0)) \in \text{Aut}_D(V_i) \oplus \text{Aut}_D(V_{-i})$$

the set $S(k)$ is mapped into $T(k)$ and under ϕ the apartment of S is mapped into the apartment of T . Let S' and T' be the maximal k_0 -split sub-tori

of $\text{Res}_{k|k_0}(S)$ and $\text{Res}_{k|k_0}(T)$ respectively. The set $S'(k_0)$ is mapped into $(T' \cap \mathbf{G})(k_0)$ under (7.1). The image of ϕ only consists of self-dual lattice functions. Hence ϕ seen as a map from $\mathfrak{B}^1(\text{Res}_{k|k_0}(\mathbf{GL}_D(V_i)), k_0)$ to $\mathfrak{B}^1(\mathbf{G}, k_0)$ is toral. \square

From now on we assume $i \in I_0$. Here we need the notion of tori adapted to a decomposition of V .

DEFINITION 12.4. — Assume we are given a decomposition

$$V = V'^+ \oplus V'^- \oplus V'^0$$

such that V'^+ and V'^- are maximal totally isotropic and $V'^+ \oplus V'^-$ is orthogonal to V'^0 with respect to h . A maximal k_0 -split torus T of \mathbf{G} is adapted to (V'^+, V'^-, V'^0) if there is a Witt decomposition $\{W_l \mid l \in L\}$ corresponding to T with anisotropic part V'^0 such that

$$\bigoplus_l (W_l \cap V'^+) = V'^+ \text{ and } \bigoplus_l (W_l \cap V'^-) = V'^-.$$

We say that an apartment of $\mathfrak{B}^1(G)$ is adapted to (V'^+, V'^-, V'^0) if every lattice function in this apartment is split by (V'^+, V'^-, V'^0) .

Proof $i \in I_0$. — We have $E = E_i$. There are a central division algebra Δ over E and a finite-dimensional right vector space W such that $\text{End}_{E \otimes_k D}(V)$ is E -algebra isomorphic to $\text{End}_\Delta(W)$. We identify the E -algebras $\text{End}_{E \otimes_k D}(V)$ and $\text{End}_\Delta(W)$ via a fixed isomorphism and we fix a signed hermitian form h_E which corresponds to the restriction σ_E of σ to the E -algebra $\text{End}_\Delta(W)$. Let r be the Witt index of h_E . We fix a decomposition

$$(12.1) \quad W = (W^+ \oplus W^-) \oplus W^0$$

such that W^+ and W^- are maximal isotropic subspaces of W contained in the orthogonal complement of W^0 . Let e_+, e_- and e_0 be the projections to the vector spaces W^+, W^- and W^0 via the direct sum (12.1). We define

$$V^+ := e^+V, \quad V^- := e^-V \text{ and } V^0 := e^0V.$$

Consider the following diagram.

$$\begin{array}{ccccc} \mathfrak{B}^1(H) & \longleftarrow & \mathfrak{B}^1(\mathbf{U}((h_E)|_{W^0 \times W^0}), E_0) \times \mathfrak{B}^1(\mathbf{GL}_\Delta(W^+), E) & \longrightarrow & \mathfrak{B}(\mathbf{GL}_\Delta(W^+), E) \\ \downarrow j & & \downarrow \alpha & & \downarrow \\ \mathfrak{B}^1(G) & \longleftarrow & \mathfrak{B}^1(\mathbf{U}(h|_{V^0 \times V^0}), k_0) \times \mathfrak{B}^1(\mathbf{GL}_D(V^+), k) & \longrightarrow & \mathfrak{B}(\mathbf{GL}_D(V^+), k) \end{array}$$

where the vertical maps are induced by j . The right horizontal maps map a pair (x, Λ) to the class of Λ . The right vertical arrow satisfies the CLF-property and its image only consists of E^\times -fixed points of $\mathfrak{B}(\mathbf{GL}_D(V^+), k)$, both properties inherited from j . Thus the map in the right column is j^E ,

i.e. the inverse of j_E , because otherwise we could construct a CLF-map from $\mathfrak{B}(\mathbf{GL}_D(V^+), k)^{E^\times}$ to $\mathfrak{B}(\mathbf{GL}_\Delta(W^+), E)$ different from j_E , but such a CLF-map is unique by [2, II.1.1.]. Now j^E maps apartments into apartments which implies that j maps apartments adapted to (W^+, W^-, W^0) into apartments adapted to (V^+, V^-, V^0) .

We now prove that j is toral if $E|k$ is separable. Let us assume that $E|k$ is separable. This implies that the right column j^E is toral by [2, 5.1] which implies the torality of α because the only maximal E_0 -split torus of the anisotropic group $\mathbf{U}((h_E)|_{W^0 \times W^0})$ is the trivial group. The torality of α implies the torality of j on tori adapted to (W^+, W^-, W^0) . Hence j is toral because the triple (W^+, W^-, W^0) was chosen arbitrarily. \square

Appendix A. The centralizer of a separable Lie algebra element

We prove the representation of the centralizer as a product of general linear groups and Weil restrictions of unitary groups if β is separable. We still rely on the notation from section 2. For more details about the Weil restriction we recommend [23, 1.3.] and [10, 20.5.]. The definitions of D'_i and V'_i are given after notation 5.1.

PROPOSITION A.1. — *Let us assume that β is separable. The centralizer $Z_{\mathbf{G}}(\beta)$ is k_0 -isomorphic to*

$$\prod_{i \in I_0} \text{Res}_{(E_i)_0|k_0}(\mathbf{U}(\sigma|_{\text{End}_{E_i \otimes_k D}(V_i)})) \times \prod_{i \in I_+} \text{Res}_{(E_i)_0|k_0}(\mathbf{GL}_{D'_i}(V'_i));$$

in particular it is reductive and defined over k_0 .

Before we come to the proof, we need some preparations on restriction of scalars.

LEMMA A.2. — *Let D' be a skew field of finite index such that the center, denoted by E , is a non-Archimedean local field. Let V' be a finite-dimensional right D' -vector space. Assume further that σ' is an involution on $\text{End}_{D'}(V')$ whose set of fixed points in E is E_0 . Let k_0 be a subfield of E_0 such that $E_0|k_0$ is separable and finite. Then $\mathbf{U}(\sigma' \otimes_{k_0} \text{id}_\Omega)$ is k_0 -isomorphic to $\text{Res}_{E_0|k_0}(\mathbf{U}(\sigma'))$, for any algebraic closure Ω of E .*

Remark A.3. — If \mathbf{V} is an affine variety defined over E_0 there is a $\text{Gal}(\Omega|k_0)$ -equivariant isomorphism

$$\text{Res}_{E_0|k_0}(\mathbf{V}) \cong \prod_{\gamma} \mathbf{V}^{\gamma},$$

defined over the normal hull of $E_0|k_0$, where γ passes through the set Γ of all field embeddings $E_0 \hookrightarrow \Omega$ fixing k_0 , and one obtains \mathbf{V}^γ from \mathbf{V} in the following way. We choose an automorphism $\bar{\gamma}$ of Ω whose restriction to E_0 is γ . Let P_1, \dots, P_l be polynomials in t variables with coefficients in E_0 such that

$$\mathbf{V} := \{x \in \Omega^t \mid P_i(x) = 0, i = 1, \dots, l\},$$

for some $t \in \mathbb{N}$. One defines

$$\mathbf{V}^\gamma := \{(\bar{\gamma}(x_j))_{1 \leq j \leq t} \mid x \in \mathbf{V}\} = \{x \in \Omega^t \mid P_i^\gamma(x) = 0, i = 1, \dots, l\},$$

where P_i^γ is the polynomial obtained from P_i by applying γ to the coefficients.

Proof of Lemma A.2. — The Ω -algebra $\text{End}_{D'}(V') \otimes_{k_0} \Omega$ is canonically isomorphic to $\text{End}_{D'}(V') \otimes_{E_0} E_0 \otimes_{k_0} \Omega$ and $E_0 \otimes_{k_0} \Omega$ is isomorphic to $\Omega^{[E_0 : k_0]}$ via

$$e \otimes_{k_0} w \mapsto (\gamma(e)w)_{\gamma \in \Gamma}.$$

We denote by Ω^γ the left E_0 -vector space Ω under the action

$$e \cdot \omega := \gamma(e)\omega.$$

Thus

$$\text{End}_{D'}(V') \otimes_{E_0} E_0 \otimes_{k_0} \Omega \cong \prod_{\gamma} \text{End}_{D'}(V') \otimes_{E_0} \Omega^\gamma.$$

The fact that σ' fixes E_0 implies that $\sigma' \otimes_{k_0} \text{id}_\Omega$ is the product of involutions $\sigma' \otimes_{E_0} \text{id}_{\Omega^\gamma}$. It is enough to prove

$$\mathbf{U}(\sigma' \otimes_{E_0} \text{id}_{\Omega^\gamma}) = \mathbf{U}(\sigma' \otimes_{E_0} \text{id}_\Omega)^\gamma.$$

To show the latter, we fix γ and we choose an extension $\bar{\gamma}$ to Ω . The ring isomorphism

$$\Phi: \text{End}_{D'}(V') \otimes_{E_0} \Omega \rightarrow \text{End}_{D'}(V') \otimes_{E_0} \Omega^\gamma$$

sending $f \otimes_{E_0} \omega$ to $f \otimes_{E_0} \bar{\gamma}(\omega)$ satisfies

$$\Phi \circ (\sigma' \otimes_{E_0} \text{id}_\Omega) = (\sigma' \otimes_{E_0} \text{id}_{\Omega^\gamma}) \circ \Phi.$$

Thus an element g of $\text{End}_{D'}(V') \otimes_{E_0} \Omega$ lies in $\mathbf{U}(\sigma' \otimes_{E_0} \text{id}_\Omega)$ if and only if $\Phi(g)$ lies in $\mathbf{U}(\sigma' \otimes_{E_0} \text{id}_{\Omega^\gamma})$, which proves the lemma. \square

Proof of Proposition A.1. — Without loss of generality we assume that $I_0 \cup I_+ = \{i\}$.

Case 1: $I_0 = \{i\}$. We fix an algebraic closure Ω of E . We have

$$\begin{aligned} Z_{\text{End}_D(V) \otimes_{k_0} \Omega}(\beta \otimes_{k_0} 1) &\cong Z_{\text{End}_D(V)}(\beta) \otimes_{k_0} \Omega \\ &\cong \text{End}_{D'_i}(V'_i) \otimes_{k_0} \Omega. \end{aligned}$$

The involution defining $\mathbf{U}(\sigma)$ on $\text{End}_D(V) \otimes_{k_0} \Omega$ is $\sigma \otimes_{k_0} \text{id}$. The above Ω -algebra isomorphism defines an involution $\sigma'_i \otimes_{k_0} \text{id}$ on the right side of the equation where σ'_i is an involution on $\text{End}_{D'_i}(V'_i)$ whose set of fixed points in E is E_0 . By Lemma A.2 the group $\mathbf{U}(\sigma'_i \otimes_{k_0} \text{id})$ is the Weil restriction of $\mathbf{U}(\sigma'_i)$ from E_0 to k_0 .

Case 2: $I_+ = \{i\}$. The algebra E is a direct product of two fields E_i and E_{-i} and we define E_0 to be the set of fixed points of σ in E . In the following part of the proof we use the bijections

$$E_i \rightarrow E_0 \rightarrow E_{-i}, \quad e_i \mapsto e_i + \sigma(e_i) \mapsto \sigma(e_i).$$

We fix an algebraic closure Ω of E_0 . As in the proofs of Lemma A.2 and case 1, we get the product decomposition

$$\begin{aligned} Z_{\text{End}_D(V) \otimes_{k_0} \Omega}(\beta \otimes_{k_0} 1) &\cong Z_{\text{End}_D(V)}(\beta) \otimes_{k_0} \Omega \\ &\cong (\text{End}_{D'_i}(V'_i) \oplus (\text{End}_{D'_i}(V'_i))^{op}) \otimes_{k_0} \Omega \\ &\cong \prod_{\gamma} ((\text{End}_{D'_i}(V'_i) \oplus (\text{End}_{D'_i}(V'_i))^{op}) \otimes_{E_0} \Omega^{\gamma}) \\ &\cong \prod_{\gamma} ((\text{End}_{D'_i}(V'_i) \otimes_{E_i} \Omega^{\gamma}) \oplus (\text{End}_{D'_i}(V'_i) \otimes_{E_{-i}} \Omega^{\gamma})^{op}). \end{aligned}$$

The unitary group

$$\left\{ g \in (\text{End}_{D'_i}(V'_i) \otimes_{E_i} \Omega) \oplus (\text{End}_{D'_{-i}}(V'_{-i}) \otimes_{E_{-i}} \Omega)^{op} \mid g(\sigma \otimes_{E_0} \text{id}_{\Omega})(g) = 1 \right\}$$

is E_i -isomorphic to $\mathbf{GL}_{D'_i}(V'_i)$. We conclude as in the proof of Lemma A.2. □

Appendix B. Rational characters

In this section we show that the extended and the non-extended buildings of G are equal in almost all cases. The following statement is common knowledge, but we give a prove for the sake of completeness. Let $X^*(?)_{k_0}$ denote the set of k_0 -rational characters of $?$. Let us recall that by definition $\mathfrak{B}^1(G)$ is different from $\mathfrak{B}(G)$ if and only if the set $X^*(\mathbf{G}^0)_{k_0}$ is non-trivial (see [6, 4.2.16]).

DEFINITION B.1. — *The special unitary group $\mathbf{SU}(h)$ of h is a connected reductive group defined over k_0 , whose set of rational points, which*

we denote by $SU(h)$, is the intersection of $U(h)(k_0)$ with the kernel of the reduced norm on $\text{End}_D(V)$.

Remark B.2 ([14], 2.15).

- (1) If σ is a unitary involution, then \mathbf{G} is a k -form of $\mathbf{GL}_{md}(\bar{k})$.
- (2) If σ is symplectic, then \mathbf{G} is k -isomorphic to $\mathbf{SU}(h)$ which is a k -form of the symplectic group $\mathbf{Sp}_{md}(\bar{k})$, in particular \mathbf{G} is semisimple.
- (3) If σ is orthogonal, then \mathbf{G}^0 is k -isomorphic to $\mathbf{SU}(h)$ which is a k -form of the special orthogonal group $\mathbf{SO}_{md}(\bar{k})$, in particular \mathbf{G}^0 is semisimple if $md \neq 2$.

We further need the following theorem.

PROPOSITION B.3. — $X^*(\mathbf{G}^0)_{k_0} \neq 1$ if and only if

- (B.1) $m = 2$ and $d = 1$ and σ is orthogonal and h is isotropic.

LEMMA B.4. — Let $L|L'$ be a field extension and $n \in \mathbb{N}$. Let \bar{L} be an algebraic closure of L . Let ϕ be an L' -algebra monomorphism from $M_n(L)$ into $M_n(\bar{L})$. Then there is an element ψ in $\text{Gal}(\bar{L}|L')$ inducing an L' -algebra automorphism of $M_n(\bar{L})$ via

$$\Psi((a_{ij})_{ij}) := (\psi(a_{ij}))_{ij},$$

such that $\Psi \circ \phi$ is an L -algebra monomorphism from $M_n(L)$ into $M_n(\bar{L})$.

Proof. — The map from $M_n(L') \otimes_{L'} \bar{L}$ to $M_n(\bar{L})$ defined by

$$x \otimes_{L'} y \mapsto \phi(x)y$$

is surjective, and thus $\text{im}(\phi)$ contains a \bar{L} -basis of $M_n(\bar{L})$. In particular $\phi(L)$ is a subset of the center \bar{L} of $M_n(\bar{L})$. Now choose $\psi \in \text{Gal}(\bar{L}|L')$ such that $\psi^{-1}|_L$ equals ϕ . □

B.3. — A semisimple group is perfect and has therefore no characters. By remark B.2 the only cases left are:

- (1) σ is unitary.
- (2) σ is orthogonal and $md = 2$ and not (B.1).
- (3) Situation (B.1).

Case 1: We have $D = k$ by Remark 2.1. There is an isomorphism of \bar{k} -algebras with involution

$$(\text{End}_k(V) \otimes_{k_0} \bar{k}, \sigma \otimes_{k_0} \text{id}_{\bar{k}}) \cong (M_m(\bar{k}) \times M_m(\bar{k}), \tilde{\sigma})$$

with

$$\tilde{\sigma}(B, C) = (C^T, B^T),$$

and we have \bar{k} -isomorphisms

$$\mathbf{G} = \mathrm{U}(\sigma \otimes_{k_0} \mathrm{id}_{\bar{k}}) \cong \mathrm{U}(\bar{\sigma}) \cong \mathrm{GL}_m(\bar{k}).$$

The last isomorphism is induced by the projection to the first coordinate. Let χ be a k_0 -rational character of \mathbf{G} . A character of $\mathrm{GL}_m(\bar{k})$ is a power of the determinant. Thus, because of Lemma B.4, $\chi|_G$ must be a power of $\det|_G$ or $\rho \circ \det|_G$. We now fix a basis of V to get a k -isomorphism from $\mathrm{M}_m(k)$ to $\mathrm{End}_k(V)$. The involution $\sigma \circ ()^\rho$ is conjugate to the transpose map, which implies

$$\chi(x)^{-1} = \chi(\sigma(x)) = \rho(\chi(x)) = \chi(x),$$

for all $x \in G$. The last equality follows from $\chi(G) \subseteq k_0$. In particular the only possible values of χ on G are 1 and -1 . Thus χ is trivial, because \mathbf{G} is connected and G is Zariski-dense in \mathbf{G} by [1, 18.3].

Case 2: We have $k = k_0$, since σ is orthogonal. If $d = 2$ There is an element $a \in \mathrm{SU}(h) \setminus \{1, -1\}$, because $\mathrm{SU}(h)$ is Zariski-dense in $\mathbf{SU}(h)$ by [1, 18.3]. We have that $k[a]$ is its own centralizer in D , because the index of D is two, in particular the commutative group $\mathrm{SU}(h)$ is a subset of $k[a]$. if we introduce a k -basis of D which contains 1 and a , then the identity from $\mathrm{SU}(h)$ to $\mathrm{U}(\sigma|_{k[a]})$ can be extended to a k -isomorphism from $\mathbf{SU}(h)$ to $\mathrm{U}(\sigma|_{k[a]})$. By case 1 there is no k -rational character on $\mathbf{SU}(h)$.

Let us now assume $d = 1$ and $\mathbf{SU}(h)$ is anisotropic. There is a k -basis of V such that the Gram matrix of h is of the form

$$\begin{pmatrix} e & 0 \\ 0 & f \end{pmatrix},$$

and we identify $\mathrm{End}_k(V)$ with $\mathrm{M}_2(k)$. A short calculation shows that

$$\mathbf{SU}(h) = \left\{ \begin{pmatrix} a & cf \\ -ce & a \end{pmatrix} \mid a, c \in \bar{k} \text{ s.t. } a^2 + efc^2 = 1 \right\}.$$

We fix square roots \sqrt{e} and $\sqrt{-f}$. The conjugation with

$$\begin{pmatrix} \sqrt{e} & \sqrt{-f} \\ \frac{\sqrt{e}}{2} & -\frac{\sqrt{-f}}{2} \end{pmatrix}$$

maps $\mathbf{SU}(h)$ to \mathbf{SO}_2^{is} . The explicit formula for the map is

$$\begin{pmatrix} a & cf \\ -ce & a \end{pmatrix} \mapsto \begin{pmatrix} a - c\sqrt{-ef} & 0 \\ 0 & a + c\sqrt{-ef} \end{pmatrix}, \quad \sqrt{-ef} := \sqrt{-e}\sqrt{f}.$$

Thus a rational character of $\mathbf{SU}(h)$ is of the form

$$\begin{pmatrix} a & cf \\ -ce & a \end{pmatrix} \mapsto (a + c\sqrt{-ef})^z,$$

for some integer z . The inverse of $(a + c\sqrt{-ef})$ is $(a - c\sqrt{-ef})$. If z is positive, we apply the binomial expansion to get coefficients α and γ in k such that

$$(a + c\sqrt{-ef})^z = \alpha + \gamma\sqrt{-ef}.$$

The element γ is zero, because $\sqrt{-ef} \notin k$ since h is anisotropic. Thus a k -rational character χ of $\mathbf{SU}(h)$ satisfies

$$\chi(x) = \chi(x^{-1}),$$

for all $x \in \mathbf{SU}(h)$. The density of $\mathbf{SU}(h)$ in $\mathbf{SU}(h)$ and the connectivity of $\mathbf{SU}(h)$ imply that χ is trivial.

Case 3: Here \mathbf{G} is k_0 -isomorphic to \mathbf{O}_2^{is} implying that \mathbf{G}^0 has non-trivial k_0 -rational characters. \square

BIBLIOGRAPHY

- [1] A. BOREL, *Linear algebraic groups*, Grad. Texts in Math., vol. 126, Springer-Verlag, New York, 1991, 2nd enl. ed., x+308 pages.
- [2] P. BROUSSOUS & B. LEMAIRE, “Building of $\mathrm{GL}(m, D)$ and centralizers”, *Transform. Groups* **7** (2002), no. 1, p. 15-50.
- [3] P. BROUSSOUS, V. SÉCHERRE & S. STEVENS, “Smooth representations of $\mathrm{GL}(m, D)$, V: Endo-classes”, arXiv:1004.5032v1, 2010.
- [4] P. BROUSSOUS & S. STEVENS, “Buildings of classical groups and centralizers of Lie algebra elements”, *J. Lie Theory* **19** (2009), no. 1, p. 55-78.
- [5] K. BROWN, *Buildings*, Springer-Verlag, New York, 1989, viii+215 pages.
- [6] F. BRUHAT & J. TITS, “Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d’une donnée radicielle valuée”, *Inst. Hautes Études Sci. Publ. Math.* (1984), no. 60, p. 197-376.
- [7] ———, “Schémas en groupes et immeubles des groupes classiques sur un corps local”, *Bull. Soc. Math. France* **112** (1984), no. 2, p. 259-301.
- [8] ———, “Schémas en groupes et immeubles des groupes classiques sur un corps local. II. Groupes unitaires”, *Bull. Soc. Math. France* **115** (1987), no. 2, p. 141-195.
- [9] C. BUSHNELL & P. KUTZKO, *The admissible dual of $\mathrm{GL}(N)$ via compact open subgroups*, Ann. of Math. Studies, vol. 129, Princeton Univ. Press, Princeton, NJ, 1993, xii+313 pages.
- [10] M.-A. KNUS, A. MERKURJEV, M. ROST & J.-P. TIGNOL, *The book of involutions*, AMS Colloquium Publications, vol. 44, Amer. Math. Soc., Providence, RI, 1998, xxii+593 pages.
- [11] E. LANDVOGT, “Some functorial properties of the Bruhat-Tits building”, *J. Reine Angew. Math.* **518** (2000), p. 213-241.
- [12] B. LEMAIRE, “Comparison of lattice filtrations and Moy-Prasad filtrations for classical groups”, *J. Lie Theory* **19** (2009), no. 1, p. 29-54.
- [13] A. MOY & G. PRASAD, “Unrefined minimal K -types for p -adic groups”, *Invent. Math.* **116** (1994), no. 1-3, p. 393-408.
- [14] V. PLATONOV & A. RAPINCHUK, *Algebraic groups and number theory*, Pure and Appl. Math., vol. 139, Acad. press, Inc., BOSTON, 1994, Transl. from the 1991 Russ. orig. by R. Rowen, xii+614 pages.

- [15] G. PRASAD & J.-K. YU, “On finite group actions on reductive groups and buildings”, *Invent. Math.* **147** (2002), no. 3, p. 545-560.
- [16] W. SCHARLAU, *Quadratic and Hermitian Forms*, Springer-Verlag, Berlin and Heidelberg, 1985, x+421 pages.
- [17] V. SÉCHERRE, “Représentations lisses de $GL(m, D)$. I. Caractères simples”, *Bull. Soc. Math. France* **132** (2004), no. 3, p. 327-396.
- [18] ———, “Représentations lisses de $GL(m, D)$. II. β -extensions”, *Compos. Math.* **141** (2005), no. 6, p. 1531-1550.
- [19] ———, “Représentations lisses de $GL(m, D)$. III. Types simples”, *Ann. Sci. École Norm. Sup. (4)* **38** (2005), no. 6, p. 951-977.
- [20] V. SÉCHERRE & S. STEVENS, “Représentations lisses de $GL_m(D)$. IV. Représentations supercuspidales”, *J. Inst. Math. Jussieu* **7** (2008), no. 3, p. 527-574.
- [21] ———, “Smooth Representations of $GL_m(D)$ VI: Semisimple Types”, *Int. Math. Res. Not.*, 2011.
- [22] S. STEVENS, “Semisimple characters for p-adic classical groups”, *Duke Math. J.* **127** (2005), no. 1, p. 123-173.
- [23] A. WEIL, *Adeles and algebraic groups*, Prog. in Math., vol. 23, Birkhäuser Boston, Mass., 1982, iii+126 pages.

Manuscrit reçu le 28 février 2011,
révisé le 9 août 2011,
accepté le 21 février 2012.

Daniel SKODLERACK
Universität Münster
Mathematisches Institut
Einsteinstrasse 62
48149 Münster (Germany)
skodlerack@uni-muenster.de