



ANNALES

DE

L'INSTITUT FOURIER

Jean RUPPENTHAL & Martin SERA

Modifications of torsion-free coherent analytic sheaves

Tome 67, n° 1 (2017), p. 237-265.

http://aif.cedram.org/item?id=AIF_2017__67_1_237_0



© Association des Annales de l'institut Fourier, 2017,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

MODIFICATIONS OF TORSION-FREE COHERENT ANALYTIC SHEAVES

by Jean RUPPENTHAL & Martin SERA (*)

ABSTRACT. — We study the transformation of torsion-free coherent analytic sheaves under proper modifications. More precisely, we study direct images of inverse image sheaves, and torsion-free preimages of direct image sheaves. Under some conditions, it is shown that torsion-free coherent sheaves can be realized as the direct image of locally free sheaves under modifications. Thus, it is possible to study coherent sheaves modulo torsion by reducing the problem to study vector bundles on manifolds. We apply this to reduced ideal sheaves and to the Grauert–Riemenschneider canonical sheaf of holomorphic n -forms.

RÉSUMÉ. — Nous étudions la transformation de faisceaux analytiques cohérents à travers des modifications propres. Plus précisément, nous étudions les images directes de faisceaux images-inverses, et les préimages sans torsion de faisceaux images directes. Sous certaines conditions, il est démontré que les faisceaux cohérents sans torsion peuvent être réalisés comme images directes de faisceaux localement libres à travers des modifications. Ainsi, il est possible d'étudier les faisceaux cohérents modulo torsion en ramenant le problème à l'étude de fibrés vectoriels sur des variétés. Nous appliquons ceci aux faisceaux d'idéaux réduits et au faisceau canonique de n -formes holomorphes de Grauert–Riemenschneider.

1. Introduction

In bimeromorphic geometry, the use of locally free coherent analytic sheaves is limited: the direct image of a locally free sheaf under a proper

Keywords: modifications, coherent analytic sheaves, torsion-free preimage.

Math. classification: 32C35, 32S45, 32L05, 32L10.

(*) The authors are grateful to Matei Toma and Daniel Greb for discovering a mistake in an earlier version of this paper and for helpful discussions on the topic, and to the unknown referee for several suggestions which helped to improve the paper. We also thank Joël Merker for translating the abstract. This research was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), grant RU 1474/2 within DFG's Emmy Noether Programme.

modification is not locally free any more. Instead, it is reasonable to consider the wider category of torsion-free coherent analytic sheaves. The restriction to torsion-free sheaves makes sense for bimeromorphic considerations as the torsion of a coherent analytic sheaf is supported on analytically thin subsets. To exemplify the use of torsion-free sheaves, just recall that an irreducible⁽¹⁾ reduced compact space X is a Moishezon space if and only if it carries a positive torsion-free coherent analytic sheaf \mathcal{S} with $\text{supp}(\mathcal{S}) = X$ (see e.g. in [15, Thm. 6.14]).

Let $\pi: Y \rightarrow X$ be a proper modification of a complex space X (cf. Section 2 for notations). Then the direct image $\pi_*\mathcal{F}$ of a torsion-free coherent analytic sheaf \mathcal{F} remains torsion-free. The problem is here that the analytic inverse image sheaf $\pi^*\mathcal{S}$ of a torsion-free coherent analytic sheaf \mathcal{S} is not torsion-free in general. For a counter-example, see e.g. the example in [11, §1], i.e., the pullback of the maximal ideal sheaf of the origin in \mathbb{C}^2 under blow-up of the origin is not torsion-free. One can say more or less that $\pi^*\mathcal{S}$ is torsion-free in a point $y \in Y$ if and only if \mathcal{S} is locally free in $\pi(y)$ (see [18] or Remark 3.21 below).

To take care of the torsion which arises when taking analytic inverse images, it is useful to consider the torsion-free preimage sheaf:

DEFINITION 1.1. — *Let $\pi: Y \rightarrow X$ be a holomorphic map between complex spaces such that Y is locally irreducible. Let \mathcal{S} be a coherent analytic sheaf on X . Then*

$$\pi^T\mathcal{S} := \pi^*\mathcal{S} / \mathcal{T}(\pi^*\mathcal{S}),$$

where $\mathcal{T}(\pi^*\mathcal{S})$ is the torsion sheaf of $\pi^*\mathcal{S}$, is called the torsion-free preimage sheaf of \mathcal{S} under π .

Torsion-free preimages under proper modifications have been first studied by H. Rossi [20], H. Grauert and O. Riemenschneider [11, 19]. The main motivation is as follows: Let \mathcal{S} be a torsion-free coherent analytic sheaf on an irreducible complex space X . Then Rossi showed that there exists a proper modification $\varphi_{\mathcal{S}}: Y \rightarrow X$ such that $\varphi_{\mathcal{S}}^T\mathcal{S}$ is locally free (see Section 2 for the details). Combining this with a resolution of singularities $\sigma: M \rightarrow Y$ which exists due to Hironaka, we obtain a resolution of singularities $\pi = \varphi_{\mathcal{S}} \circ \sigma: M \rightarrow X$ such that $\pi^T\mathcal{S}$ is locally free. Thus, it

⁽¹⁾We call a complex space *locally irreducible* if each stalk of its structure sheaf is an integral domain (see e.g. [10, Chap. 1, §1.5]). In particular, it is then automatically reduced. On the other hand, we say that a complex space X is (*globally*) *irreducible* if the underlying reduced space, $\text{red}(X)$, consists of just one irreducible component. If there are more than one components, then X is called *reducible*.

is possible to study coherent analytic sheaves modulo torsion by reducing the problem to study vector bundles on manifolds.

In view of this idea, it seems very interesting to study the connection between \mathcal{S} and its torsion-free preimage $\pi^T \mathcal{S}$ closer, and we have found the following relation which to our knowledge has not been observed in the literature:

THEOREM 1.2. — *Let $\pi: Y \rightarrow X$ be a proper modification of a complex space X , and let \mathcal{F} and \mathcal{G} be torsion-free coherent analytic sheaves on X and Y , respectively.*

(1) *If $\mathcal{F} = \pi_* \mathcal{G}$, then*

$$\mathcal{F} \cong \pi_* \pi^T \mathcal{F}.$$

(2) *If $\mathcal{G} = \pi^T \mathcal{F}$, then*

$$\pi^T \pi_* \mathcal{G} \cong \mathcal{G}.$$

Theorem 1.2 can be shown directly by standard facts on modifications and torsion-free sheaves. But we obtain it here as a simple byproduct of considerations on the functor π^T (see Lemma 6.1) and the fact that the natural maps

$$\begin{aligned} \mathcal{F} &\rightarrow \pi_* \pi^T \mathcal{F}, \\ \pi^T \pi_* \mathcal{G} &\rightarrow \mathcal{G} \end{aligned}$$

both are injective (Theorem 4.1(i) and Lemma 5.1; the proof of Lemma 5.1 presented here is due to Matei Toma). We give also counter-examples to show that these injections are not bijective in general (Remark 4.2 and Remark 5.2).

In order to understand relations as in Theorem 1.2 better, it turns out useful to study properties of linear spaces associated to coherent analytic sheaves. Among other things, we obtain the following equivalence:

THEOREM 1.3. — *Let X be a connected factorial Cohen–Macaulay space and \mathcal{S} a coherent analytic sheaf on X generated by $\text{rk } \mathcal{S} + m$ sections, $m \leq 2$, such that the singular locus of \mathcal{S} is at least of codimension $m+1$ in X . Then the following conditions are equivalent:*

- (1) \mathcal{S} is torsion-free.
- (2) The linear (fiber) space $L(\mathcal{S})$ associated to \mathcal{S} is globally irreducible and reduced (i.e., it consists only of its primary component).
- (3) $L(\mathcal{S})$ is locally irreducible.

(4) For all $p \in X$, there is a neighborhood $U \subset X$ such that

$$0 \rightarrow \mathcal{O}_U^m \rightarrow \mathcal{O}_U^{\text{rk } \mathcal{S} + m} \rightarrow \mathcal{S}_U \rightarrow 0$$

is exact, i.e., the homological dimension of \mathcal{S} is at most one.

R. Axelsson and J. Magnússon proved in [2, Prop. 3.11] that a condition on the codimension of the singular locus of \mathcal{S} is necessary to obtain the irreducibility of $L(\mathcal{S})$. In this context, we can highly recommend their papers [1] and [2], which deal with complex analytic cones, a generalization of complex linear spaces.

If, moreover, the singular locus of \mathcal{S} is at least $(m+2)$ -codimensional, then $L(\mathcal{S})$ is normal (see Theorem 3.18). This is a particularly interesting situation because of the following statement.

THEOREM 1.4. — *Let X be a locally irreducible complex space, \mathcal{S} a torsion-free coherent analytic sheaf on X such that the linear space associated to \mathcal{S} is normal, and $\pi: Y \rightarrow X$ a proper modification of X . Then the canonical homomorphism $\mathcal{S} \rightarrow \pi_*(\pi^T \mathcal{S})$ is bijective, i.e.,*

$$\mathcal{S} \cong \pi_*(\pi^T \mathcal{S}).$$

In this situation, \mathcal{S} can actually be represented as the direct image of a locally free sheaf.

Let us mention two applications of Theorem 1.2. First, we will study ideal sheaves. Let $\pi: Y \rightarrow X$ be a proper modification of a locally irreducible complex space X , $A \subset X$ an analytic subset with ideal sheaf \mathcal{I}_A , and $B := \pi^{-1}(A)$ the analytic preimage with ideal sheaf \mathcal{I}_B . Then $\pi^T \mathcal{I}_A = \mathcal{I}_B$ (cf. Lemma 7.2), and Theorem 1.2 yields that $\mathcal{I}_B \cong \pi^T \pi_* \mathcal{I}_B$.

If we assume moreover that X is normal and that A is either a locally complete intersection or a normal analytic set and that $\sigma: Y \rightarrow X$ is the monoidal transformation with respect to \mathcal{I}_A , then $\mathcal{I}_B = \sigma^T \mathcal{I}_A$ (is locally free) and we have (see Lemma 7.3):

$$\mathcal{I}_A \cong \sigma_* \mathcal{I}_B \cong \sigma_* \sigma^T \mathcal{I}_A.$$

Second, let X be a locally irreducible complex space of pure dimension n , and \mathcal{K}_X the Grauert–Riemenschneider canonical sheaf on X (as introduced in [11]). Then there exists a resolution of singularities $\pi: M \rightarrow X$ (with only normal crossings) such that $\pi^T \mathcal{K}_X$ is locally free, and so there is an effective divisor D with support on the exceptional set of the modification such that

$$(1.1) \quad \mathcal{K}_X \cong \pi_* \pi^T \mathcal{K}_X = \pi_* \Omega_M^n(-D) = \pi_*(\Omega_M^n \otimes \mathcal{O}(-D))$$

(see Theorem 8.1). Let us explain briefly the meaning of (1.1). By definition of the Grauert–Riemenschneider canonical sheaf, we know already that $\mathcal{K}_X \cong \pi_*\Omega_M^n$. (1.1) tells us that we can as well consider the push-forward of holomorphic n -forms which vanish to the order of D on the exceptional set. This is a useful information, particularly if π is explicitly given so that D can be calculated explicitly. An example: If X is already a manifold (i.e., $\mathcal{K}_X = \Omega_X^n$) and $\pi: M \rightarrow X$ is the blow-up along a submanifold of codimension s in X with exceptional set E , then (see e.g. [4, Prop. VII.12.7]):

$$\pi^T \mathcal{K}_X = \pi^* \mathcal{K}_X = \Omega_M^n(- (s - 1)E),$$

and so

$$\Omega_X^n = \mathcal{K}_X \cong \pi_*\Omega_M^n(- (s - 1)E).$$

Considerations of this kind are particularly important in the study of canonical sheaves on singular complex spaces (see [21]). We will set up the relation (1.1) also for holomorphic n -forms with values in locally free coherent analytic sheaves (see Theorem 8.3).

Using Theorem 1.4, we are able to generalize Takegoshi’s relative version [23] of the Grauert–Riemenschneider vanishing theorem in several directions. This is elaborated by the second author in [22].

The content of the present paper is organized as follows. After a brief review of monoidal transformations with respect to coherent analytic sheaves in Section 2, we will study linear spaces associated to torsion-free coherent analytic sheaves in Section 3. There, we will prove Theorem 1.3. Then we study direct images of (torsion-free) analytic preimage sheaves (including the proof of Theorem 1.4) in Section 4, and torsion-free analytic preimages of direct image sheaves in Section 5. In Section 6, we show that the analytic inverse image functor preserves monomorphisms and epimorphisms and use this fact in combination with the previous considerations to prove Theorem 1.2. Section 7 and Section 8 contain the applications described above. We complement the paper by analogous considerations on the non-analytic inverse image functor in Section 9.

2. Monoidal transformations

Let us recall some preliminaries on monoidal transformations of complex spaces with respect to coherent analytic sheaves.

DEFINITION 2.1. — A proper surjective holomorphic map $\varphi: X \rightarrow Y$ of complex spaces X and Y is called a (proper) modification if there are closed analytic sets $A \subset X$ and $B \subset Y$ such that

- (1) $B = \varphi(A)$,
- (2) $\varphi|_{X \setminus A}: X \setminus A \rightarrow Y \setminus B$ is biholomorphic,
- (3) A and B are analytically rare, and
- (4) A and B are minimal with the properties (1–3).

A is called the exceptional set of φ and B the center of the modification.

Rossi showed in [20] that coherent analytic sheaves can be made locally free by use of modifications. This process has been treated more systematically by Riemenschneider [19]. Following [19, §2], we define:

DEFINITION 2.2. — Let X be a complex space and \mathcal{S} a coherent analytic sheaf on X . Then a pair $(X_{\mathcal{S}}, \varphi_{\mathcal{S}})$ of a complex space $X_{\mathcal{S}}$ and a proper modification $\varphi_{\mathcal{S}}: X_{\mathcal{S}} \rightarrow X$ is called the monoidal transformation of X with respect to \mathcal{S} if the following two conditions are fulfilled:

- (1) the torsion-free preimage $\varphi_{\mathcal{S}}^T \mathcal{S} = \varphi_{\mathcal{S}}^* \mathcal{S} / \mathcal{T}(\varphi_{\mathcal{S}}^* \mathcal{S})$ is locally free on $X_{\mathcal{S}}$,
- (2) if $\pi: Y \rightarrow X$ is any proper modification such that (1) holds then there is a unique holomorphic mapping $\psi: Y \rightarrow X_{\mathcal{S}}$ such that $\pi = \varphi_{\mathcal{S}} \circ \psi$.

So, if $X_{\mathcal{S}}$ exists, it is uniquely determined up to biholomorphism by (2). But existence was first proven by Rossi (see Thm. 3.5 in [20]) and then studied further by Riemenschneider (see Thm. 2 in [19]):

THEOREM 2.3. — Let X be an (irreducible) complex space, \mathcal{S} a coherent analytic sheaf on X and $A = \text{Sing } \mathcal{S} := \{x \in X: \mathcal{S} \text{ is not locally free at } x\}$ the singular locus of \mathcal{S} . Then there exists the monoidal transformation $(X_{\mathcal{S}}, \varphi_{\mathcal{S}})$ of X with respect to \mathcal{S} . $X_{\mathcal{S}}$ is a reduced (irreducible) complex space and $\varphi_{\mathcal{S}}$ is a projective proper modification such that

$$\varphi_{\mathcal{S}}: X_{\mathcal{S}} \setminus \varphi_{\mathcal{S}}^{-1}(A) \rightarrow X \setminus A$$

is biholomorphic. If $U \subset X$ is an open subset, then $(\varphi_{\mathcal{S}}^{-1}(U), \varphi_{\mathcal{S}})$ is the monoidal transformation of U with respect to \mathcal{S}_U .

3. Linear spaces of torsion-free coherent analytic sheaves

For a coherent analytic sheaf \mathcal{S} on a complex space X , we work with the linear (fiber) space $S := L(\mathcal{S})$ associated to \mathcal{S} (in the sense of Fischer [6, 7])

and Grothendieck [12]). Note that if \mathcal{E} is a locally free sheaf, then the linear space $L(\mathcal{E})$ is the dual of the vector bundle which has \mathcal{E} as sheaf of sections.

Linear fiber spaces are a special case of *complex analytic cones*, introduced by R. Axelsson and J. Magnússon in [1]. Their further study of complex analytic cones in [2] offers particularly also new, clarifying insights into the theory of linear spaces.

3.1. Primary component of a linear space

In the following, we will always assume that X is a locally irreducible complex space. Thus, X decomposes into disjoint connected components which can be considered separately (see e.g. [10, Chap. 9 §2]). So, we can assume that X is connected, thus also globally irreducible. For a coherent analytic sheaf \mathcal{S} , let A be the thin analytic set in X where \mathcal{S} is not locally free (see [9]). We call $\text{Sing } S = \text{Sing } \mathcal{S} := A$ the *singular locus* of $S = L(\mathcal{S})$ and \mathcal{S} , respectively. As X is irreducible, $X' := X \setminus (A \cup X_{\text{sing}})$ and $A^c := X \setminus A$ are connected. $S_U \cong U \times \mathbb{C}^r$, for small open sets $U \subset A^c$, implies that S_{A^c} is also connected. The set S_A is an analytic subset of S . Let E be the irreducible component of $\text{red}(S)$ which contains S_{A^c} . $\text{PC}(S) := E$ is called the *primary component* of S (following the notation of [18]). We have the decomposition $S = E \cup S_A$.

Remark 3.1. — Let S be a linear space associated to a coherent analytic sheaf \mathcal{S} . Let $s \in \text{Hom}(S_U, U \times \mathbb{C}) \cong \mathcal{S}(U)$ be a section. Then the primary component E of S determines s up to torsion, i.e., if $s|_E = 0$, then $s \in \mathcal{T}(\mathcal{S})$. This is clear as $s|_E = 0$ implies that s is supported only on an analytically thin set.

The following observation can be deduced easily from the definition of the primary component. It is also an immediate consequence of Thm. 3.10 in [2], so that we do not need to include a proof here:

LEMMA 3.2. — *Let X be a locally irreducible complex space and \mathcal{S} a coherent analytic sheaf on X . Let $S = L(\mathcal{S})$ be the linear space associated to \mathcal{S} and E its primary component. If \mathcal{S} has a torsion element, then $E \neq S$. In particular, S is reducible or non-reduced.*

Obviously, a linear space can be reducible but reduced (for instance, $\{xz = 0\} \subset \mathbb{C}_x \times \mathbb{C}_z$). The case that a linear space is irreducible and non-reduced can occur as well: Let $S \subset \mathbb{C}_{x,y}^2 \times \mathbb{C}_{z,w}^2$ be the linear space given by the ideal sheaf generated by $h_1(x, y; z, w) := y^2z - xyw$ and

$h_2(x, y, z, w) := xyz - x^2w$. Then the primary component $E := \text{PC}(S)$ is given by $g(x, y, z, w) := yz - xw$, i.e., $E = \text{red } S$ and S is irreducible. Yet, g does not vanish on S (in the unreduced sense) while $g^2 = yzg - xwg = zh_1 - wh_2$ does.

Remark 3.3. — The converse of Lemma 3.2 is not true:

Let \mathcal{I} be the ideal sheaf generated by x^2, xy, y^2 on $\mathbb{C}_{x,y}^2$ and $S := L(\mathcal{I})$ the linear space associated to \mathcal{I} . Since \mathcal{I} can not be generated by 2 elements, we get $\text{rk } S_0 = \text{rk } \mathcal{I}_0 = 3$. Hence, S_0 is a 3-dimensional analytic subset of S . On the other hand, the primary component has dimension $2 + \text{rk } S = 3$. Hence, S is not irreducible. More precisely, S is given in $\mathbb{C}_{x,y}^2 \times \mathbb{C}_z^3$ by the ideal sheaf generated by $h_1(x, y, z) := yz_1 - xz_2$ and $h_2(x, y, z) := yz_2 - xz_3$ where $z = (z_1, z_2, z_3)$. Since $y(z_2^2 - z_1z_3) = z_2h_2$, the primary component $\text{PC}(S)$ is defined by the functions h_1, h_2 and $z_2^2 - z_1z_3$. This shows also that the fibers of $\text{PC}(S)$ are not linear (the fiber over the origin is just $\{z_2^2 = z_1z_3\}$). So, the primary component is in general not a linear space, neither in the sense of Fischer [6, 7] nor in the sense of Grauert [9].⁽²⁾

Actually, we see that S is the linear space associated to a torsion-free sheaf, and it is reduced but not irreducible. Considering analogously the ideal sheaf given by x^2, xy^2, y^4 , then it turns out that the associated linear space is neither irreducible nor reduced.

In general, for an ideal sheaf \mathcal{I} on a reduced complex space X , the primary component of the associated linear space is given by the analytic spectrum of the Rees algebra of \mathcal{I} (see [2, Exp. 3.12(1)]):

$$\text{PC}(L(\mathcal{I})) \cong \text{Specan} \left(\bigoplus_{m \geq 0} \mathcal{I}^m \right).$$

In particular, $S(\mathcal{I}) \cong \bigoplus_{m \geq 0} \mathcal{I}^m$ implies that $L(\mathcal{I}) \cong \text{PC}(L(\mathcal{I}))$ (e.g. if \mathcal{I} is generated by an \mathcal{O}_X -regular sequence or by two elements, see [3, §3] and [13, (8) and Thm. 3.1] resp.).

Using Rossi's monoidal transformation, we can make the following observation about the primary component:

⁽²⁾ In contrast to Fischer's notion of a linear space, where it is required that $+: S \times_X S \rightarrow S$ is a holomorphic map, Grauert requires in [9] (only) that the addition $+: S \oplus_X S \rightarrow S$ is holomorphic. That gives a different category of linear spaces (which is no longer dually equivalent to the category of coherent analytic sheaves). In [17, p. 238], Rabinowitz claims that the primary component of a linear space is a linear space in the sense of Grauert [9], but not in the sense of Fischer [6, 7]. Our example shows that even this is not the case. More details and criteria for a Grauert linear space to be linear in Fischer's sense can be found in [2, §3.3].

THEOREM 3.4. — *Let X be a locally irreducible complex space and \mathcal{S} a coherent analytic sheaf on X . Then the primary component E of the linear space S associated to \mathcal{S} is locally irreducible.*

Proof. — As above, we can assume that X is connected, i.e., irreducible. Let

$$\varphi := \varphi_{\mathcal{S}}: X_{\mathcal{S}} \rightarrow X$$

be the monoidal transformation of X with respect to \mathcal{S} . This implies that φ is biholomorphic on $X_{\mathcal{S}} \setminus \varphi^{-1}(A)$, where $A \subset X$ is the thin analytic subset where \mathcal{S} is not locally free. Then

$$\varphi^*S = X_{\mathcal{S}} \times_X S$$

is the linear space associated to $\varphi^*\mathcal{S}$, and there is a proper holomorphic projection

$$\text{pr}: \varphi^*S \rightarrow S.$$

Now consider the natural surjective homomorphism

$$\varphi^*\mathcal{S} \longrightarrow \varphi^T\mathcal{S} = \varphi^*\mathcal{S} / \mathcal{I}(\varphi^*\mathcal{S})$$

which induces a closed embedding of the linear space $V := L(\varphi^T\mathcal{S})$ into φ^*S . Note that V coincides with φ^*S on $X_{\mathcal{S}} \setminus \varphi^{-1}(A)$. Thus, the vector bundle V is just the primary component of φ^*S , and V is clearly locally and globally irreducible (because the base space $X_{\mathcal{S}}$ is connected and locally irreducible).

As pr is a proper holomorphic mapping, we have that $\text{pr}(V)$ is an irreducible analytic subset of S by Remmert’s proper mapping theorem and the fact that holomorphic images of irreducible sets are again irreducible (see [10, Chap. 9, §1.3]). But $\text{pr}(V)$ coincides with E , the primary component of S , over $X \setminus A$. Thus:

$$(3.1) \quad \text{pr}(V) = E,$$

and so $\text{pr}|_V: V \rightarrow E$ is a proper modification. Using this and the fact that V is clearly locally irreducible, it is easy to see that E is also locally irreducible: For an open connected set $W \subset E$, $\text{pr}|_V^{-1}(W) \subset V$ is again open and connected, thus irreducible since φ is a proper modification of the irreducible X . But then $W = \text{pr}|_V(\text{pr}|_V^{-1}(W))$ is also irreducible by the same argument as above (holomorphic images of irreducible sets are irreducible). □

LEMMA 3.5. — *Let X be a locally irreducible complex space and \mathcal{S} a torsion-free coherent analytic sheaf on X . Then $S = L(\mathcal{S})$ is locally irreducible if and only if the primary component of S is a linear space.*

Proof. — Let $E \subset S$ denote the primary component of S . As \mathcal{S} is torsion-free, [2, Thm. 3.10] implies that $E = S$ if and only if the primary component E is a linear space. Alternatively, this assertion can be proven by elementary computations with [7, Lem. 1]. The statement of the lemma follows now with Theorem 3.4. \square

As we have seen in the counter-example Remark 3.3, the primary component need not be a linear space (even in the sense of Grauert). Though, it appears as the analytic spectrum of a connected graded \mathcal{O}_X -algebra of finite presentation. So, it is locally isomorphic to a subcone of a trivial linear space defined by quasihomogeneous polynomials (see [1, Cor. 1.13]). Since a symmetric algebra and its torsion-free reduction is generated by elements of degree one, R. Axelsson and J. Magnússon actually showed in the proof of [1, Cor. 1.13] also the following:

LEMMA 3.6. — *Let X be a (locally irreducible) complex space and \mathcal{S} a coherent analytic sheaf on X . Then the primary component E of the linear space $S = L(\mathcal{S})$ associated to \mathcal{S} is fiberwise homogeneous and E is locally defined as analytic set in $U \times \mathbb{C}^N$ by holomorphic fiberwise homogeneous functions for $U \subset X$ small enough.*

If $E \times_X E$ is locally irreducible, one can prove that the primary component E is a linear space. Yet, for an irreducible fiber space $E \rightarrow X$, the fiber product of $E \times_X E$ need not be reduced (not to mention locally irreducible; for a counter-example, see [6, §4]). Therefore, the restriction of the addition need not be holomorphic.

3.2. Linear spaces of small corank – Proof of Theorem 1.3

For a linear space to be (locally) irreducible, it is necessary that the associated coherent analytic sheaf is torsion-free. In the following, we will prove that this is a sufficient criterion under certain additional assumptions.

DEFINITION 3.7. — *For a coherent analytic sheaf \mathcal{S} on a complex space X , we define the corank of \mathcal{S} in a point $p \in X$, $\text{cork}_p \mathcal{S}$, as the difference of the minimal number of generators of \mathcal{S}_p and the rank of \mathcal{S} , and the global corank $\text{cork} \mathcal{S} := \sup_{p \in X} \text{cork}_p \mathcal{S}$. The corank of a linear space is defined as the corank of the associated coherent analytic sheaf.*

We get the following relation between the corank and the homological dimension of a coherent analytic sheaf:

LEMMA 3.8. — *Let X be a complex space and \mathcal{S} a coherent analytic sheaf on X . Then, for all $p \in X$, the following is equivalent:*

- (1) *There exists a neighborhood U of p such that the following sequence is exact:*

$$\mathcal{O}_U^{\text{cork}_p \mathcal{S}} \xrightarrow{\alpha} \mathcal{O}_U^{\text{rk} \mathcal{S} + \text{cork}_p \mathcal{S}} \rightarrow \mathcal{S}_U \rightarrow 0.$$

- (2) *The homological dimension of \mathcal{S} in p is less or equal 1, i.e., (by definition) there exists a neighborhood U of p such that*

$$0 \rightarrow \mathcal{O}_U^m \xrightarrow{\alpha} \mathcal{O}_U^N \rightarrow \mathcal{S}_U \rightarrow 0$$

is exact for suitable m and N .

If (1) and (2) are fulfilled, then α in (1) is injective, i.e., m and N in (2) can be chosen to be $m = \text{cork}_p \mathcal{S}$ and $N = \text{rk} \mathcal{S} + m$.

Proof. — For the implication (1) \Rightarrow (2), we just need to show that α is injective: In points where \mathcal{S} is locally free, α is injective (due to the rank / dimension). Hence, $\mathcal{Ker} \alpha$ has support on a proper analytic set in U , i.e., is a torsion sheaf or the zero sheaf. Since \mathcal{O}_U does not contain any torsion sheaf, α is a monomorphism. (Alternatively, one can apply Lemma 3.19.)

(2) \Rightarrow (1): By the uniqueness of the minimal resolution (see e.g. [5, Thm. 20.2]), we can assume that N is equal to the minimal number of generators of \mathcal{S} in p , i.e.,

$$N = \text{rk}_p \mathcal{S} \stackrel{\text{def}}{=} \text{cork}_p \mathcal{S} + \text{rk} \mathcal{S}.$$

The injectivity of α implies $N - m = \text{rk}(\mathcal{O}^N / \alpha(\mathcal{O}^m)) = \text{rk} \mathcal{S}$, i.e.,

$$m = N - \text{rk} \mathcal{S} = \text{cork}_p \mathcal{S}. \quad \square$$

Remark 3.9. — We will work with Cohen–Macaulay spaces (for a definition and some crucial properties, see e.g. [16, §5]). Let us recall the following facts about Cohen–Macaulay spaces which will be used below:

- (i) A complex space X is Cohen–Macaulay in $p \in X$ if and only if for any (or at least one) non-zero-divisor f in the maximal ideal sheaf \mathfrak{m}_p , $\{f = 0\}$ is Cohen–Macaulay in p .
- (ii) If X is Cohen–Macaulay and A is an analytic subset of X with $\text{codim} A \geq 2$, then $\mathcal{O}(X) \rightarrow \mathcal{O}(X \setminus A)$ is bijective.
- (iii) A Cohen–Macaulay space is normal if and only if its singular set is at least 2-codimensional.

LEMMA 3.10. — *Let X be a normal or Cohen–Macaulay space and $S \subset X \times \mathbb{C}^N$ be a linear space over X with at least 2-codimensional singular*

locus in X and defined by one fiberwise linear function $h \in \mathcal{O}(X \times \mathbb{C}^N)$. Then S is locally irreducible. In particular, the coherent analytic sheaf associated to S is torsion-free.

Proof. — Let $A \subset X$ denote the singular locus of S (as linear space) and E denote the primary component of S . Lemma 3.6 implies that E is given by the ideal sheaf (h, g_1, \dots, g_m) with g_i holomorphic on $X \times \mathbb{C}^N$ and fiberwise homogeneous (shrink X if necessary). On the regular part $X' := X \setminus A$ of S , we get $S_{X'} = E_{X'}$, i.e., $g_{i,(p,z)} \in (h)_{(p,z)} \forall (p, z) \in X' \times \mathbb{C}^N$. Therefore, $f_i := g_i/h_i$ is a holomorphic function on $X' \times \mathbb{C}^N$. Since we assumed X to be normal or Cohen–Macaulay and A is of codimension 2 in X , f_i can be extended to a holomorphic function on $X \times \mathbb{C}^N$. We obtain $g_i \in (h)$ and $E = S$. Now, Lemma 3.2 implies the second statement. \square

Note that for the proof of Lemma 3.10, we hardly used the fact that S is given by a principal ideal sheaf. If S is defined by more than two functions while the corank of S is 1, it can happen that $\mathcal{S} = \mathcal{L}(S)$ has torsion elements with support on a 2-codimensional set. Since the singular locus of a torsion-free coherent analytic sheaf on a normal complex space is at least 2-codimensional (see Cor. of [14, Lem. 1.1.8]), we get the following corollary from Lemma 3.10.

COROLLARY 3.11. — *Let \mathcal{S} be a torsion-free coherent analytic sheaf on a normal complex space X such that $\mathcal{O}_X \rightarrow \mathcal{O}_X^N \rightarrow \mathcal{S} \rightarrow 0$ is exact (i.e., the homological dimension of \mathcal{S} is at most 1, see Lemma 3.8). Then the linear space associated to \mathcal{S} is locally irreducible.*

For a sheaf \mathcal{S} of homological dimension one, but with arbitrary corank, the associated linear space $L(\mathcal{S})$ is not necessarily locally irreducible (see Remark 3.3, where the ideal sheaf is of homological dimension one and of corank two). If $\text{codim}\{x : \text{cork}_x \mathcal{S} \geq k\} > k$ for all $k \geq 1$, then $L(\mathcal{S})$ is irreducible (see Prop. 3.11 in [2]). But, it is not clear whether $L(\mathcal{S})$ is reduced. At least, we can prove that \mathcal{S} is torsion-free:

LEMMA 3.12. — *Let X be a normal or Cohen–Macaulay space, and let \mathcal{S} be a coherent analytic sheaf on X such that the homological dimension of \mathcal{S} is at most one and $\text{Sing } \mathcal{S}$ has at least codimension 2 in X . Then \mathcal{S} is torsion-free.*

Proof. — Let A denote the singular locus of \mathcal{S} . By Lemma 3.8, every point $p \in X$ has a neighborhood U such that

$$(3.2) \quad 0 \rightarrow \mathcal{O}_U^m \xrightarrow{\alpha} \mathcal{O}_U^N \rightarrow \mathcal{S}_U \rightarrow 0,$$

with $m := \text{cork}_p \mathcal{S}$ and $N := \text{rk } \mathcal{S} + m$. Therefore, the associated linear space $S := L(\mathcal{S}_U)$ is defined by m holomorphic functions h_j on $U \times \mathbb{C}^N$ which are fiberwise linear. Let s be a torsion section of \mathcal{S} , i.e., there exists an $r \in \mathcal{O}_X^*(U)$ with $rs = 0$ for small enough U . We can understand $s(x, z) = \sum_{i=1}^N s_i(x)z_i$ as fiberwise linear holomorphic function on $U_x \times \mathbb{C}_z^N$, and will show that s is zero on S , i.e., s is in the ideal sheaf generated by h_j , $j = 1, \dots, m$:

$rs = 0$ on S implies that there exist m holomorphic functions a_j on U (shrinking U if necessary) such that

$$rs = \sum_{j=0}^m a_j h_j.$$

Since s is represented by the tuple (s_i) in \mathcal{O}_U^N , $(a_j) \in \mathcal{O}_U^m$ is the preimage $\alpha^{-1}(r \cdot (s_i)) = r\alpha^{-1}((s_i))$ under α in (3.2). (a_j) is uniquely determinate since α is injective. The support of $\mathcal{S}(\mathcal{S})$ is contained in A , i.e., $s = 0$ on S over $U' := U \setminus A$. In particular, for all $x \in U'$, there exists the decomposition

$$s_x = \sum_{j=0}^m b_{j,x} h_{j,x}.$$

A priori, b_j depends on x . Yet, since they are uniquely determined (α is injective), they are independent and exist on U' . In particular, $b_j = \frac{a_j}{r}$ on U' . Since A is at least 2 codimensional and $U \subset X$ is normal or Cohen-Macaulay, $\frac{a_j}{r}$ are holomorphic maps on U (see e.g. Remark 3.9(ii)). Hence,

$$s = \sum_{j=0}^m \frac{a_j}{r} h_j,$$

as desired. □

We call a normal complex space *factorial* if its structure sheaf is factorial (also called unique factorization domain). In this case, hypersurfaces are (locally) given as the zero set of one holomorphic function. The simplest examples for factorial spaces are manifolds.

THEOREM 3.13. — *Let S be a linear space over a factorial complex space X which is locally defined by one holomorphic fiberwise linear function in $X \times \mathbb{C}^{\text{rk } S+1}$ (i.e., the associated ideal sheaf is a principal ideal sheaf). Then the primary component of S is a linear space.*

Proof. — Let $S \subset X \times \mathbb{C}^N$ be given by the fiberwise linear $h \in \mathcal{O}(X \times \mathbb{C}^N)$. The primary component E of S is an irreducible hypersurface. Since X (and, hence, $X \times \mathbb{C}^N$) is factorial, the ideal sheaf \mathcal{I}_E is generated by one element g . By Lemma 3.6, we get g is fiberwise homogeneous. Moreover, g divides h . Hence, it has to be fiberwise linear. This implies that E is a linear space. □

LEMMA 3.14. — *Let X be a factorial complex space and $S \subset X \times \mathbb{C}^N$ a linear space associated to a torsion-free coherent analytic sheaf on X . Then S can be defined by locally irreducible holomorphic fiberwise linear functions.*

Proof. — Let the linear space S be defined by fiberwise linear $h_1, \dots, h_m \in \mathcal{O}(X \times \mathbb{C}^N)$. Let $S_i := \text{PC}(\{h_i = 0\})$ be defined by the fiberwise linear $g_i \in \mathcal{O}(X \times \mathbb{C}^N)$ (using Theorem 3.13). We will prove $S = \bigcap S_i$, i.e., $(h_i)_{i=1}^m = (g_i)_{i=1}^m$:

Since $g_j | h_j$, we get $(h_i)_{i=1}^m \subset (g_j)_{j=1}^m$. On the other hand, g_j vanishes on S_j . Hence, it vanishes on $\text{PC}(S)$, as well. Since the coherent analytic sheaf $\mathcal{L}(S)$ associated to S is torsion-free, we get $g_j = 0$ on S (in the non-reduced sense; see Remark 3.1), i.e., $g_j \in (h_i)_{i=1}^m$. \square

THEOREM 3.15. — *Let \mathcal{S} be a torsion-free coherent analytic sheaf of corank 1 on a factorial complex space X . Then the linear space associated to \mathcal{S} is locally irreducible and, for small enough open $U \subset X$, there exists an exact sequence*

$$0 \rightarrow \mathcal{O}_U \rightarrow \mathcal{O}_U^{\text{rk } \mathcal{S} + 1} \rightarrow \mathcal{S}_U \rightarrow 0,$$

i.e., the homological dimension of \mathcal{S} is at most 1 (see Lemma 3.8).

Proof. — Lemma 3.14 implies, for small enough open $U \subset X$, that the linear space S associated to \mathcal{S}_U can be defined by irreducible fiberwise linear $h_1, \dots, h_m \in \mathcal{O}(U \times \mathbb{C}^N)$ with $N = \text{rk } \mathcal{S} + 1$. Yet, the primary component E of S is already an irreducible hypersurface in $U \times \mathbb{C}^N$. Hence, E coincides with $S_i := \{h_i = 0\}$ and is a linear space. Lemma 3.5 implies $S = E = S_i$.

We obtain the exact sequence $\mathcal{O}_U \xrightarrow{h_i^*} \mathcal{O}_U^N \rightarrow \mathcal{S}_U \rightarrow 0$. Lemma 3.8, (1) \Rightarrow (2) or Lemma 3.19 give the injectivity of h_i^* . \square

Let us generalize this for sheaves with corank 2:

THEOREM 3.16. — *Let X be a factorial Cohen–Macaulay space and let S be a linear space of corank 2 on X such that $\text{Sing } S$ has at least codimension 3 in X and the coherent analytic sheaf $\mathcal{L}(S)$ associated to S is torsion-free. Then S is locally irreducible.*

Proof. — The proof is similar to the proof of Lemma 3.10. Let $S \subset U \times \mathbb{C}^N$ be defined by the fiberwise linear $h_1, \dots, h_m \in \mathcal{O}(U \times \mathbb{C}^N)$ for an open subset $U \subset X$ with $N = 2 + \text{rk } \mathcal{S}$. Because of Lemma 3.14, we can assume that $S_i := \{h_i = 0\}$ is locally irreducible. In particular, S_i is Cohen–Macaulay. Let us assume $h_1, h_2 \neq 0$ and $h_2 \notin (h_1)$. Since h_1 is irreducible,

$S_{12} := S_1 \cap S_2$ is a linear space with the same rank as S . We will prove that S_{12} coincides with $E := \text{PC}(S)$:

By Lemma 3.6, E is defined by g_1, \dots, g_k (in particular, $h_i \in (g_1, \dots, g_k)$). Let A denote the singular locus of S and $U' = U \setminus A$. Since $E \subset S_{12}$ and $\dim E_{U'} = \dim S_{12,U'}$, we get $E = S_{12}$ over U' . Hence, we obtain $g_{j,(p,z)} \in (h_1, h_2)_{(p,z)}$ for all $(p, z) \in S_{1,U'}$. This means $\frac{g_j}{h_2}$ is a holomorphic function on $S_{1,U'}$. By assumption, $\text{codim}_{S_1}((A \times \mathbb{C}^N) \cap S_1) \geq \text{codim}_{U \times \mathbb{C}^N}(A \times \mathbb{C}^N) - 1 \geq 2$. Taking into account that S_1 is Cohen–Macaulay, we conclude that $\frac{g_j}{h_2}$ is holomorphic on S_1 and, hence, $g_j \in (h_1, h_2)$, i.e., $E = S_{12}$. Lemma 3.5 implies the claim. \square

COROLLARY 3.17. — *Let X be a factorial Cohen–Macaulay space and \mathcal{S} be a torsion-free coherent analytic sheaf \mathcal{S} of corank 2 on X with at least 3-codimensional singular locus. Then the linear space associated to \mathcal{S} is locally irreducible and, for small enough open $U \subset X$, there exists an exact sequence*

$$0 \rightarrow \mathcal{O}_U^2 \rightarrow \mathcal{O}_U^{\text{rk } \mathcal{S} + 2} \rightarrow \mathcal{S}_U \rightarrow 0,$$

i.e., the homological dimension of \mathcal{S} is at most 1.

Keeping the counter-example Remark 3.3 in mind, one sees the assumption on the codimension is necessary.

Proof of Theorem 1.3. — Lemma 3.2 gives the implication (2) \Rightarrow (1), and Theorem 3.15 and Corollary 3.17 yield the implication (1) \Rightarrow (3, 4). (4) \Rightarrow (1) is obtained by Lemma 3.12. There is one implications left.

(3) \Rightarrow (2): Assume that (3) is satisfied, i.e., that $S := L(\mathcal{S})$ is locally irreducible. By definition, this implies that S is reduced. But S is connected. So, there can be just one irreducible component, i.e., (2) holds also. \square

If the codimension of the singular set of \mathcal{S} is big enough, we can prove normality of the linear space associated to \mathcal{S} .

THEOREM 3.18. — *Let X be a factorial Cohen–Macaulay space and \mathcal{S} be a torsion-free coherent analytic sheaf on X of corank at most 2 with $\text{codim } \text{Sing } \mathcal{S} \geq 2 + \text{cork } \mathcal{S}$. Then the linear space associated to \mathcal{S} is normal.*

Proof. — Let $S \subset U \times \mathbb{C}^N$ be the linear space associated to \mathcal{S} on an open subset U of X with $N = \text{rk } \mathcal{S} + \text{cork } \mathcal{S}$. Theorem 3.15 or Corollary 3.17 implies that S is Cohen–Macaulay. Let $A := \text{Sing } S = \text{Sing } \mathcal{S}_U \subset U$ denote the singular locus of S as linear space. The singular set S_{sing} of S as analytic subset of $U \times \mathbb{C}^N$ is contained in $((A \times \mathbb{C}^N) \cap S) \cup S_{U_{\text{sing}} \setminus A}$. We get

$$\text{codim}_S((A \times \mathbb{C}^N) \cap S) \geq \text{codim}_U A - \text{cork } \mathcal{S} \geq 2$$

and $\text{codim}_S S_{U_{\text{sing}} \setminus A} \geq \text{codim } U_{\text{sing}} \geq 2$ since $U \subset X$ is normal and $S_{U \setminus A}$ is a vector bundle. Hence, $\text{codim}_S S_{\text{sing}} \geq 2$. Remark 3.9(iii) (see e.g. [16, Cor. 5.2]) implies that S is actually normal. \square

3.3. More preliminaries on torsion

Throughout the paper, we will use the following observation without mentioning explicitly. Let $\psi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of analytic sheaves on a (locally irreducible) complex space (X, \mathcal{O}_X) . Then ψ induces a canonical map

$$\widehat{\psi}: \mathcal{F} / \mathcal{T}(\mathcal{F}) \rightarrow \mathcal{G} / \mathcal{T}(\mathcal{G})$$

because the torsion sheaf $\mathcal{T}(\mathcal{F})$ of \mathcal{F} is mapped by ψ into the torsion sheaf $\mathcal{T}(\mathcal{G})$ of \mathcal{G} : $r_x \psi(s_x) = \psi(r_x s_x) = 0$ for germs $r_x \in \mathcal{O}_{X,x}$, $s_x \in \mathcal{T}(\mathcal{F})_x$ with $r_x s_x = 0$. Note that particularly $\mathcal{T}(\mathcal{F}) \subset \ker \psi$ if \mathcal{G} is torsion-free.

LEMMA 3.19. — *Let X be a locally irreducible complex space, \mathcal{F} and \mathcal{G} coherent analytic sheaves on X such that there exists a morphism $\psi: \mathcal{F} \rightarrow \mathcal{G}$ which is a monomorphism on an open dense subset of X . If \mathcal{F} is torsion-free, then ψ is a monomorphism. If not, ψ induces a monomorphism $\widehat{\psi}: \mathcal{F} / \mathcal{T}(\mathcal{F}) \hookrightarrow \mathcal{G} / \mathcal{T}(\mathcal{G})$.*

Proof. — The second statement follows from the considerations above and the torsion-free case. Hence, we can assume that \mathcal{F} is torsion-free. Let F and G denote the linear spaces associated to \mathcal{F} and \mathcal{G} , respectively. Theorem 3.4 implies that $\text{PC}(F)$ and $\text{PC}(G)$ are locally irreducible. Let ψ be a monomorphism on the open dense subset W of X with $W \subset X \setminus (\text{Sing } \mathcal{F} \cap \text{Sing } \mathcal{G})$. Thus, ψ induces a holomorphic fiberwise linear map $\psi^*: G \rightarrow F$ such that $\psi^*_W: G_W \rightarrow F_W$ is a surjective map of vector bundles. Let s be a section in $\mathcal{Ker } \psi$, i.e., $\psi^* \circ s$ vanishes on G . We get that s vanishes on F_W and, hence, on $\text{PC}(F)$. Since \mathcal{F} is torsion free, we obtain $s = 0$ (cf. Remark 3.1). \square

Alternatively, one can prove this lemma by using only sheaf-theoretical terminology and arguments (cf. the proof of Lemma 5.1).

Note the following trivial consequence of the definition of the pullback:

LEMMA 3.20. — *Let $S \subset U \times \mathbb{C}^N$ be a linear space over a complex space U , and let $\pi: V \rightarrow U$ be a holomorphic map. Then the pullback $\pi^*(S) = V \times_U S$ can be embedded in $V \times \mathbb{C}^N$.*

Remark 3.21. — Let \mathcal{S} be a coherent analytic sheaf over a locally irreducible complex space X . Let $\varphi = \varphi_{\mathcal{S}}: X_{\mathcal{S}} \rightarrow X$ be the monoidal transformation of X with respect to \mathcal{S} , i.e., $\mathcal{E} := \varphi^T \mathcal{S}$ is locally free. Note that $X_{\mathcal{S}}$ is again locally irreducible. Then $\varphi^* \mathcal{S}$ has torsion in a point q if and only if \mathcal{S} is not locally free in $\varphi(q)$ (see [18]). We will give a short, alternative proof with the statements from above. Let S, S^* and E denote the linear complex spaces associated to $\mathcal{S}, \varphi^* \mathcal{S}$ and \mathcal{E} , respectively. If \mathcal{S} is not locally free in $\varphi(q)$, then $\dim E_q < \dim S_{\varphi(q)}$ (as $\dim E_q = \text{rk } E = \dim S_{\tilde{q}}$ in all points \tilde{q} where \mathcal{S} is locally free, see [11, §1.1]). Lemma 3.20 implies $\dim S_q^* = \dim S_{\varphi(q)} > \dim E_q$. Since $\text{PC}(S^*) = E$, we obtain that S^* is reducible in $(q, 0)$, i.e., $\varphi^* \mathcal{S}$ has torsion in q by Lemma 3.5. The other implication of the claim is trivial.

4. Direct images of torsion-free preimage sheaves

In this section, we will prove Theorem 1.4 and more:

THEOREM 4.1. — *Let X be a locally irreducible complex space, \mathcal{S} a torsion-free coherent analytic sheaf on X and $\pi: Y \rightarrow X$ a proper modification of X .*

- (i) *Then the canonical homomorphisms $\mathcal{S} \rightarrow \pi_*(\pi^* \mathcal{S})$ and $\mathcal{S} \rightarrow \pi_*(\pi^T \mathcal{S})$ both are injective, where $\pi^T \mathcal{S}$ is the torsion-free preimage sheaf of \mathcal{S} under π .*
- (ii) *If the linear space $L(\mathcal{S})$ associated to \mathcal{S} is locally irreducible and $L(\pi^* \mathcal{S})$ is reduced, then $\pi_*(\pi^* \mathcal{S}) \rightarrow \pi_*(\pi^T \mathcal{S})$ is injective.*
- (iii) *If the linear space $L(\mathcal{S})$ is normal, then*

$$\mathcal{S} \cong \pi_*(\pi^T \mathcal{S}).$$

Proof. — We can assume that X is connected. Let S denote the linear space associated to \mathcal{S} , $A \subset Y$ the set where π is not biholomorphic and A^c the complement. $S^* = Y \times_X S$ is the linear space associated to $\mathcal{S}^* := \pi^* \mathcal{S}$. Let $\text{pr}: S^* \rightarrow S$ denote the projection, let E be the linear space associated to $\mathcal{E} := \pi^T \mathcal{S}$, let U be an open Stein set in X , and let $V := \pi^{-1}(U)$. The construction of the linear spaces implies

$$\begin{aligned} \text{Hom}(S_U, U \times \mathbb{C}) &\cong \mathcal{S}(U), \\ \text{Hom}(S_V^*, V \times \mathbb{C}) &\cong \pi^* \mathcal{S}(V) = (\pi_*(\pi^* \mathcal{S}))(U) \\ \text{and } \text{Hom}(E_V, V \times \mathbb{C}) &\cong \mathcal{E}(V) = (\pi_* \mathcal{E})(U). \end{aligned}$$

Let N be an integer big enough, so that S_U can be realized as a subset of $U \times \mathbb{C}^N$. We obtain closed embeddings $E_V \subset S_V^* \subset V \times \mathbb{C}^N$, and (q, z) is

in S_V^* if and only if $(\pi(q), z) \in S$. Since obviously $\text{pr}(E_{A^c}) = S_{\pi(A^c)}$ and proper holomorphic images of irreducible sets are irreducible, we obtain (cf. (3.1))

$$(4.1) \quad \text{pr}(\text{PC}(E_V)) = \text{PC}(S_U).$$

(i) $\mathcal{S} \hookrightarrow \pi_*(\pi^*\mathcal{S})$ and $\mathcal{S} \hookrightarrow \pi_*(\pi^T\mathcal{S})$: This follows from Lemma 3.19.

(ii) Assume S is locally irreducible and S^* is reduced. To prove that the natural map given by the restriction $\text{Hom}(S_V^*, V \times \mathbb{C}) \rightarrow \text{Hom}(E_V, V \times \mathbb{C})$ is injective, we apply $\text{pr}(\text{PC}(E_V)) = S_U$ (use (4.1) and that S is irreducible):

Let $s \in \text{Hom}(S_V^*, V \times \mathbb{C})$ with $s|_E = 0$, i.e., $s|_{\text{PC}(E)} = 0$. Let $s(q, z) = (q, f(q, z)) \in \text{Hom}(S_V^*, V \times \mathbb{C})$ be not the zero section, i.e., there is a point $(q', z_0) \in S^*$ with $f(q', z_0) \neq 0$. There is a $q'' \in \pi^{-1}(\pi(q'))$ such that $(q'', z_0) \in E$. Since $\pi^{-1}(\pi(q')) \times \{z_0\}$ is a compact analytic set in S_V^* , we get $f(q', z_0) = f(q'', z_0)$, i.e., $f|_E \neq 0$ and $s|_E \neq 0$.

(iii) Assume that S is normal. Fix a section $s(q, z) = (q, f(q, z)) \in \text{Hom}(E_V, V \times \mathbb{C})$. Since $\text{pr}: \text{PC}(E_V) \rightarrow \text{PC}(S_U) = S_U$ is a proper modification (surjectivity is (4.1)), the map $\tilde{f} := f \circ \text{pr}^{-1}: S_U \rightarrow \mathbb{C}$ is a bounded meromorphic function, i.e., it is weakly holomorphic. Since S_U is normal, \tilde{f} is holomorphic. Obviously, it is linear in the second argument. Hence, pr^{-1} gives a map $(\text{pr}^{-1})^*: \text{Hom}(E_V, V \times \mathbb{C}) \rightarrow \text{Hom}(S_U, U \times \mathbb{C})$, $s \mapsto \tilde{s}$ with $\tilde{s}(p, z) = (p, \tilde{f}(p, z))$. Since $f \circ \text{pr}^{-1} = \tilde{f} = 0$ implies $f = 0$, this map is injective. It is now easy to see that $(\text{pr}^{-1})^*: \pi_*\pi^T\mathcal{S} \hookrightarrow \mathcal{S}$ is the inverse to the natural mapping $\mathcal{S} \hookrightarrow \pi_*\pi^T\mathcal{S}$. \square

Remark 4.2. — Without the additional assumption about normality, the natural map $\mathcal{S} \hookrightarrow \pi_*\pi^T\mathcal{S}$ is not necessarily bijective. The following counter-example is derived from one due to Mircea Mustațǎ.

Let $\mathcal{S} = (x^3, y^3)$ be the ideal sheaf on $\mathbb{C}_{x,y}^2$ generated by the functions x^3 and y^3 , and let $\pi: M \rightarrow \mathbb{C}^2$ be the blow up of the origin, i.e.,

$$M = \{(x, y; [t_1 : t_2]) \in \mathbb{C}^2 \times \mathbb{C}\mathbb{P}^1 : xt_2 = yt_1\}.$$

Then

$$S = L(\mathcal{S}) = \{(x, y; z_1, z_2) \in \mathbb{C}^2 \times \mathbb{C}^2 : z_2x^3 = z_1y^3\},$$

and sections in \mathcal{S} correspond to sections in $\text{Hom}(S, \mathbb{C}^2 \times \mathbb{C})$ via the assignment $x^3 \mapsto [(x, y; z_1, z_2) \mapsto z_1]$, $y^3 \mapsto [(x, y; z_1, z_2) \mapsto z_2]$. Now,

$$S^* = L(\pi^*\mathcal{S}) = \{(x, y; [t_1:t_2]; z_1, z_2) \in \mathbb{C}^2 \times \mathbb{C}\mathbb{P}^1 \times \mathbb{C}^2 : xt_2=yt_1, z_2x^3=z_1y^3\},$$

$$E = L(\pi^T\mathcal{S}) = \{(x, y; [t_1:t_2]; z_1, z_2) \in \mathbb{C}^2 \times \mathbb{C}\mathbb{P}^1 \times \mathbb{C}^2 : xt_2=yt_1, z_2t_1^3=z_1t_2^3\}.$$

Thus, $S^* = E \cup T$ with

$$T = \{(x, y; [t_1:t_2]; z_1, z_2) \in \mathbb{C}^2 \times \mathbb{C}\mathbb{P}^1 \times \mathbb{C}^2 : x = y = 0\}.$$

In $\text{Hom}(E, M \times \mathbb{C})$, we have now also the section

$$\left\{ \left(\frac{t_2}{t_1} z_1 : t_1 \neq 0 \right); \left(\frac{t_1^2}{t_2} z_2 : t_2 \neq 0 \right) \right\},$$

corresponding to x^2y in $\pi^T \mathcal{S}$, and the section

$$\left\{ \left(\frac{t_2^2}{t_1} z_1 : t_1 \neq 0 \right); \left(\frac{t_1}{t_2} z_2 : t_2 \neq 0 \right) \right\},$$

corresponding to xy^2 , but these two do not extend to $S^* = E \cup T$ because there is no relation between z_1 and z_2 on T . In $\text{Hom}(S^*, M \times \mathbb{C})$, however, we have the section

$$\left\{ \left(y \frac{t_2}{t_1} z_1 : t_1 \neq 0 \right); \left(y \frac{t_1^2}{t_2} z_2 : t_2 \neq 0 \right) \right\},$$

corresponding to x^2y^2 in $\pi^* \mathcal{S}$. Moreover, it is easy to check that x, x^2, y, y^2, xy are neither contained in $\pi_* \pi^* \mathcal{S}$ nor in $\pi_* \pi^T \mathcal{S}$. Hence, we have:

$$\mathcal{S} \subsetneq (x^3, x^2y^2, y^3) = \pi_* \pi^* \mathcal{S} \subsetneq (x^3, x^2y, xy^2, y^3) = \pi_* \pi^T \mathcal{S}.$$

Let us present a counter-example for (ii) in Theorem 4.1 if S is not irreducible:

Let $X = \{x^3 = y^2\} \subset \mathbb{C}^2$ be the cusp, $\pi: \mathbb{C} \rightarrow X$, $\pi(t) := (t^2, t^3)$ the normalization and let $\widehat{\mathcal{O}}_X$ denote the sheaf of weakly holomorphic functions on X . Then one can compute that $\pi_* \pi^* \widehat{\mathcal{O}}_X$ has torsion elements with support in 0. Yet, $\pi_* \pi^T \widehat{\mathcal{O}}_X = \pi_* \mathcal{O}_{\mathbb{C}} = \widehat{\mathcal{O}}_X$ is torsion-free. Hence, there can not exist an injective morphism $\pi_* \pi^* \widehat{\mathcal{O}}_X \rightarrow \pi_* \pi^T \widehat{\mathcal{O}}_X$.

Remark 4.3. — In order to get the isomorphism $\mathcal{S} = \pi_* \pi^T \mathcal{S}$, normality is a natural assumption. For example, if \mathcal{E} is locally free, we obtain $\pi_* \pi^* \mathcal{E} \cong \mathcal{E}$ if and only if $\pi_* \mathcal{O}_Y \cong \mathcal{O}_X$.

5. Torsion-free preimages of direct image sheaves

LEMMA 5.1. — *Let $\pi: Y \rightarrow X$ be a proper modification between complex spaces Y, X , and \mathcal{E} a torsion-free coherent analytic sheaf on Y . Then the canonical homomorphism $\pi^* \pi_* \mathcal{E} \rightarrow \mathcal{E}$ induces a canonical injection*

$$(5.1) \quad \pi^T \pi_* \mathcal{E} \hookrightarrow \mathcal{E},$$

where $\pi^T \pi_* \mathcal{E} = \pi^* \pi_* \mathcal{E} / \mathcal{F}(\pi^* \pi_* \mathcal{E})$ is the torsion-free preimage of $\pi_* \mathcal{E}$.

The following proof was communicated to us by Matei Toma. Alternatively, Lemma 5.1 follows also from Lemma 3.19.

Proof. — Let \mathcal{T} denote the torsion sheaf of $\pi^*\pi_*\mathcal{E}$, and let $\psi: \pi^*\pi_*\mathcal{E} \rightarrow \mathcal{E}$ denote the natural map. Since \mathcal{E} is torsion-free, $\psi(\mathcal{T}) = 0$ and, hence, ψ factors through $\widehat{\psi}: \pi^T\pi_*\mathcal{E} \rightarrow \mathcal{E}$. Since ψ is an isomorphism outside of a thin analytic set $A \subset X$, an element in the kernel of ψ has support in A . Therefore, the kernel is a subset of \mathcal{T} , i.e., $\widehat{\psi}$ is injective. \square

Remark 5.2. — Let us give a counter-example showing that (5.1) is in general not an isomorphism. Consider a modification $\pi: M \rightarrow \mathbb{C}^n$ where M is a complex manifold with canonical sheaves Ω_M^n and $\Omega_{\mathbb{C}^n}^n \cong \mathcal{O}_{\mathbb{C}^n}$. Then $\pi_*\Omega_M^n = \Omega_{\mathbb{C}^n}^n \cong \mathcal{O}_{\mathbb{C}^n}$ so that $\pi^T\pi_*\Omega_M^n \cong \pi^T\mathcal{O}_{\mathbb{C}^n} \cong \mathcal{O}_M$. But $\mathcal{O}_M \neq \Omega_M^n$ in general.

However, we can be a bit more precise in Lemma 5.1 by use of the following observation if \mathcal{E} is locally free of rank 1:

LEMMA 5.3. — *Let X be a complex space and $i: \mathcal{F} \hookrightarrow \mathcal{G}$ an injective morphism between two coherent locally free sheaves of rank 1 over X . Then there exists an effective Cartier divisor, $D \geq 0$, such that*

$$i(\mathcal{F}) = \mathcal{G} \otimes \mathcal{O}_X(-D).$$

In particular, i is an isomorphism precisely on $X - |D|$.

Proof. — Let $\{X_\alpha\}_\alpha$ be a locally finite open cover of X such that both, \mathcal{F} and \mathcal{G} , are free over each X_α . So, there are trivializations

$$\begin{aligned} \phi_\alpha: \mathcal{F}|_{X_\alpha} &\xrightarrow{\sim} \mathcal{O}_{X_\alpha}, \\ \psi_\alpha: \mathcal{G}|_{X_\alpha} &\xrightarrow{\sim} \mathcal{O}_{X_\alpha}, \end{aligned}$$

and for $X_{\alpha\beta} := X_\alpha \cap X_\beta \neq \emptyset$, we have transition functions $F_{\beta\alpha} := \phi_\beta \circ \phi_\alpha^{-1} \in \mathcal{O}^*(X_{\alpha\beta})$ and $G_{\beta\alpha} := \psi_\beta \circ \psi_\alpha^{-1} \in \mathcal{O}^*(X_{\alpha\beta})$ satisfying the cocycle conditions. In trivializations

$$\psi_\alpha \circ i|_{X_\alpha} \circ \phi_\alpha^{-1}: \mathcal{O}_{X_\alpha} \rightarrow \mathcal{O}_{X_\alpha}$$

is given by a holomorphic function $i_\alpha \in \mathcal{O}(X_\alpha)$, vanishing nowhere identically, with (unreduced) divisor (i_α) . It is easy to see that $G_{\beta\alpha} \cdot i_\alpha = i_\beta \cdot F_{\beta\alpha}$ on $X_{\alpha\beta}$, so that $i_\alpha/i_\beta = F_{\beta\alpha}/G_{\beta\alpha} \in \mathcal{O}^*(X_{\alpha\beta})$. Thus $D := \{(X_\alpha, i_\alpha)\}_\alpha$ defines in fact an effective Cartier divisor with support $|D|$.

To see that $i(\mathcal{F}) = \mathcal{G} \otimes \mathcal{O}_X(-D)$, note that $\mathcal{G} \otimes \mathcal{O}_X(-D)$ is a coherent subsheaf of \mathcal{G} because $\mathcal{O}_X(-D)$ is a sheaf of ideals in \mathcal{O}_X , and that

$$\psi_\alpha \otimes 1: \mathcal{G} \otimes \mathcal{O}_X(-D)|_{X_\alpha} \xrightarrow{\sim} \mathcal{O}_{X_\alpha} \otimes \mathcal{O}_{X_\alpha}(-(i_\alpha)). \quad \square$$

So, we can deduce the following direct consequence of Lemma 5.1:

THEOREM 5.4. — *Let $\pi: Y \rightarrow X$ be a proper modification of X , \mathcal{E} a locally free analytic sheaf of rank 1 on Y and assume that $\pi^T \pi_* \mathcal{E}$ is also locally free. Then there exists an effective Cartier divisor D on Y such the following holds: The canonical homomorphism $\pi^* \pi_* \mathcal{E} \rightarrow \mathcal{E}$ induces a canonical injection*

$$i: \pi^T \pi_* \mathcal{E} \hookrightarrow \mathcal{E}$$

and

$$i(\pi^T \pi_* \mathcal{E}) = \mathcal{E} \otimes \mathcal{O}_Y(-D).$$

In particular, i is an isomorphism precisely on $Y - |D|$, and $|D|$ is contained in the exceptional set of π .

6. π^T preserves injectivity and surjectivity

In this section, we consider properties of π^T as functor and use them to prove Theorem 1.2.

LEMMA 6.1. — *Let $\pi: Y \rightarrow X$ be a proper modification of a complex space X such that Y is locally irreducible. Let*

$$\psi: \mathcal{F} \rightarrow \mathcal{G}$$

be a morphism of coherent analytic sheaves. If ψ is an epimorphism, then the induced mapping

$$\pi^T \psi: \pi^T \mathcal{F} \rightarrow \pi^T \mathcal{G}$$

is also an epimorphism. If ψ is a monomorphism, then $\pi^T \psi$ is also a monomorphism.

Proof. — Let ψ be an epimorphism, i.e., surjective. Recall that π^* is right-exact. So, $\pi^* \psi: \pi^* \mathcal{F} \rightarrow \pi^* \mathcal{G}$ is still surjective. But then it is easy to see that the induced mapping $\pi^T \psi: \pi^T \mathcal{F} \rightarrow \pi^T \mathcal{G}$ is also surjective.

For the second statement, let ψ be injective. Let $f_x \in (\pi^T \mathcal{F})_x$ such that $\pi^T \psi(f_x) = 0$. This means that there is an open set $U \subset Y$ and a representative $f \in \pi^T \mathcal{F}(U)$ such that $\pi^T \psi(f) = 0$. But $\pi^T \psi$ is injective on a dense open subset $W \subset X$. Thus, $f = 0$ on $U \cap W$, i.e., f has support on a thin set. But $\pi^T \mathcal{F}$ is torsion-free. So, $f_x = 0$ and $f = 0$. \square

Note that π^T is not exact. A simple counter-example is as follows. Let \mathfrak{m} be the maximal ideal sheaf of the origin in \mathbb{C}^2 . Then $0 \rightarrow \mathfrak{m} \hookrightarrow \mathcal{O}_{\mathbb{C}^2} \rightarrow \mathcal{O}_{\mathbb{C}^2}/\mathfrak{m} \rightarrow 0$ is exact. Let π be just the identity on \mathbb{C}^2 . So, we have $\pi^T \mathfrak{m} = \mathfrak{m}$, $\pi^T \mathcal{O}_{\mathbb{C}^2} = \mathcal{O}_{\mathbb{C}^2}$ and $\pi^T(\mathcal{O}_{\mathbb{C}^2}/\mathfrak{m}) = 0$. The resulting sequence $0 \rightarrow \mathfrak{m} \hookrightarrow \mathcal{O}_{\mathbb{C}^2} \rightarrow 0$ is clearly not exact.

Proof of Theorem 1.2. — Let $\pi: Y \rightarrow X$ be a proper modification between locally irreducible complex spaces. Let \mathcal{F} and \mathcal{G} be coherent analytic sheaves on X and Y , respectively.

(i) The case $\mathcal{F} = \pi_*\mathcal{G}$: By Theorem 4.1(i), the natural map $\mathcal{F} \rightarrow \pi_*\pi^T\mathcal{F}$ is injective. Moreover, Lemma 5.1 yields injectivity of the natural map

$$\pi^T\mathcal{F} = \pi^T\pi_*\mathcal{G} \rightarrow \mathcal{G}.$$

Since π_* is left-exact, we obtain the second natural injection

$$\pi_*\pi^T\mathcal{F} \hookrightarrow \pi_*\mathcal{G} = \mathcal{F}.$$

(ii) The case $\mathcal{G} = \pi^T\mathcal{F}$: As above, Lemma 5.1 gives $\pi^T\pi_*\mathcal{G} \hookrightarrow \mathcal{G}$. By Theorem 4.1(i) we have also the natural injection

$$\mathcal{F} \hookrightarrow \pi_*\pi^T\mathcal{F} = \pi_*\mathcal{G}.$$

But π^T preserves injectivity (Lemma 6.1) so that we obtain the injection

$$\mathcal{G} = \pi^T\mathcal{F} \hookrightarrow \pi^T\pi_*\mathcal{G}. \quad \square$$

7. Application to ideal sheaves

In this section, we discuss the application of Theorem 1.2 to reduced ideal sheaves. As a preparation, note the following:

LEMMA 7.1. — *Let $\pi: Y \rightarrow X$ be a holomorphic mapping between complex spaces Y, X . Then $\pi^*\mathcal{O}_X = \mathcal{O}_Y$.*

Proof. — As $\pi^{-1}\mathcal{O}_X \subset \mathcal{O}_Y$, we have that $\pi^*\mathcal{O}_X = \pi^{-1}\mathcal{O}_X \otimes_{\pi^{-1}\mathcal{O}_X} \mathcal{O}_Y = \mathcal{O}_Y$, because $\pi^{-1}\mathcal{O}_X$ contains (the germ of) the function 1 at any point of Y . \square

Coming to ideal sheaves, let us start with the following observation:

LEMMA 7.2. — *Let X be a locally irreducible complex space and $A \subset X$ an (unreduced) analytic subspace with ideal sheaf \mathcal{I}_A . Let $\pi: Y \rightarrow X$ be a proper modification and $B := \pi^{-1}(A)$ the unreduced analytic preimage with ideal sheaf \mathcal{I}_B . Then:*

$$\pi^T\mathcal{I}_A = \mathcal{I}_B.$$

Proof. — Consider the short exact sequence of sheaves over X :

$$0 \rightarrow \mathcal{I}_A \xrightarrow{\alpha} \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{I}_A \rightarrow 0.$$

By right-exactness of π^* , we deduce the exact sequence

$$\pi^* \mathcal{I}_A \xrightarrow{\pi^* \alpha} \pi^* \mathcal{O}_X \rightarrow \pi^*(\mathcal{O}_X/\mathcal{I}_A) \rightarrow 0.$$

Now, we use Lemma 7.1 twice: $\pi^* \mathcal{O}_X = \mathcal{O}_Y$ and $\pi|_B^* \mathcal{O}_A = \mathcal{O}_B$ (which implies that $\pi^*(\mathcal{O}_X/\mathcal{I}_A) = \mathcal{O}_Y/\mathcal{I}_B$ using the definition of the analytic preimage, see e.g. Prop. 0.27 in [8]). As \mathcal{O}_Y is torsion-free, it is clear that

$$(7.1) \quad \mathcal{F}(\pi^* \mathcal{I}_A) \subset \text{Ker } \pi^* \alpha.$$

Consider $\pi^T \alpha: \pi^T \mathcal{I}_A \rightarrow \pi^T \mathcal{O}_X = \pi^* \mathcal{O}_X = \mathcal{O}_Y$. By (7.1), it follows that $\mathcal{I}m(\pi^* \alpha) = \mathcal{I}m(\pi^T \alpha)$, and Lemma 6.1 tells us that $\pi^T \alpha$ is injective. So, we obtain the short exact sequence

$$0 \rightarrow \pi^T \mathcal{I}_A \xrightarrow{\pi^T \alpha} \mathcal{O}_Y \rightarrow \mathcal{O}_Y/\mathcal{I}_B \rightarrow 0,$$

telling us that in fact $\pi^T \mathcal{I}_A = \mathcal{I}_B$. □

It is clear that \mathcal{I}_A (and $\pi^T \mathcal{I}_A = \mathcal{I}_B$) are torsion-free, and so we obtain from Theorem 1.2(ii) that:

$$(7.2) \quad \mathcal{I}_B \cong \pi^T \pi_* \mathcal{I}_B.$$

Under some additional assumptions, we have also:

LEMMA 7.3. — *Let X be a normal complex space, and let A be a locally complete intersection or a normal analytic set in X with (reduced) ideal sheaf \mathcal{I}_A . Let $\sigma: \tilde{X} \rightarrow X$ denote the blow up of X with center A , i.e. the monoidal transformation with respect to \mathcal{I}_A , and let \mathcal{I}_B be the (reduced) ideal sheaf associated to the reduced exceptional set $B := \sigma^{-1}(A)$. Then:*

$$(7.3) \quad \mathcal{I}_A \cong \sigma_* \mathcal{I}_B.$$

Proof. — The statement is local with respect to X , so we can assume that A is the zero-set of reduced holomorphic functions f_0, \dots, f_m . $(\mathcal{I}_A)_p$ is generated by the germs $f_{0,p}, \dots, f_{m,p}$, and $\tilde{X} \subset X \times \mathbb{C}\mathbb{P}^m$ (see e.g. [19, §2.5] for the monoidal transformation of ideal sheaves). We show that

$$\mathcal{O}_A \cong \sigma_* \mathcal{O}_B.$$

(I) A is a complete intersection, i.e., $m + 1 = \text{codim } A$: This implies $B = A \times \mathbb{C}\mathbb{P}^m$. For all open subsets $U \subset A$, we obtain

$$\mathcal{O}_A(U) \cong \mathcal{O}_{A \times \mathbb{C}\mathbb{P}^m}(U \times \mathbb{C}\mathbb{P}^m) = \mathcal{O}_B(\sigma^{-1}(U)).$$

(II) A is normal: In this case, B is an analytic subset of $A \times \mathbb{C}\mathbb{P}^m$, and by the surjectivity $\sigma(B) = A$, we get the injection $\mathcal{O}_A \cong \sigma_* \mathcal{O}_{A \times \mathbb{C}\mathbb{P}^m} \hookrightarrow \sigma_* \mathcal{O}_B$.

On the other hand, a section in $\mathcal{O}_B(\sigma^{-1}(U))$ gives a weakly holomorphic function on A : With part (I) applied on the regular part A_{reg} of A , we get a holomorphic function on A_{reg} which is bounded in points of A_{sing} . Since we assumed A to be normal, we get $\sigma_*\mathcal{O}_B \cong \widehat{\mathcal{O}}_A \cong \mathcal{O}_A$.

Thus, $\mathcal{O}_A \cong \sigma_*\mathcal{O}_B$ as desired. In other words: $\mathcal{O}_X/\mathcal{I}_A \cong \sigma_*(\mathcal{O}_{\widetilde{X}}/\mathcal{I}_B)$. We obtain the exact commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \sigma_*\mathcal{I}_B & \longrightarrow & \sigma_*\mathcal{O}_{\widetilde{X}} & \longrightarrow & \sigma_*(\mathcal{O}_{\widetilde{X}}/\mathcal{I}_B) \\
 & & & & \parallel \wr & & \parallel \wr \\
 0 & \longrightarrow & \mathcal{I}_A & \longrightarrow & \mathcal{O}_X & \longrightarrow & \mathcal{O}_X/\mathcal{I}_A \longrightarrow 0.
 \end{array}$$

It follows that in fact $\sigma_*\mathcal{I}_B \cong \mathcal{I}_A$. □

In the situation of Lemma 7.3, we can now apply Theorem 1.2(i) to $\mathcal{I}_A \cong \sigma_*\mathcal{I}_B$ and obtain:

$$(7.4) \quad \mathcal{I}_A \cong \sigma_*\sigma^T \mathcal{I}_A.$$

8. Holomorphic n -forms on singular spaces

As a consequence of Theorem 1.2 and Theorem 5.4, we get also the following application to holomorphic n -forms:

THEOREM 8.1. — *Let X be a complex space of pure dimension n and \mathcal{K}_X the Grauert–Riemenschneider canonical sheaf on X . Then there exist a resolution of singularities $\pi: M \rightarrow X$ and an effective divisor, $D \geq 0$, with support on the exceptional set of the resolution such that*

$$(8.1) \quad \pi^T \mathcal{K}_X \cong \Omega_M^n(-D) = \Omega_M^n \otimes \mathcal{O}_M(-D),$$

where Ω_M^n is the canonical sheaf of holomorphic n -forms on M , and (8.1) is induced by the natural mapping $\pi^* \mathcal{K}_X = \pi^* \pi_* \Omega_M^n \rightarrow \Omega_M^n$. Moreover, we get

$$\pi_* \Omega_M^n = \mathcal{K}_X \cong \pi_* \Omega_M^n(-D).$$

Proof. — Let $\pi: M \rightarrow X$ be a resolution of singularities such that $\pi^T \mathcal{K}_X$ is locally free. Such a resolution exists due to Rossi and Hironaka (apply first Rossi’s Theorem 2.3 and then Hironaka’s resolution of singularities). Recall that $\mathcal{K}_X = \pi_* \Omega_M^n$ by definition of the Grauert–Riemenschneider canonical sheaf. So, the assertion follows directly from Theorem 5.4 and Theorem 1.2. □

The following observation is also useful:

LEMMA 8.2. — *Let \mathcal{F}, \mathcal{G} be torsion-free coherent analytic sheaves on a locally irreducible complex space X and let $\pi: Y \rightarrow X$ be a proper modification of X such that $\pi^T\mathcal{G}$ is locally free. Then*

$$\pi^T(\mathcal{F} \otimes \mathcal{G}) = \pi^T\mathcal{F} \otimes \pi^T\mathcal{G}$$

and there is a natural injection

$$\mathcal{F} \otimes \mathcal{G} \hookrightarrow \pi_*(\pi^T\mathcal{F} \otimes \pi^T\mathcal{G}).$$

Proof. — Note that Y is also locally irreducible. Consider the two natural surjections $\pi^*\mathcal{F} \rightarrow \pi^T\mathcal{F}$ and $\pi^*\mathcal{G} \rightarrow \pi^T\mathcal{G}$. These yield a natural surjection

$$\pi^*(\mathcal{F} \otimes \mathcal{G}) = \pi^*\mathcal{F} \otimes \pi^*\mathcal{G} \longrightarrow \pi^T\mathcal{F} \otimes \pi^T\mathcal{G}$$

which is an isomorphism on an open dense subset of Y . Since the tensor product of a torsion-free and a locally free sheaf is torsion-free,⁽³⁾ we obtain by use of Lemma 3.19 a natural isomorphism

$$(8.2) \quad \pi^T(\mathcal{F} \otimes \mathcal{G}) = \frac{\pi^*(\mathcal{F} \otimes \mathcal{G})}{\mathcal{I}(\pi^*(\mathcal{F} \otimes \mathcal{G}))} \xrightarrow{\sim} \pi^T\mathcal{F} \otimes \pi^T\mathcal{G}.$$

The second statement follows by taking the direct image of (8.2) under π and Theorem 4.1(i). □

This lemma and the projection formula gives directly the following corollary of Theorem 8.1.

THEOREM 8.3. — *Let X be a complex space of pure dimension n , \mathcal{K}_X the Grauert–Riemenschneider canonical sheaf on X and \mathcal{F} a torsion-free coherent analytic sheaf on X . Then there exists a resolution of singularities $\pi: M \rightarrow X$ and an effective divisor, $D \geq 0$, with support on the exceptional set of the resolution such that*

$$\pi^T(\mathcal{F} \otimes \mathcal{K}_X) \cong \pi^T\mathcal{F} \otimes \Omega_M^n(-D).$$

If \mathcal{F} is locally free, then

$$\pi_*(\pi^*\mathcal{F} \otimes \Omega_M^n) \cong \mathcal{F} \otimes \mathcal{K}_X \cong \pi_*(\pi^*\mathcal{F} \otimes \Omega_M^n(-D)).$$

(3) The tensor product of two torsion-free sheaves need not be torsion-free. E.g. let \mathcal{I} be the ideal sheaf generated of (z^2, zw) on $\mathbb{C}_{z,w}^2$ and \mathcal{J} be the ideal sheaf generated by (w^2, zw) . Then $z^2 \otimes w^2 - zw \otimes zw \in \mathcal{I} \otimes \mathcal{J}$ is not zero; yet, $z \cdot (z^2 \otimes w^2 - zw \otimes zw) = z^3 \otimes w^2 - z^2 \otimes zw^2 = 0$.

9. Non-analytic preimages and direct images

In this section, we will finally study non-analytic preimages of direct image sheaves and vice versa. For our purpose, the following definition is useful:

DEFINITION 9.1. — *Let \mathcal{F} be a sheaf on a complex space X . We say that \mathcal{F} satisfies the property (id) if the following holds: For any irreducible open set $W \subset X$ and sections $s, t \in \mathcal{F}(W)$, the equality $s = t$ on a non-empty open subset of W implies that $s = t$ on W .*

Property (id) means that the identity theorem generalizes to sections of \mathcal{F} . Actually, the identity theorem for irreducible complex spaces (cf. e.g. [10, Chap. 9, §1.3]) implies that the structure sheaf \mathcal{O}_X of a complex space satisfies (id). Moreover, we have:

LEMMA 9.2. — *Let X be a locally irreducible complex space. Then a coherent analytic sheaf \mathcal{F} on X satisfies the property (id) if and only if it is torsion-free.*

Proof. — Let \mathcal{F} be a torsion-free coherent sheaf on X and $F := L(\mathcal{F})$ the associated linear space. Then, by Remark 3.1, a section of \mathcal{F} is uniquely defined by its values on the locally irreducible primary component $E := \text{PC}(F)$ of F . I.e., for $W \subset X$ and $s, t \in \mathcal{F}(W) = \text{Hom}(F_W, W \times \mathbb{C})$, $s|_E = t|_E$ is equivalent to $s = t$. So, the desired property follows by the identity theorem applied to E .

Conversely, it is clear that sheaves with torsion on a locally irreducible space do not satisfy (id). \square

For non-coherent sheaves, the equivalence of Lemma 9.2 does not hold in general: The sheaf \mathcal{C} of continuous functions on an irreducible complex space X is torsion-free as \mathcal{O}_X -module sheaf, but it does not satisfy (id).

The property (id) is useful in the context of non-analytic preimages:

LEMMA 9.3. — *Let $\pi: Y \rightarrow X$ be a proper modification of a locally irreducible complex space X , and \mathcal{F} a sheaf on X satisfying (id). Then for $U \subset Y$ open:*

$$(9.1) \quad \pi^{-1}\mathcal{F}(U) = \varinjlim_{V \supset \pi(U)} \mathcal{F}(V),$$

where the limit runs over the open neighborhoods of $\pi(U)$.

Proof. — As X is locally irreducible, we can assume that X and Y are connected. Recall that $\pi^{-1}\mathcal{F}$ is the sheaf associated to the presheaf

$$U \mapsto F(U) := \varinjlim_{V \supset \pi(U)} \mathcal{F}(V)$$

where $U \subset Y$ is open and the limit runs over the open neighborhoods of $\pi(U)$. We have to show that the presheaf F is canonical (i.e., it is already a sheaf).

(i) Existence/Gluing-axiom: Let $U \subset Y$ be covered by open sets U_i , $i \in I$, and let $s_i \in F(U_i)$ satisfy $s_i = s_j$ on $U_{ij} := U_i \cap U_j$. By definition of the inductive limit, $s_i \in F(U_i)$ means there are an open set $V_i \supset \pi(U_i)$ and a section $f_i \in \mathcal{F}(V_i)$ with $s_i = [f_i]$ (s_i is represented by f_i). A priori, we just get $f_i = f_j$ on $\pi(U_{ij}) \subset V_{ij}$, where $V_{ij} = V_i \cap V_j$. Without loss of generality, we can assume that each connected component of V_{ij} contains an open subset of $\pi(U_{ij})$ (π is a modification). So, (id) for \mathcal{F} implies that $f_i = f_j$ on V_{ij} . As \mathcal{F} is a sheaf, there is a section $f \in \mathcal{F}(V)$ with $f|_{V_i} = f_i$, where $V := \bigcup_{i \in I} V_i \supset \pi(U)$. f represents an $s \in F(U)$ with $s|_{U_i} = s_i$.

(ii) Uniqueness-axiom: Let $U \subset Y$ be a connected open set, covered by open sets U_i , $i \in I$, and let $s, t \in F(U)$ satisfy $s = t$ on U_i for all $i \in I$. By definition of the inductive limit, there are a connected open set $V \supset \pi(U)$ and sections $f, g \in \mathcal{F}(V)$ with $s = [f]$ and $t = [g]$. We get $f = g$ on $\pi(U_i)$. Since $\pi(U_i)$ contains an open subset of V , (id) implies that $f = g$ on V . (We have not directly used the uniqueness-axiom for \mathcal{F} because it is contained in (id)). □

As a special case, we have:

LEMMA 9.4. — *Let $\pi: Y \rightarrow X$ be a proper modification of a locally irreducible complex space X , and \mathcal{G} a sheaf on Y satisfying (id). Then for $U \subset Y$ open:*

$$(9.2) \quad \pi^{-1}\pi_*\mathcal{G}(U) = \varinjlim_{V \supset \pi(U)} \mathcal{G}(\pi^{-1}(V)),$$

where the limit runs over the open neighborhoods of $\pi(U)$, and the canonical homomorphism $\pi^{-1}\pi_*\mathcal{G} \rightarrow \mathcal{G}$ is injective so that $\pi^{-1}\pi_*\mathcal{G}$ is a subsheaf of \mathcal{G} .

Proof. — As X is locally irreducible, we can assume that X and Y are connected. Here, $\pi^{-1}\pi_*\mathcal{G}$ is the sheaf associated to the presheaf

$$U \mapsto F(U) := \varinjlim_{V \supset \pi(U)} \mathcal{G}(\pi^{-1}(V))$$

where $U \subset Y$ is open and the limit runs over the open neighborhoods of $\pi(U)$. The canonical homomorphism $\pi^{-1}\pi_*\mathcal{G} \rightarrow \mathcal{G}$ is then induced by the restrictions $\mathcal{G}(\pi^{-1}(V)) \rightarrow \mathcal{G}(U)$.

By Lemma 9.3, F is canonical, i.e., (9.2) holds. It is now easy to see that the canonical homomorphism $\psi: \pi^{-1}\pi_*\mathcal{G} \rightarrow \mathcal{G}$ is injective. Let $s_x \in$

$(\pi^{-1}\pi_*\mathcal{G})_x$. Then (9.2) implies that s_x is represented by a section $s \in \mathcal{G}(U)$, where U is an open neighborhood of $K_x := \pi^{-1}(\pi(x))$. But our assumptions yield that K_x is connected, and so we can assume that U is a connected neighborhood of K_x . Assume that $\psi(s_x) = 0$. This means that s is vanishing on a neighborhood of the point x . But then $s = 0$ as U is connected (and \mathcal{G} satisfies (id)). \square

Lemma 9.4 allows for the following interpretation of $\pi^{-1}\pi_*\mathcal{G}$: The sections of $\pi^{-1}\pi_*\mathcal{G}$ are the sections of \mathcal{G} which extend along fibers of the modification π . This is of particular interest for the choices $\mathcal{G} = \mathcal{O}_M$ or $\mathcal{G} = \Omega_M^n$ when $\pi: M \rightarrow X$ is a resolution of singularities, giving the useful injections $\pi^{-1}\pi_*\mathcal{O}_M \hookrightarrow \mathcal{O}_M$ and $\pi^{-1}\pi_*\Omega_M^n \hookrightarrow \Omega_M^n$, respectively.

For the direct image of a non-analytic inverse image sheaf, Lemma 9.3 implies

$$(\pi_*\pi^{-1}\mathcal{F})(U) = (\pi^{-1}\mathcal{F})(\pi^{-1}(U)) = \varinjlim_{V \supset U} \mathcal{F}(V) = \mathcal{F}(U),$$

i.e., we obtain:

COROLLARY 9.5. — *Let $\pi: Y \rightarrow X$ be a proper modification of a locally irreducible complex space X , and \mathcal{F} a sheaf on X satisfying (id). Then*

$$\pi_*\pi^{-1}\mathcal{F} = \mathcal{F}.$$

BIBLIOGRAPHY

- [1] R. AXELSSON & J. MAGNÚSSON, “Complex analytic cones”, *Math. Ann.* **273** (1986), no. 4, p. 601-627.
- [2] ———, “A closure operation on complex analytic cones and torsion”, *Ann. Fac. Sci. Toulouse Math. (6)* **7** (1998), no. 1, p. 5-33.
- [3] J. BARSHAY, “Graded algebras of powers of ideals generated by A -sequences”, *J. Algebra* **25** (1973), p. 90-99.
- [4] J.-P. DEMAILLY, “Complex Analytic and Differential Geometry”, Institut Fourier, Université de Grenoble I. OpenContent AG-Book, <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>, 2012.
- [5] D. EISENBUD, *Commutative algebra*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995, With a view toward algebraic geometry, xvi+785 pages.
- [6] G. FISCHER, “Eine Charakterisierung von holomorphen Vektorraumbündeln”, *Bayer. Akad. Wiss. Math.-Natur. Kl. S.-B.* **1966** (1967), p. 101-107.
- [7] ———, “Lineare Faserräume und kohärente Modulgarben über komplexen Räumen”, *Arch. Math. (Basel)* **18** (1967), p. 609-617.
- [8] ———, *Complex analytic geometry*, Lecture Notes in Mathematics, Vol. 538, Springer-Verlag, Berlin, 1976, vii+201 pages.
- [9] H. GRAUERT, “Über Modifikationen und Exzeptionelle Analytische Mengen”, *Math. Ann.* **146** (1962), p. 331-368.

- [10] H. GRAUERT & R. REMMERT, *Coherent analytic sheaves*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 265, Springer-Verlag, Berlin, 1984, xviii+249 pages.
- [11] H. GRAUERT & O. RIEMENSCHNEIDER, “Verschwindungssätze für analytische Kohomologiegruppen auf komplexen Räumen”, *Invent. Math.* **11** (1970), p. 263-292.
- [12] A. GROTHENDIECK, “Techniques de construction en géométrie analytique. V: Fibres vectoriels, fibres projectifs, fibres en drapeaux.”, in *Sem. H. Cartan 13 (1960/61)*, no. 12, 1962, p. 15 (French).
- [13] C. HUNEKE, “On the symmetric and Rees algebra of an ideal generated by a d -sequence”, *J. Algebra* **62** (1980), no. 2, p. 268-275.
- [14] C. OKONEK, M. SCHNEIDER & H. SPINDLER, *Vector bundles on complex projective spaces*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2011, Corrected reprint of the 1988 edition, With an appendix by S. I. Gelfand, viii+239 pages.
- [15] T. PETERNELL, “Modifications”, in *Several complex variables, VII*, Encyclopaedia Math. Sci., vol. 74, Springer, Berlin, 1994, p. 285-317.
- [16] T. PETERNELL & R. REMMERT, “Differential calculus, holomorphic maps and linear structures on complex spaces”, in *Several complex variables, VII*, Encyclopaedia Math. Sci., vol. 74, Springer, Berlin, 1994, p. 97-144.
- [17] J. H. RABINOWITZ, “Moišezon spaces and positive coherent sheaves”, *Proc. Amer. Math. Soc.* **71** (1978), no. 2, p. 237-240.
- [18] ———, “On monoidal transformations of coherent sheaves”, *Proc. Amer. Math. Soc.* **74** (1979), no. 2, p. 389-390.
- [19] O. RIEMENSCHNEIDER, “Characterizing Moišezon spaces by almost positive coherent analytic sheaves”, *Math. Z.* **123** (1971), p. 263-284.
- [20] H. ROSSI, “Picard variety of an isolated singular point”, *Rice Univ. Studies* **54** (1968), no. 4, p. 63-73.
- [21] J. RUPPENTHAL, “ L^2 -theory for the $\bar{\partial}$ -operator on compact complex spaces”, *Duke Math. J.* **163** (2014), no. 15, p. 2887-2934.
- [22] M. SERA, “A generalization of Takegoshi’s relative vanishing theorem”, *J. Geom. Anal.* **26** (2016), no. 3, p. 1891-1912.
- [23] K. TAKEGOSHI, “Relative vanishing theorems in analytic spaces”, *Duke Math. J.* **52** (1985), no. 1, p. 273-279.

Manuscrit reçu le 20 avril 2015,
révisé le 24 mars 2016,
accepté le 6 avril 2016.

Jean RUPPENTHAL
ruppenthal@uni-wuppertal.de
University of Wuppertal
Department of Mathematics and Informatics
Gaußstr. 20, 42119 Wuppertal (Germany)

Martin SERA
sera@math.uni-wuppertal.de
University of Wuppertal
Department of Mathematics and Informatics
Gaußstr. 20, 42119 Wuppertal (Germany)