



ANNALES

DE

L'INSTITUT FOURIER

Xu WANG

A curvature formula associated to a family of pseudoconvex domains

Tome 67, n° 1 (2017), p. 269-313.

http://aif.cedram.org/item?id=AIF_2017__67_1_269_0



© Association des Annales de l'institut Fourier, 2017,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

A CURVATURE FORMULA ASSOCIATED TO A FAMILY OF PSEUDOCONVEX DOMAINS

by Xu WANG (*)

ABSTRACT. — We shall give a definition of the curvature operator for a family of weighted Bergman spaces $\{\mathcal{H}_t\}$ associated to a smooth family of smoothly bounded strongly pseudoconvex domains $\{D_t\}$. In order to study the “boundary term” in the curvature operator, we shall introduce the notion of geodesic curvature for the associated family of boundaries $\{\partial D_t\}$. As an application, we get a variation formula for the norms of Bergman projections of currents with compact support. A flatness criterion for $\{\mathcal{H}_t\}$ and its applications to triviality of fibrations are also given in this paper.

RÉSUMÉ. — Nous définissons l’opérateur de courbure pour une famille d’espaces de Bergman pondérés $\{\mathcal{H}_t\}$ associés à une famille lisse de domaines lisses bornés strictement pseudoconvexes $\{D_t\}$. Afin d’étudier le “terme au bord” dans l’opérateur de courbure, nous introduisons la notion de courbure géodésique pour la famille des bords associés. Comme application, nous obtenons une formule de variation pour les normes de projections de Bergman des courants à support compact. Un critère de platitude pour $\{\mathcal{H}_t\}$ et ses applications à la trivialité des fibrations sont également données dans cet article.

1. Introduction

In 2005, Berndtsson [3] found that the functional version of the classical Brunn–Minkowski inequality, i.e. the Prekopa theorem [43], can be seen as a special case of the subharmonicity property of the Bergman kernel (see [38, 39] for early results with different point of view). It opens another door (so called *complex Brunn–Minkowski theory*) of studying complex geometry by using the Brunn–Minkowski theory in convex geometry. Below,

Keywords: Brunn–Minkowski theory, Prekopa theorem, $\bar{\partial}$ -equation, Hörmander theory.
Math. classification: 32A25, 32L25, 32G05.

(*) Research supported by the Knut and Alice Wallenberg Foundation, and the China Postdoctoral Science Foundation.

we shall give a short account of the complex Brunn–Minkowski theory, and a simple example to show our motivation to write this paper.

Heuristically speaking, the Prekopa theorem can be seen as a version of inverse Hölder inequality. In [8], Berndtsson gave another form of the classical Hölder inequality:

THEOREM A (Hölder inequality). — *Let $\phi(t, x)$ be convex in t . Then*

$$(1.1) \quad t \mapsto \log \int_{\mathbb{R}^n} e^{\phi(t,x)} dx, \quad dx = dx^1 \wedge \cdots \wedge dx^n,$$

is convex (if the integral is convergent).

Proof. — The proof follows by differentiating with respect to t :

$$\left(\log \int e^\phi \right)_{tt} = \left(\int e^\phi \right)^{-2} \left(\int e^\phi \int (\phi_{tt} e^\phi + \phi_t^2 e^\phi) - \left(\int \phi_t e^\phi \right)^2 \right).$$

Notice that, by the Cauchy–Schwarz inequality, we have

$$(1.2) \quad \left(\int \phi_t e^\phi \right)^2 \leq \int e^\phi \int \phi_t^2 e^\phi.$$

Thus $\phi_{tt} \geq 0$ implies that $(\log \int e^\phi)_{tt} \geq 0$. □

The following result is due to Prekopa:

THEOREM B (Prekopa theorem). — *Let $\phi(t, x)$ be convex in t and x . Then*

$$(1.3) \quad t \mapsto -\log \int_{\mathbb{R}^n} e^{-\phi(t,x)} dx,$$

is convex.

There are many ways to prove Theorem B. A famous observation of Brascamp–Lieb (see [16]) is: one may use a weighted L^2 -estimates of the d -operator to prove Theorem B.

In [2], Berndtsson showed that one may also use Hörmander’s weighted L^2 -estimates of the $\bar{\partial}$ -operator (see [29]) to prove Theorem B. Moreover, in [3], he established the following complex version of Theorem B:

THEOREM C (Berndtsson’s theorem). — *Let $\phi(t, z)$ be a plurisubharmonic function on a pseudoconvex domain $D \subset \mathbb{C}_t^m \times \mathbb{C}_z^n$. Then*

$$(1.4) \quad (t, z) \mapsto \log K^t(z, z),$$

is plurisubharmonic or equal to $-\infty$ identically on D , where each K^t denotes the weighted Bergman kernel associated to the fibre $D_t := D \cap (\{t\} \times \mathbb{C}^n)$ and the weight $\phi^t := \phi|_{D_t}$.

In [3], Berndtsson gave two proofs of Theorem C. A crucial step in his first proof is also the Hörmander’s L^2 -estimates of the $\bar{\partial}$ -operator. Later in [4], he pointed out that it will be more natural to look at Theorem C as a curvature property of the direct image bundle (see [4, Theorem 1.1 and 1.2]). This is a milestone in the complex Brunn–Minkowski theory (see [12]).

The complex Brunn–Minkowski theory has proved to be very useful in several complex variables and complex geometry (see [5, 9, 10, 13, 14, 15] and references therein). This paper is an attempt to study the curvature formula of the direct image bundle associated to general Stein-fibrations (see [23, 40, 41, 34, 44, 52] for other generalizations and related results). The new results are the *boundary term* of the curvature formula and its relation with *interpolation family* of convex bodies.

Let us start by looking at an almost trivial case of Theorem B. Let

$$(1.5) \quad \mathcal{F} := \{[a(t), b(t)]\}_{0 \leq t \leq 1},$$

be a family of line segments. Let

$$D := \{(t, x) \in \mathbb{R}^2 : a(t) < x < b(t), 0 < t < 1\},$$

be the total space. Assume that $b(t) > a(t)$ for each $0 \leq t \leq 1$ and a, b are smooth on a neighborhood of $[0, 1]$. Put

$$\theta(a) = \frac{d^2 a}{dt^2}, \quad \theta(b) = -\frac{d^2 b}{dt^2}.$$

Let us introduce the following definitions:

DEFINITION 1.1. — We call θ the geodesic curvature of \mathcal{F} .

DEFINITION 1.2. — We call \mathcal{F} an interpolation family if $\theta \equiv 0$.

Remark. — \mathcal{F} is an interpolation family if and only if both a and b are affine functions.

DEFINITION 1.3. — We call \mathcal{F} a trivial family if there exists a real constant c such that for every $0 < t < 1$, $[a(t), b(t)] = [a(0), b(0)] + ct$.

Put

$$(1.6) \quad \phi(t, x) = 0, \text{ on } D, \quad \phi(t, x) = \infty, \text{ on } \mathbb{R}^2 \setminus D,$$

then convexity of D is equivalent to convexity of ϕ . Thus Theorem B implies that if D is convex then

$$(1.7) \quad \Phi : t \mapsto -\log(b(t) - a(t)) = -\log \int_{\mathbb{R}} e^{-\phi(t,x)} dx$$

is convex on $(0, 1)$. Moreover, by direct computation,

$$(1.8) \quad \ddot{\Phi} = \frac{(b-a)(\ddot{a}-\ddot{b})+(\dot{a}-\dot{b})^2}{(b-a)^2}, \quad \ddot{\Phi} := \frac{d^2\Phi}{dt^2}, \quad \dot{a} := \frac{da}{dt}.$$

We call

$$(1.9) \quad Geo := \frac{(b-a)(\ddot{a}-\ddot{b})}{(b-a)^2} = \frac{\theta(a)+\theta(b)}{b-a},$$

the geodesic term in $\ddot{\Phi}$ and

$$(1.10) \quad R := \frac{(\dot{a}-\dot{b})^2}{(b-a)^2}$$

the remaining term in $\ddot{\Phi}$. Thus we have:

PROPOSITION 1.4. — *The remaining term in $\ddot{\Phi}$ is always non-negative. Moreover, if the total space D is convex then the geodesic term in $\ddot{\Phi}$ is also non-negative.*

PROPOSITION 1.5. — *Assume that the total space D is convex then affine-ness of Φ is equivalent to triviality of \mathcal{F} .*

In this paper, we shall study the counterparts of the above notions in complex geometry. In the next section, we shall define the notion of geodesic curvature (see Definition 2.8) for a smooth family of smoothly bounded Stein domains (see Definition 2.5). Then Definition 2.10, *interpolation family of Stein domains*, can be seen as a generalization of Definition 1.2; and Definition 2.16, *trivial family of Stein domains*, can be seen as a generalization of Definition 1.3.

Our main result, Theorem 2.11, is a curvature formula associated to variation of Stein manifolds. Let $\{D_t\}$ be a smooth family of smoothly bounded n -dimensional Stein domains. Let \mathcal{L} be a holomorphic line bundle on the total space D . Let h be a smooth Hermitian metric on \mathcal{L} . We shall consider the associated family of Bergman spaces

$$\mathcal{H} := \{\mathcal{H}_t\},$$

where each \mathcal{H}_t is the space of L^2 -holomorphic $\mathcal{L}|_{D_t}$ -valued $(n, 0)$ -forms on D_t . Then Theorem 2.11 reads that:

MAIN THEOREM. — *Assume that \mathcal{L} is flat or relatively ample. Then the curvature of \mathcal{H} contains two terms: the geodesic term and the remaining term. The remaining term is always semi-positive in the sense of Nakano. Moreover, if the total space is pseudoconvex and \mathcal{L} is semi-positive on the total space then the geodesic term is also semi-positive in the sense of Nakano.*

Thus our main result can be seen as a generalization of Proposition 1.4. In Section 2.5, we shall give a definition of the holomorphic section of the dual of \mathcal{H} (see Definition 2.14). Then our main application, Corollary 2.15, can be stated as follows:

APPLICATION. — *If the total space is pseudoconvex and \mathcal{L} is non-negative on the total space then $\log \|f\|$ is plurisubharmonic for every holomorphic section f of the dual of \mathcal{H} .*

Corollary 2.15 can be seen as a generalization of Theorem C (see the remark behind [4, Theorem 1.1]). In Section 5.2, we shall also use Corollary 2.15 to study variation of the Bergman projection of currents with compact support. In particular, we shall give a variation formula for the derivatives of the Bergman kernel (see Theorem 5.2).

In Section 6, we shall discuss the counterparts of Proposition 1.5 in complex case. We shall show that under some assumptions (see Theorem 2.17), flatness of \mathcal{H} and triviality of D are equivalent. As a direct corollary, we shall give a triviality criterion for a class of holomorphic motions (see Corollary 2.18) of planar domains.

Acknowledgement

I would like to thank Bo-Yong Chen for introducing me this topic, Bo Berndtsson for many inspiring discussions relating the curvature of the direct images, his useful comments on this paper and suggestions on paper writing. Thanks are also given to Qing-Chun Ji for his constant support and encouragement.

I would also like to thank Laszlo Lempert for pointing out a mistake in the first version of this paper regarding the assumptions of Lemma 2.9. After the first version of this manuscript was completed, Laszlo Lempert kindly sent me a preprint [51] of Dat Vu Tran. In [51], Dat Vu Tran gave a careful study of curvature of Hilbert fields (see also [33]) and showed that the curvature operator associated to a family of planar domains with

pseudoconvex global space is semi-positive, which covers our main theorem in case each fibre is one-dimensional.

Last but not least, thanks are due to the referee for pointing out several inaccuracies in this paper and hopefully making this paper more readable.

2. Basic definitions and results

2.1. List of notations

1. $\pi : \mathcal{X} \rightarrow \mathbb{B}$ is a holomorphic submersion.
2. D is an open subset in \mathcal{X} , $D_t := D \cap \pi^{-1}(t)$.
3. \mathcal{L} is a holomorphic line bundle over \mathcal{X} and $L_t := \mathcal{L}|_{D_t}$.
4. \mathcal{H}_t is the space of L^2 holomorphic L_t -valued $(n, 0)$ -forms on D_t .
5. $\mathcal{H} := \{\mathcal{H}_t\}_{t \in \mathbb{B}}$.
6. i_t : the inclusion mapping $D_t \hookrightarrow D$.
7. t : coordinate system on \mathbb{B} , t^j components of t , $\partial_{t^j} = \partial/\partial t^j$.
8. $\sum d\bar{t}^j \otimes \bar{\partial}_{t^j}$ is the $\bar{\partial}$ -operator on \mathcal{H} .
9. $\sum dt^j \otimes D_{t^j}$ is the $(1, 0)$ -part of the Chern connection on \mathcal{H} .
10. $\Theta_{j\bar{k}} := [D_{t^j}, \bar{\partial}_{t^k}]$: curvature operators on \mathcal{H} .
11. $\theta_{j\bar{k}}(\rho)$: geodesic curvature of $\{\partial D_t\}$.
12. η, ζ, μ : local coordinates on a fixed fibre D_t , μ^λ : components of μ .
13. $e^{-\phi}$: local representative of a Hermitian metric h on a line bundle \mathcal{L} .
14. $\phi_j := \partial\phi/\partial t^j$, $\phi_{j\lambda} := \partial^2\phi/\partial t^j \partial \mu^\lambda$, $\phi_{\lambda\bar{\nu}} := \partial^2\phi/\partial \mu^\lambda \partial \bar{\mu}^\nu$.
15. $(\rho^{\lambda\nu})$, $(\phi^{\lambda\nu})$: inverse matrix of $(\rho_{\lambda\bar{\nu}})$, $(\phi_{\lambda\bar{\nu}})$ respectively.
16. $\delta_V := V \lrcorner$ means contraction of a form with a vector field V ;
17. $\alpha, \beta \in \mathbb{N}^n$, $|\alpha| := \alpha_1 + \dots + \alpha_n$, $f_\alpha := \partial^{|\alpha|} f / (\partial \mu^1)^{\alpha_1} \dots (\partial \mu^n)^{\alpha_n}$.

2.2. Set up

Let $\pi : \mathcal{X} \rightarrow \mathbb{B}$ be a holomorphic submersion from an $(n+m)$ -dimensional complex manifold \mathcal{X} to the unit ball \mathbb{B} in \mathbb{C}^m . Let D be an open subset of \mathcal{X} . Put

$$D_t = D \cap \pi^{-1}(t).$$

The following assumption will be used throughout this paper.

The restriction of π to the closure of D (with respect to the topology

(A1) *structure of the total space \mathcal{X}) is proper, and D_t is non-empty for every t in \mathbb{B} .*

Let \mathcal{L} be a holomorphic line bundle over \mathcal{X} . Put

$$L_t := \mathcal{L}|_{D_t}.$$

We shall consider the following family of vector spaces associated to $\{D_t\}_{t \in \mathbb{B}}$:

$$\mathcal{H}_t := \{f \in H^0(D_t, K_{D_t} + L_t) : \int_{D_t} i^{n^2} \{f, f\} < \infty\},$$

where $\{\cdot, \cdot\}$ is the canonical sesquilinear pairing (see [18, p. 268]) with respect to h . If we fix a local holomorphic frame, say e , of \mathcal{L} , and write $h(e, e) = e^{-\phi}$, then

$$\{f(z) \otimes e, f(z) \otimes e\} = e^{-\phi(z)} f(z) \wedge \overline{f(z)}.$$

Put

$$\mathcal{H} = \{\mathcal{H}_t\}_{t \in \mathbb{B}}.$$

Remark. — We know that each fibre \mathcal{H}_t is an infinite dimensional Hilbert space, moreover, the element in \mathcal{H}_t may not be smooth up to the boundary. Thus it is not easy to give a good definition of the curvature operator on \mathcal{H} (see [33] for a careful study of this subject). In order to make things easier, we shall only define the curvature operator on sections that are smooth up to the boundary.

Let i_t be the inclusion mapping

$$(2.1) \quad i_t : D_t \hookrightarrow D.$$

We shall introduce the following definition:

DEFINITION 2.1. — We call $u : t \mapsto u^t \in \mathcal{H}_t$ a smooth section of \mathcal{H} if there exists an \mathcal{L} -valued $(n, 0)$ -form, say \mathbf{u} , such that

$$(2.2) \quad i_t^* \mathbf{u} = u^t, \quad \forall t \in \mathbb{B},$$

and \mathbf{u} is smooth up to the boundary of D . We shall denote by $\Gamma(\mathcal{H})$ the space of smooth sections of \mathcal{H} .

Remark. — One may choose \mathbf{u} in the above definition such that \mathbf{u} is smooth on the total space \mathcal{X} .

In order to give a precise definition of the $\bar{\partial}$ -operator on \mathcal{H} , we shall introduce the following definition (see [7], see also [31, p. 46] or [53] for the admissible coordinate method):

DEFINITION 2.2. — We call a smooth \mathcal{L} -valued $(n, 0)$ -form \mathbf{u} on \mathcal{X} a representative of $u \in \Gamma(\mathcal{H})$ if $i_t^*(\mathbf{u}) = u^t$ for all $t \in \mathbb{B}$.

$\bar{\partial}$ -operator on \mathcal{H} . Let \mathbf{u} be a representative of $u \in \Gamma(\mathcal{H})$. Then one may write

$$(2.3) \quad \bar{\partial}\mathbf{u} = \sum dt^j \wedge \eta_j + d\bar{t}^j \wedge \nu_j.$$

Since $\mathbf{u} \wedge dt$, $dt := dt^1 \wedge \cdots \wedge dt^m$, does not depend on the choice of \mathbf{u} and

$$(2.4) \quad \bar{\partial}(\mathbf{u} \wedge dt) = \sum d\bar{t}^j \wedge \nu_j \wedge dt,$$

we know that each $i_t^* \nu_j$ does not depend on the choice of ν_j . Let us define

$$(2.5) \quad \bar{\partial}_{t^j} u : t \mapsto i_t^* \nu_j.$$

Then the $\bar{\partial}$ -operator on \mathcal{H} can be defined as

$$\bar{\partial}u := \sum d\bar{t}^j \otimes \bar{\partial}_{t^j} u.$$

From the definition, we know that $\bar{\partial}u \equiv 0$ on \mathbb{B} if and only if $\mathbf{u} \wedge dt$ is holomorphic on D . We shall introduce the following definition:

DEFINITION 2.3. — *Let u be a smooth section of \mathcal{H} . We call u a holomorphic section of \mathcal{H} if $\mathbf{u} \wedge dt$ is holomorphic on D .*

Chern connection on \mathcal{H} . We shall write the $(1, 0)$ -part of the Chern connection on \mathcal{H} as $\sum dt^j \otimes D_{t^j}$. By definition, each D_{t^j} should satisfy

$$(2.6) \quad \partial_{t^j}(u, v) = (D_{t^j} u, v) + (u, \bar{\partial}_{t^j} v), \quad \forall u, v \in \Gamma(\mathcal{H}),$$

where $\partial_{t^j} = \partial/\partial t^j$ and (\cdot, \cdot) denotes the inner product on \mathcal{H}_t .

DEFINITION 2.4. — *We say that the Chern connection is well defined on \mathcal{H} if for every $1 \leq j \leq m$, there exists a \mathbb{C} -linear operator $D_{t^j} : \Gamma(\mathcal{H}) \rightarrow \Gamma(\mathcal{H})$ such that (2.6) is true.*

Remark. — By using Hamilton’s theorem (see [28]), in Section 4 (see Proposition 4.1), we shall prove that the Chern connection is well defined on \mathcal{H} if D satisfies (A1) and the following assumption:

There is a smooth real valued function ρ on \mathcal{X} such that for each $t \in \mathbb{B}$, $\rho|_{\pi^{-1}(t)}$ is strictly plurisubharmonic in a neighborhood of the (A2) closure (with respect to the topology on $\pi^{-1}(t)$) of D_t . Moreover, $D_t = \{\rho < 0\} \cap \pi^{-1}(t)$ and the gradient of $\rho|_{\pi^{-1}(t)}$ has no zero point on ∂D_t .

Remark. — In Section 3.2, we shall prove that (A1) and (A2) together implies that every smooth vector field on the base \mathbb{B} has a smooth lift on \mathcal{X} that tangent to the boundary of D . Thus in this case, $\{D_t\}$ is locally trivial as a smooth family.

DEFINITION 2.5. — $\{D_t\}$ is said to be a smooth family of smoothly bounded Stein domains if D satisfies (A1) and (A2).

Assume that D satisfies (A1) and (A2). Then we can define the curvature operators on \mathcal{H} as follows:

$$(2.7) \quad \Theta_{j\bar{k}} u := [D_{tj}, \bar{\partial}_{tk}]u = D_{tj}\bar{\partial}_{tk}u - \bar{\partial}_{tk}D_{tj}u, \quad \forall u \in \Gamma(\mathcal{H}).$$

2.3. Previous results

We shall give a short account of Berndtsson’s results on “geodesic” formula for $\Theta_{j\bar{k}}$. Let us recall the following notions in his curvature formula.

Geodesic curvature in the space of Kähler metrics. Let us denote by $\Theta(\mathcal{L}, h)$ the curvature of (\mathcal{L}, h) . If we write h locally as $e^{-\phi}$ then we have

$$(2.8) \quad \Theta(\mathcal{L}, h) = \partial\bar{\partial}\phi.$$

If D is a product, say $D = D_0 \times \mathbb{B}$, and

$$(2.9) \quad i\partial\bar{\partial}\phi|_{D_0 \times \{t\}} > 0, \quad \forall t \in \mathbb{B},$$

then $\{i\partial\bar{\partial}\phi|_{D_0 \times \{t\}}\}$ can be seen as a family of Kähler metrics on D_0 . Assume further that $m = 1$. Then there exists a smooth function, say $c(\phi)$, such that

$$(2.10) \quad \frac{(i\partial\bar{\partial}\phi)^{n+1}}{(n+1)!} = c(\phi) \frac{(i\partial\bar{\partial}\phi)^n}{n!} \wedge idt \wedge d\bar{t}.$$

By [20, Proposition 3], if $\{i\partial\bar{\partial}\phi|_{D_0 \times \{t\}}\}$ is S^1 invariant then

The path $\{i\partial\bar{\partial}\phi|_{D_0 \times \{t\}}\}$ defines a geodesic in the space of Kähler metrics on D_0 if and only if $c(\phi) \equiv 0$.

In general, $c(\phi)$ is called the *geodesic curvature* in the space of Kähler metrics. The geodesic curvature plays a crucial role on variation of Kähler metrics on projective manifolds; see [20, 37, 45], to cite just a few. Another way to look at the geodesic curvature is to use the notion of *horizontal lift*.

Horizontal lift. The notion of horizontal lift is introduced by Siu in [47]. In [7], Berndtsson found that one may also define the notion of horizontal lift with respect to a relative Kähler form (a smooth d -closed $(1, 1)$ -form that is positive on each fibre). Let us recall his definition:

Let ω be a relative Kähler form on \mathcal{X} . A $(1, 0)$ -vector field V on \mathcal{X} is said to be horizontal with respect to ω if $\langle V, W \rangle_\omega = 0$ for every $(1, 0)$ -vector

field W such that $\pi_*(W) = 0$. Let v be a vector field on \mathbb{B} . We call V a horizontal lift of v if V is horizontal with respect to ω and $\pi_*V=v$.

By Berndtsson’s formula (see [7, p. 3]), if we write $\omega = i\partial\bar{\partial}\phi$ locally then for each $1 \leq j \leq m$, $\partial/\partial t^j$ has a unique horizontal lift, say V_j , as follows:

$$(2.11) \quad V_j = \partial/\partial t^j - \sum \phi_{j\bar{\nu}}\phi^{\bar{\nu}\lambda}\partial/\partial\mu^\lambda.$$

Moreover, if $m = 1$ then

$$(2.12) \quad \langle V_1, V_1 \rangle_{i\partial\bar{\partial}\phi} = c(\phi).$$

By this formula, it is natural to define the notion of geodesic curvature for general base dimension m and general fibration π .

Geodesic curvature of $\{h|_{L_t}\}$. Let us assume that

$$i\Theta(\mathcal{L}, h)|_{D_t} > 0, \forall t \in \mathbb{B}, \text{ or } \Theta(\mathcal{L}, h) \equiv 0 \text{ on } D,$$

In case $i\Theta(\mathcal{L}, h)|_{D_t} > 0, \forall t \in \mathbb{B}$, let V_j^h be the horizontal lift of the base vector fields $\partial/\partial t^j$ with respect to $i\Theta(\mathcal{L}, h)$. We shall define the geodesic curvature of $\{h|_{L_t}\}$ as:

$$(2.13) \quad c_{j\bar{k}}(h) := \langle V_j^h, V_k^h \rangle_{i\Theta(\mathcal{L}, h)}; \text{ and } c_{j\bar{k}}(h) := 0 \text{ if } \Theta(\mathcal{L}, h) \equiv 0 \text{ on } D.$$

Another notion in Berndtsson’s curvature formula is the following:

Remaining term in Hörmander’s L^2 -estimate (product case).

Assume that $D = D_0 \times \mathbb{B}$. Then the vector fields $\partial/\partial t^j$ are well defined on \mathcal{X} . Fix

$$(2.14) \quad u_j \in \Gamma(\mathcal{H}), \quad 1 \leq j \leq m, \quad (\text{see Definition 2.1}).$$

Let a be the L^2 -minimal solution of

$$\bar{\partial}^t(\cdot) = c,$$

where $\bar{\partial}^t$ denotes the Cauchy–Riemann operator on $D_t = D_0 \times \{t\}$ and

$$(2.15) \quad c := \sum (\partial/\partial t^j \lrcorner \Theta(\mathcal{L}, h))|_{D_t} \wedge u_j = \sum \bar{\partial}^t \phi_j \wedge u_j.$$

Then we have the following remaining term in Hörmander’s L^2 -estimate

$$R := \|c\|_{i\Theta(\mathcal{L}, h)|_{D_t}}^2 - \|a\|^2.$$

By Hörmander’s theorem (see [29]), if D_0 is pseudoconvex then R is non-negative. Thus in our case, R is always non-negative. Now we can state the following theorem of Berndtsson (see [3] and [4, Theorem 1.1]):

THEOREM 2.6. — Assume that $D = D_0 \times \mathbb{B}$, where D_0 is a strongly pseudoconvex domain with smooth boundary. If $i\Theta(\mathcal{L}, h)|_{D_0 \times \{t\}} > 0, \forall t \in \mathbb{B}$, then we have

$$(2.16) \quad \sum (\Theta_{j\bar{k}} u_j, u_k) = \sum (c_{j\bar{k}}(h) u_j, u_k) + R, \quad R \geq 0.$$

If $\Theta(\mathcal{L}, h) \equiv 0$ on D then

$$(2.17) \quad \sum (\Theta_{j\bar{k}} u_j, u_k) \equiv 0.$$

Remark. — (2.16) can be found in the proof of [4, Theorem 1.1]. (2.17) is a direct application of [4, (2.4)]. A special case of (2.16) for variation of the Bergman kernel is given in [3].

The counterpart of Theorem 2.6 for a proper Kähler fibration was given in [7] by Berndtsson. Let us recall the following notions in his formulae:

Remaining term in Hörmander’s L^2 -estimate (Polarized fibration). Let

$$(2.18) \quad \pi : D \rightarrow \mathbb{B},$$

be a proper holomorphic submersion. Assume that \mathcal{L} is a relatively ample line bundle on D , i.e. $i\Theta(\mathcal{L}, h)|_{D_t} > 0, \forall t \in \mathbb{B}$. By the Ohsawa–Takegoshi extension theorem (see [42, 48]), we know that the dimension of $\mathcal{H}_t := H^0(D_t, K_{D_t} + L_t)$ does not depend on t and our bundle \mathcal{H} is just the holomorphic vector bundle associated to the zero-th direct image sheaf $\pi_* \mathcal{O}(K_{D/\mathbb{B}} + \mathcal{L})$. For each j , let V_j^h be the horizontal lift of $\partial/\partial t^j$ with respect to $i\Theta(\mathcal{L}, h)$. Let us denote by

$$(2.19) \quad \kappa : T_{\mathbb{B}} \rightarrow \{H^1(D_t, T_{D_t})\}_{t \in \mathbb{B}},$$

the Kodaira–Spencer map associated to the holomorphic fibration π . By [30, Theorem 5.4], we know that each $(\bar{\partial} V_j^h)|_{D_t}$ can be seen as a representative of the Kodaira–Spencer class $\kappa(\partial/\partial t^j)$. For each $1 \leq j \leq m$, let u_j be a smooth section of \mathcal{H} . Put

$$b = \sum (\bar{\partial} V_j^h)|_{D_t} \lrcorner u_j.$$

Let a be the L^2 -minimal solution of

$$\bar{\partial}^t(\cdot) = \partial_\phi^t b,$$

where ∂_ϕ^t is the restriction of the $(1, 0)$ -part of the Chern connection of \mathcal{L} on D_t . Then we have the following Hörmander type-remaining term:

$$R^h := \|b\|_{i\Theta(\mathcal{L}, h)|_{D_t}}^2 - \|a\|^2 \geq 0.$$

Remaining term in Hörmander’s L^2 -estimate (Kähler fibration).

In this case, let us assume that the total space of the proper holomorphic submersion $\pi : D \rightarrow \mathbb{B}$ possesses a Kähler form ω . Let \mathcal{L} be a flat line bundle over D , i.e. $\Theta(\mathcal{L}, h) \equiv 0$ on D . By [4, Theorem 8.1], we know that $\pi_*\mathcal{O}(K_{D/\mathbb{B}} + \mathcal{L})$ is locally free. Let \mathcal{H} be the associated vector bundle. For each j , let u_j be a smooth section of \mathcal{H} . Put

$$b = \sum (\bar{\partial}V_j^\omega)|_{D_t \dashv} u_j,$$

where each V_j^ω is the horizontal lift of $\partial/\partial t^j$ with respect to the Kähler form ω . Consider

$$\bar{\partial}^t(a) = \partial_\phi^t b,$$

where a is the L^2 -minimal solution. Then the associated Hörmander remaining term is

$$R^\omega := \|b\|_{\omega|_{D_t}}^2 - \|a\|^2 \geq 0.$$

Now we can state the following theorem of Berndtsson (see [7]):

THEOREM 2.7. — *Let $\pi : D \rightarrow \mathbb{B}$ be a proper holomorphic submersion. Let (\mathcal{L}, h) be a holomorphic line bundle over D . If \mathcal{L} is relatively ample then we have*

$$(2.20) \quad \sum (\Theta_{j\bar{k}} u_j, u_k) = \sum (c_{j\bar{k}}(h) u_j, u_k) + R^h.$$

If \mathcal{L} is flat then

$$(2.21) \quad \sum (\Theta_{j\bar{k}} u_j, u_k) \equiv R^\omega.$$

Remark. — If \mathcal{L} is trivial then (2.21) is just Griffiths’ formula (see [25, p. 33]). In general, if the total space D is Kähler and there is a smooth Hermitian metric on \mathcal{L} with non-negative curvature then by [4, Theorem 1.2], we know that

$$\sum (\Theta_{j\bar{k}} u_j, u_k) \geq 0.$$

If the boundary of each fibre is non-empty then, in general, the boundary term should also appear in the curvature formula (see [39]). Our main result is a study of the curvature of \mathcal{H} for fibrations with boundary.

2.4. Basic notions for fibrations with boundary

Let $D = \{D_t\}_{t \in \mathbb{B}}$ be a smooth family of smoothly bounded Stein domains (see Definition 2.5). We shall define the notion of “geodesic curvature” of $\{\partial D_t\}$ by using the notion of *horizontal lift with respect to the Levi-form on the boundary* of D . Let ρ be the defining function in (A2). We call an

(1, 0)-tangent vector field V on ∂D horizontal with respect to the Levi-form if

$$\langle V, W \rangle_{i\partial\bar{\partial}\rho} = 0, \text{ on } \partial D,$$

for every (1, 0)-tangent vector field W on ∂D such that $\pi_*(W) = 0$.

Remark. — From the above definition, the notion of horizontal lift with respect to the Levi-form on the boundary is compatible with the usual notion of horizontal lift if we only consider the category of tangent vector fields on ∂D . Same as before, we shall also define the notion of geodesic curvature in the following sense:

DEFINITION 2.8. — Assume that D satisfies (A1) and (A2) and for each j , $\partial/\partial t^j$ has a horizontal lift to ∂D with respect to the Levi-form $i\partial\bar{\partial}\rho$ on ∂D . Then we call

$$(2.22) \quad \theta_{j\bar{k}}(\rho) := \langle V_j^\rho, V_k^\rho \rangle_{i\partial\bar{\partial}\rho},$$

the geodesic curvature of $\{\partial D_t\}_{t \in \mathbb{B}}$ with respect to the Levi form $i\partial\bar{\partial}\rho$ on ∂D .

Now a natural question is whether each base vector field has a horizontal lift (with respect to the Levi-form) to ∂D or not. We have the following lemma:

LEMMA 2.9 (Key Lemma). — Assume that D satisfies (A1) and (A2). Put

$$(2.23) \quad \omega := i\partial\bar{\partial}(-\log -\rho),$$

where ρ is the defining function in (A2). For each j , let V_j be the horizontal lift (on D) of $\partial/\partial t^j$ with respect to ω . Then each V_j is smooth up to the boundary of D and $V_j|_{\partial D}$ is horizontal with respect to the Levi form $i\partial\bar{\partial}\rho$ on ∂D . In particular, every smooth base vector field has a unique smooth horizontal lift with respect to the Levi form and the geodesic curvature $\theta_{j\bar{k}}(\rho)$ is well defined on ∂D .

As a generalization of Definition 1.2, we shall introduce the following definition:

DEFINITION 2.10. — Assume that D satisfies (A1) and (A2). We call $\{D_t\}_{t \in \mathbb{B}}$ an interpolation family in \mathcal{X} if $\theta_{j\bar{k}}(\rho) \equiv 0$ on ∂D .

Remaining term in Hörmander’s L^2 -estimate (fibration with boundary). For each $1 \leq j \leq m$, let u_j be a smooth section of \mathcal{H} (see Definition 2.1). Put

$$(2.24) \quad c = \sum (V_j \lrcorner \Theta(\mathcal{L}, h))|_{D_t} \wedge u_j, \quad b = \sum (\bar{\partial}V_j)|_{D_t} \lrcorner u_j,$$

where each V_j is the vector field in Lemma 2.9. Let a be the L^2 -minimal solution of

$$\bar{\partial}^t(\cdot) = \partial_\phi^t b + c,$$

in $L^2(D_t, K_{D_t} + L_t)$. Put

$$\omega^t := i\bar{\partial}\bar{\partial}(-\log -\rho)|_{D_t}.$$

Then we shall define

$$(2.25) \quad R := \|c\|_{i\Theta(\mathcal{L}, h)|_{D_t}}^2 + \|b\|_{\omega^t}^2 - \|a\|_{\omega^t}^2,$$

if $i\Theta(\mathcal{L}, h)|_{D_t} > 0$; and define

$$R := \|b\|_{\omega^t}^2 - \|a\|_{\omega^t}^2,$$

if $\Theta(\mathcal{L}, h) \equiv 0$. We shall show in Theorem A.5 that R is always non-negative.

2.5. Main theorem

THEOREM 2.11. — Assume that D satisfies (A1) and (A2). If

$$(2.26) \quad i\Theta(\mathcal{L}, h)|_{D_t} > 0, \quad \forall t \in \mathbb{B}, \quad \text{or } \Theta(\mathcal{L}, h) \equiv 0 \text{ on } D,$$

then, using the above notation, see (2.7), (2.13), (2.22), (2.25), we have the following curvature formula of \mathcal{H} :

$$(2.27) \quad \sum (\Theta_{j\bar{k}} u_j, u_k) = \sum \int_{\partial D_t} \theta_{j\bar{k}}(\rho) \langle u_j, u_k \rangle d\sigma + \sum (c_{j\bar{k}}(h) u_j, u_k) + R,$$

where $\langle \cdot, \cdot \rangle$ denotes the point-wise inner product with respect to $i\bar{\partial}\bar{\partial}\rho|_{D_t}$ and h , and the surface measure $d\sigma$ with respect to $i\bar{\partial}\bar{\partial}\rho|_{D_t}$ is defined as

$$d\sigma := \frac{\sum \rho_{\bar{\lambda}} \rho^{\bar{\lambda}\nu} \partial/\partial\mu^\nu}{\sum \rho_{\bar{\lambda}} \rho^{\bar{\lambda}\nu} \rho_\nu} \lrcorner \frac{(i\bar{\partial}\bar{\partial}\rho|_{D_t})^n}{n!}.$$

2.6. Applications

We shall show how to use our main theorem to study the complex geometry counterparts of Theorem 1.4. Let us give some positive-curvature criterion of \mathcal{H} first. Recall that, \mathcal{H} is said to be semi-positive in the sense of Nakano if $\sum(\Theta_{j\bar{k}}u_j, u_k) \geq 0$, for all smooth sections u_1, \dots, u_m of \mathcal{H} . As a direct consequence of our main theorem, we shall prove that:

COROLLARY 2.12. — *Assume that D satisfies (A1) and (A2). If D is Stein and $i\Theta(\mathcal{L}, h) \geq 0$ on D then \mathcal{H} is semi-positive in the sense of Nakano.*

Another very useful notion of positivity is the Griffiths positivity. Recall that \mathcal{H} is said to be Griffiths semi-positive if $\sum(\Theta_{j\bar{k}}u, u)\xi_j\bar{\xi}_k \geq 0$, for every smooth section u of \mathcal{H} and every $\xi \in \mathbb{C}^m$. It is known that a finite rank vector bundle is Griffiths semi-positive if and only if its dual bundle is Griffiths semi-negative. Moreover, the following is true:

Criterion for Griffiths semi-positivity. — A finite rank vector bundle is Griffiths semi-positive if and only if the log-norm of the holomorphic sections of its dual bundle are plurisubharmonic.

Then a natural question is whether there is a similar criterion in our case, i.e. for the infinite rank vector bundle \mathcal{H} . As a first step, we have to define the notion of the holomorphic section of the dual of \mathcal{H} .

DEFINITION 2.13. — *For each $t \in \mathbb{B}$, let f^t be a \mathbb{C} -linear mapping from \mathcal{H}_t to \mathbb{C} . We call $f : t \mapsto f^t$ a smooth section of the dual of \mathcal{H} if there exists a smooth section, say $P(f)$, of \mathcal{H} such that*

$$(2.28) \quad f^t(u^t) = (u^t, P(f)^t),$$

for every $u^t \in \mathcal{H}_t$ and every $t \in \mathbb{B}$. We shall write the norm of f^t as $\|f^t\| := \|P(f)^t\|$.

DEFINITION 2.14. — *Let $f : t \mapsto f^t$ be a smooth section of the dual of \mathcal{H} . We call f a holomorphic section if*

$$(2.29) \quad t \mapsto f^t(u^t)$$

is holomorphic for every holomorphic section u of \mathcal{H} .

Remark. — Inspired by [14], we shall give a careful study of those holomorphic sections of the dual of \mathcal{H} defined by a family currents with compact support in fibres.

Now we are ready to state the main application of our main theorem:

COROLLARY 2.15. — Assume that D satisfies (A1) and (A2). If D is Stein and $i\Theta(\mathcal{L}, h) \geq 0$ on D then

$$(2.30) \quad \log \|f\| : t \mapsto \log \|f^t\|$$

is plurisubharmonic for every holomorphic section f of the dual of \mathcal{H} .

Remark. — If we choose f^t as a fixed Dirac measure then we get the plurisubharmonicity of the Bergman kernel [3]. In Section 5, we shall also use Corollary 2.15 to study variation of the derivatives of the Bergman kernel.

Triviality and flatness. In case every fibre D_t is compact without boundary, we know that the criterions for flatness of \mathcal{H} are quite useful in the study of the uniqueness problems of extremal Kähler metrics (see [5, 10]). In our case, we call \mathcal{H} a flat bundle if

$$(2.31) \quad \Theta_{j\bar{k}} u \equiv 0, \quad \forall 1 \leq j, k \leq m.$$

for every smooth section u of \mathcal{H} . From Proposition 1.4, one may guess that flatness of \mathcal{H} should be related to triviality of the fibration π . Let us introduce the following definition as a generalization of Definition 1.3.

DEFINITION 2.16. — Assume that D satisfies (A1) and (A2). We call $\{D_t\}_{t \in \mathbb{B}}$ a trivial family, or D is trivial, if there exists a biholomorphic mapping $\Phi : D_0 \times \mathbb{B} \rightarrow D$ such that

$$(2.32) \quad \Phi(D_0 \times \{t\}) = D_t, \quad \forall t \in \mathbb{B},$$

and $\Phi_*(\partial/\partial t^j)$ extends to a smooth $(1, 0)$ -vector field on \mathcal{X} for every $1 \leq j \leq m$.

The following theorem can be seen as a generalization of Proposition 1.5:

THEOREM 2.17. — Assume that D satisfies (A1) and (A2). Assume further that the total space D is Stein. If $K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}$ is trivial on each fibre of π and $\Theta(\mathcal{L}, h) \equiv 0$ on D then flatness of \mathcal{H} and triviality of D are equivalent.

As a direct corollary of Theorem 2.17, we have

COROLLARY 2.18. — Let D_0 be a smooth domain in \mathbb{C} . Let

$$(2.33) \quad F : (t, z) \mapsto (t, z + a(t)\bar{z})$$

be a mapping from $\mathbb{B} \times D_0$ to $\mathbb{B} \times \mathbb{C}$. Assume that a is holomorphic on \mathbb{B} and

$$(2.34) \quad |a| < 1, \text{ on } \mathbb{B}, \quad a(0) = 0.$$

Then $\{F(\{t\} \times D_0)\}_{t \in \mathbb{B}}$ is a trivial family if and only if $a \equiv 0$ on \mathbb{B} .

Remark. — One may also give a direct proof of Corollary 2.18 by introducing the notion of Kodaira–Spencer class for deformations with boundary, we leave it to the interested reader.

3. Geodesic curvature and interpolation family

In this section, we shall give two proofs of Lemma 2.9 and show that our Definition of interpolation family (see Definition 2.10) is compatible with the usual definition of interpolation family of Hermitian norms on \mathbb{C}^n for $n \geq 1$.

Relation with Levi-flatness. By the definition of the geodesic curvature $\theta_{j\bar{k}}(\rho)$ (see Definition 2.8) of $\{\partial D_t\}$, we have:

Assume that every fibre D_t is one dimensional. Then $\{D_t\}$ is an interpolation family if and only if the boundary of D is Levi flat.

For higher fibre dimension case, the criterion for interpolation family (see [45] and references therein) is not so obvious. We will give a study of it in Section 3.3. Let us prove our Key Lemma first.

3.1. First proof of Lemma 2.9

Since V_j is a lift of $\partial/\partial t^j$, locally one may write,

$$(3.1) \quad V_j = \partial/\partial t^j - \sum v_j^\lambda \partial/\partial \mu^\lambda.$$

Put $\psi = -\log -\rho$. By definition, we know that each V_j is determined by

$$\langle V_j, \partial/\partial \mu^\nu \rangle_{i\partial\bar{\partial}\psi} \equiv 0, \text{ on } D, \forall 1 \leq \nu \leq n,$$

thus

$$(3.2) \quad v_j^\lambda = \sum \psi_{j\bar{\nu}} \psi^{\bar{\nu}\lambda}, \text{ on } D.$$

Fibre dimension one case. If $n = 1$, by direct computation, we have

$$(3.3) \quad V_j := \frac{\partial}{\partial t^j} - \frac{\rho_j \rho_{\bar{\mu}} - \rho \rho_{j\bar{\mu}}}{|\rho_\mu|^2 - \rho \rho_{\mu\bar{\mu}}} \frac{\partial}{\partial \mu}.$$

By Assumption (A2), ρ_μ has no zero point near the boundary and $\rho_{\mu\bar{\mu}} > 0$ near the boundary, thus V_j is smooth up to the boundary of D . Furthermore, (3.3) implies that $V_j(\rho) = 0$ on $\{\rho = 0\}$. Notice that, in case $n = 1$,

every tangent vector field on ∂D is horizontal with respect to the Levi-form on ∂D . Thus we know that if $n = 1$ then $V_j|_{\partial D}$ is the horizontal lift of $\partial/\partial t^j$ with respect to the Levi-form.

General case. The general case can also be proved by direct computation (see the second proof below). But there is also a simple proof as follows: If $n \geq 2$, fix $x_0 \in \partial D_0$, then by Assumption (A2), $(\rho_{\lambda\bar{\nu}})(x_0)$ is a positive definite matrix, thus one may choose local coordinates around x_0 such that

$$(\rho_{\lambda\bar{\nu}}(x_0)) = I_n, \rho_\nu(x_0) = 0, \forall \nu \geq 2,$$

where I_n is the identity matrix. Now we have

$$v_j^1(x_0) = \frac{\rho_j \rho_{\bar{1}} - \rho \rho_{j\bar{1}}}{|\rho_1|^2 - \rho}(x_0) = \frac{\rho_j}{\rho_1}(x_0), v_j^\lambda(x_0) = \rho_{j\bar{\lambda}}(x_0), \forall \lambda \geq 2.$$

By Assumption (A2), we know that $\rho_1(x_0) \neq 0$, thus V_j is smooth up to the boundary and

$$V_j(\rho)(x_0) = \rho_j(x_0) - \sum v_j^\lambda \rho_\lambda(x_0) = \rho_j(x_0) - \rho_j(x_0) = 0.$$

Moreover, we have

$$\langle V_j, \partial/\partial \mu^\lambda \rangle_{i\partial\bar{\partial}\rho}(x_0) = \rho_{j\bar{\lambda}}(x_0) - v_j^\lambda(x_0) = 0, \forall \lambda \geq 2,$$

which implies that each V_j is horizontal with respect to the Levi form. The proof is complete.

3.2. Second proof of Lemma 2.9

In this subsection, we will give an explicit formula for each V_j . Here we shall use some computations from [17]. Put

$$\rho^\alpha = \sum \rho^{\bar{\beta}\alpha} \rho_{\bar{\beta}}, \quad |\partial\rho|^2 = \sum |\rho_\alpha|^2.$$

By (3.1) and (3.2), we have

$$(3.4) \quad V_j = \partial/\partial t^j - \sum \psi_{j\bar{\nu}} \psi^{\bar{\nu}\lambda} \partial/\partial \mu^\lambda,$$

where $\psi = -\log -\rho$. One may verify that (see [17])

$$(3.5) \quad \psi^{\bar{\alpha}\beta} = (-\rho) \left(\rho^{\bar{\alpha}\beta} + \frac{\rho^{\bar{\alpha}} \rho^\beta}{\rho - |\partial\rho|^2} \right).$$

By direct computation, we have

$$(3.6) \quad \sum \psi_{j\bar{\alpha}} \psi^{\bar{\alpha}\beta} = -\frac{\rho^\beta \rho_j}{\rho - |\partial\rho|^2} + \sum \left(\rho^{\bar{\alpha}\beta} \rho_{j\bar{\alpha}} + \frac{\rho^{\bar{\alpha}} \rho^\beta \rho_{j\bar{\alpha}}}{\rho - |\partial\rho|^2} \right).$$

From (3.6), we know that each V_j is smooth up to the boundary of D and is tangent to the boundary of D . By a direct computation, we also have that each V_j is horizontal with respect to the Levi-form of the boundary of D . Thus the proof of Lemma 2.9 is complete.

3.3. Geodesic curvature for $\{\partial D_t\}$

Denote by \hat{V}_j the horizontal lift of $\partial/\partial t^j$ with respect to $i\partial\bar{\partial}\rho$. By the proof of (3.2), we have

$$(3.7) \quad \hat{V}_j = \frac{\partial}{\partial t^j} - \sum \rho_{j\bar{\alpha}} \rho^{\bar{\alpha}\beta} \frac{\partial}{\partial \mu^{\bar{\beta}}}.$$

By (3.4) and (3.6) and a direct computation, we get

$$(3.8) \quad \theta_{j\bar{k}}(\rho) = \langle V_j, V_k \rangle_{i\partial\bar{\partial}\rho} = c_{j\bar{k}}(\rho) + \frac{|\partial\rho|^2 \hat{V}_j(\rho) \overline{\hat{V}_k(\rho)}}{(\rho - |\partial\rho|^2)^2},$$

where

$$(3.9) \quad c_{j\bar{k}}(\rho) := \langle \hat{V}_j, \hat{V}_k \rangle_{i\partial\bar{\partial}\rho}.$$

Thus we have:

PROPOSITION 3.1. — *Let $\{D_t\}$ be a smooth family of smoothly bounded Stein domains. Then*

$$(3.10) \quad \sum \theta_{j\bar{k}}(\rho) \xi^j \bar{\xi}^k \geq \sum c_{j\bar{k}}(\rho) \xi^j \bar{\xi}^k, \quad \forall \xi \in \mathbb{C}^m,$$

on ∂D . Moreover, $\theta_{j\bar{k}}(\rho) \equiv c_{j\bar{k}}(\rho)$ if and only if each \hat{V}_j is tangent to the boundary of D .

3.4. Relation with interpolation of norms

Let h be a smooth Hermitian norm on the trivial vector bundle $\mathbb{B} \times \mathbb{C}^n$. Then for each $t \in \mathbb{B}$, $h^t := h|_{t \times \mathbb{C}^n}$ defines a Hermitian norm on \mathbb{C}^n . It is known that $\{h^t\}$ defines an interpolation family if and only if the curvature of h vanishes identically on \mathbb{B} (see Semmes [45]). Denote by N_t the unit ball in \mathbb{C}^n defined by h^t . We shall prove that:

PROPOSITION 3.2. — *$\{h^t\}$ defines an interpolation family if and only if the geodesic curvature of $\{\partial N_t\}$ vanishes identically.*

Proof. — Let us write

$$h^t(z) = \sum h_{\alpha\bar{\beta}}(t) z^\alpha \bar{z}^\beta.$$

By definition,

$$\rho(t, z) := h^t(z) - 1$$

is a defining function for $\{\partial N_t\}$. By direct computation, we have that $\hat{V}_j(\rho)$ vanishes identically. Thus by Proposition 3.1, the geodesic curvature, $\theta_{j\bar{k}}(\rho)$, of $\{\partial N_t\}$ is equal to $c_{j\bar{k}}(\rho)$. By (3.9),

$$(3.11) \quad c_{j\bar{k}}(\rho) = \langle \hat{V}_j, \hat{V}_{\bar{k}} \rangle_{i\partial\bar{\partial}\rho} = \rho_{j\bar{k}} - \sum \rho_{j\bar{\alpha}} \rho^{\bar{\alpha}\beta} \rho_{\bar{k}\beta},$$

thus we have

$$(3.12) \quad \theta_{j\bar{k}}(\rho) = \sum \left(h_{\alpha\bar{\beta},j\bar{k}} - \sum h_{\alpha\bar{\lambda},j} h^{\bar{\lambda}\nu} h_{\nu\bar{\beta},\bar{k}} \right) z^\alpha \bar{z}^\beta.$$

Thus $\theta_{j\bar{k}}(\rho)$ vanishes identically if and only if

$$(3.13) \quad h_{\alpha\bar{\beta},j\bar{k}} - \sum h_{\alpha\bar{\lambda},j} h^{\bar{\lambda}\nu} h_{\nu\bar{\beta},\bar{k}} \equiv 0, \quad \text{on } \mathbb{B}.$$

Notice that (3.13) is equivalent to that the curvature of h vanishes identically. The proof is complete. □

4. Curvature formula

4.1. Definition of the Chern connection

By Definition 2.4, it suffices to find a linear operator D_{tj} from $\Gamma(\mathcal{H})$ to $\Gamma(\mathcal{H})$ such that

$$(4.1) \quad \partial_{tj}(u, v) = (D_{tj}u, v) + (u, \bar{\partial}_{tj}v), \quad \forall u, v \in \Gamma(\mathcal{H}),$$

is true. By Definition 2.2, the left hand side of (4.1) can be written as

$$(4.2) \quad \partial_{tj}(\pi_*(c_n\{\mathbf{u}, \mathbf{v}\})),$$

where \mathbf{u}, \mathbf{v} are arbitrary representatives (see Definition 2.2) of u, v and $c_n = i^{n^2}$ such that $c_n\{\mathbf{u}, \mathbf{u}\}$ is a positive (n, n) -form on the total space.

Assume that D satisfies (A1) and (A2). Let V_j be the vector fields in Lemma 2.9. Since $V_j(\rho) = 0$ on ∂D , by Corollary A.4, we have

$$(4.3) \quad \partial_{tj}(\pi_*\{\mathbf{u}, \mathbf{v}\}) = \pi_*(L_{V_j}\{\mathbf{u}, \mathbf{v}\}).$$

Let $d^{\mathcal{L}}$ be the Chern connection on \mathcal{L} . Then we have

$$d\{\mathbf{u}, \mathbf{v}\} = \{d^{\mathcal{L}}\mathbf{u}, \mathbf{v}\} + (-1)^n \{\mathbf{u}, d^{\mathcal{L}}\mathbf{v}\}.$$

Using Cartan’s formula,

$$L_{V_j} = d\delta_{V_j} + \delta_{V_j}d,$$

we get that

$$L_{V_j}\{\mathbf{u}, \mathbf{v}\} = \{L_j\mathbf{u}, \mathbf{v}\} + \{\mathbf{u}, L_{\bar{j}}\mathbf{v}\},$$

where

$$L_j := d^{\mathcal{L}}\delta_{V_j} + \delta_{V_j}d^{\mathcal{L}},$$

and

$$L_{\bar{j}} := d^{\mathcal{L}}\delta_{\bar{V}_j} + \delta_{\bar{V}_j}d^{\mathcal{L}}.$$

Since \mathbf{v} is an $(n, 0)$ -form, we have

$$(4.4) \quad L_{\bar{j}}\mathbf{v} = \delta_{\bar{V}_j}\bar{\partial}\mathbf{v},$$

By (2.5), we know that $L_{\bar{j}}\mathbf{v}$ is a representative of $\bar{\partial}_{t^j}v$. Thus we have

$$(4.5) \quad \partial_{t^j}(u, v) = \pi_*(c_n\{L_j\mathbf{u}, \mathbf{v}\}) + (u, \bar{\partial}_{t^j}v), \quad \forall u, v \in \Gamma(\mathcal{H}).$$

Notice that the $(n, 0)$ -part of $L_j\mathbf{u}$ can be written as

$$(4.6) \quad (\partial_\phi\delta_{V_j} + \delta_{V_j}\partial_\phi)\mathbf{u},$$

where ∂_ϕ denotes the $(1, 0)$ -component of $d^{\mathcal{L}}$. Thus we have

$$\pi_*(c_n\{L_j\mathbf{u}, \mathbf{v}\})(t) = (i_t^*(\partial_\phi\delta_{V_j} + \delta_{V_j}\partial_\phi)\mathbf{u}, v^t).$$

Assumption (A2) implies that

$$(4.7) \quad \{v^t : v \in \Gamma(\mathcal{H})\}.$$

is dense in the Hilbert space \mathcal{H}_t (see the proof of Lemma 4.7 below).

Thus there is a *unique* element, say σ^t , in \mathcal{H}_t such that

$$(4.8) \quad (i_t^*(\partial_\phi\delta_{V_j} + \delta_{V_j}\partial_\phi)\mathbf{u}, v^t) = (\sigma^t, v^t),$$

which implies that there is a *unique* element, σ^t , in \mathcal{H}_t such that

$$(4.9) \quad \partial_{t^j}(u, v) = (\sigma^t, v^t) + (u, \bar{\partial}_{t^j}v), \quad \forall u, v \in \Gamma(\mathcal{H}).$$

Thus by Definition 2.4, we know that: \mathcal{H} has a Chern connection if and only if

$$\sigma : t \rightarrow \sigma^t,$$

defines a smooth section of \mathcal{H} , i.e.

$$\sigma \in \Gamma(\mathcal{H}).$$

By (4.8), σ^t is the Bergman projection to \mathcal{H}_t of $i_t^*((\partial_\phi\delta_{V_j} + \delta_{V_j}\partial_\phi)\mathbf{u})$. Thus by Hamilton’s theorem (see [24, 27, 28] or Appendix A.2), if $\{D_t\}$ is a smooth family of smoothly bounded Stein domains then $\sigma \in \Gamma(\mathcal{H})$ and

$$(4.10) \quad D_{t^j}u = \sigma.$$

Thus we have:

PROPOSITION 4.1. — *If D satisfies (A1) and (A2) then the Chern connection is well defined on \mathcal{H} .*

Now we are ready to compute the curvature of the Chern connection of \mathcal{H} . First we shall show how to get a curvature formula for holomorphic sections of \mathcal{H} .

4.2. Curvature formula for holomorphic sections

Let u_1, \dots, u_m be holomorphic sections (see Definition 2.3) of \mathcal{H} . By definition of the Chern connection and (2.7), we have

$$(4.11) \quad (\Theta_{j\bar{k}} u_j, u_k) = (D_{t^j} u_j, D_{t^k} u_k) - (u_j, u_k)_{j\bar{k}}.$$

By (4.5), we have

$$(4.12) \quad (u_j, u_k)_{j\bar{k}} = \pi_*(c_n\{L_j \mathbf{u}_j, L_k \mathbf{u}_k\}) + \pi_*(c_n\{L_{\bar{k}} L_j \mathbf{u}_j, \mathbf{u}_k\}).$$

Since each u_j is a holomorphic section, we have $i_t^*(L_{\bar{k}} \mathbf{u}_j) = (\bar{\partial}_{t^k} u_j)(t) \equiv 0$. Thus

$$(4.13) \quad \pi_*\{L_j L_{\bar{k}} \mathbf{u}_j, \mathbf{u}_k\} = \partial_{t^j} \pi_*\{L_{\bar{k}} \mathbf{u}_j, \mathbf{u}_k\} - \pi_*\{L_{\bar{k}} \mathbf{u}_j, L_{\bar{j}} \mathbf{u}_k\} \equiv 0,$$

which implies that

$$(4.14) \quad (u_j, u_k)_{j\bar{k}} = \pi_*(c_n\{L_j \mathbf{u}_j, L_k \mathbf{u}_k\}) - \pi_*(c_n\{[L_j, L_{\bar{k}}] \mathbf{u}_j, \mathbf{u}_k\}).$$

We shall use the following formula:

PROPOSITION 4.2.

$$[L_j, L_{\bar{k}}] = d^{\mathcal{L}} \delta_{[V_j, \bar{V}_k]} + \delta_{[V_j, \bar{V}_k]} d^{\mathcal{L}} + \langle V_j, V_k \rangle_{i\Theta(\mathcal{L}, h)}.$$

Proof. — By definition, locally we have

$$L_j = L_{V_j} - V_j(\phi), \quad L_{\bar{k}} = L_{\bar{V}_k}.$$

Thus

$$[L_j, L_{\bar{k}}] = [L_{V_j}, L_{\bar{V}_k}] - [V_j(\phi), L_{\bar{V}_k}] = L_{[V_j, \bar{V}_k]} + \bar{V}_k V_j(\phi).$$

By Cartan's formula, we have

$$L_{[V_j, \bar{V}_k]} = d^{\mathcal{L}} \delta_{[V_j, \bar{V}_k]} + \delta_{[V_j, \bar{V}_k]} d^{\mathcal{L}} + \delta_{[V_j, \bar{V}_k]} \partial \phi,$$

Thus

$$[L_j, L_{\bar{k}}] - \left(d^{\mathcal{L}} \delta_{[V_j, \bar{V}_k]} + \delta_{[V_j, \bar{V}_k]} d^{\mathcal{L}} \right) = \delta_{[V_j, \bar{V}_k]} \partial \phi + \bar{V}_k V_j(\phi).$$

By direct computation, we have

$$(4.15) \quad \delta_{[V_j, \bar{V}_k]} \partial \phi + \bar{V}_k V_j(\phi) = \langle V_j, V_k \rangle_{i\Theta(\mathcal{L}, h)}.$$

Thus the proposition follows. □

Now we can prove the following:

LEMMA 4.3. — *If $i\Theta(\mathcal{L}, h)|_{D_t} > 0, \forall t \in \mathbb{B}$, then*

$$(4.16) \quad \sum \pi_*(c_n\{[L_j, L_{\bar{k}}] \mathbf{u}_j, \mathbf{u}_k\}) = \|c\|_{i\Theta(\mathcal{L}, h)|_{D_t}}^2 + B + \sum (c_{j\bar{k}}(h) u_j, u_k),$$

where c is defined by (2.24) and B is the boundary term defined by

$$B := \sum \int_{\partial D_t} \theta_{j\bar{k}}(\rho) \langle u_j, u_k \rangle d\sigma$$

If $i\Theta(\mathcal{L}, h) \equiv 0$ on D then

$$(4.17) \quad \sum \pi_*(c_n\{[L_j, L_{\bar{k}}] \mathbf{u}_j, \mathbf{u}_k\}) = B.$$

Proof. — By the above proposition, we have

$$(4.18) \quad \sum \pi_*(c_n\{[L_j, L_{\bar{k}}] \mathbf{u}_j, \mathbf{u}_k\}) = \sum c_n \int_{\partial D_t} \{\delta_{[V_j, \bar{V}_k]} u_j, u_k\} + I,$$

where

$$(4.19) \quad I := \sum \langle \langle V_j, V_k \rangle_{i\Theta(\mathcal{L}, h)} u_j, u_k \rangle.$$

Now the boundary term can be written as

$$\sum c_n \int_{\partial D_t} \{\delta_{[V_j, \bar{V}_k]} u_j, u_k\} = \sum \int_{\partial D_t} (\delta_{[V_j, \bar{V}_k]} \partial \rho) \langle u_j, u_k \rangle d\sigma.$$

We shall prove $\delta_{[V_j, \bar{V}_k]} \partial \rho = \theta_{j\bar{k}}(\rho)$ on ∂D . In fact, by (4.15), we have

$$\delta_{[V_j, \bar{V}_k]} \partial \rho + \bar{V}_k V_j(\rho) = \langle V_j, V_k \rangle_{i\partial \bar{\partial} \rho},$$

and by our key lemma, $V_j|_{\partial D} = V_j^p$, thus $\bar{V}_k V_j(\rho) \equiv 0$ on ∂D and

$$\delta_{[V_j, \bar{V}_k]} \partial \rho \equiv \theta_{j\bar{k}}(\rho), \text{ on } \partial D.$$

Thus

$$(4.20) \quad B = \sum c_n \int_{\partial D_t} \{\delta_{[V_j, \bar{V}_k]} u_j, u_k\}.$$

and (4.18) implies (4.17). Now let us prove (4.16): By (2.13), we have

$$\langle V_j, V_k \rangle_{i\Theta(\mathcal{L}, h)} = c_{j\bar{k}}(h) + \langle V_j - V_j^h, V_k - V_k^h \rangle_{i\Theta(\mathcal{L}, h)|_{D_t}}.$$

Since

$$(V_j - V_j^h) \lrcorner (i\Theta(\mathcal{L}, h)|_{D_t}) = ((V_j - V_j^h) \lrcorner i\Theta(\mathcal{L}, h))|_{D_t} = (V_j \lrcorner i\Theta(\mathcal{L}, h))|_{D_t},$$

by (2.24), we have

$$(4.21) \quad I = \sum (c_{j\bar{k}}(h)u_j, u_k) + \|c\|_{i\Theta(\mathcal{L},h)|_{D_t}}^2.$$

Thus (4.16) follows. □

By (4.11) and (4.14), we have

$$(4.22) \quad (\Theta_{j\bar{k}}u_j, u_k) = \pi_*(c_n\{[L_j, L_{\bar{k}}]\mathbf{u}_j, \mathbf{u}_k\}) \\ + (D_{t^j}u_j, D_{t^k}u_k) - \pi_*(c_n\{L_j\mathbf{u}_j, L_k\mathbf{u}_k\}).$$

Let a^j be the $(n, 0)$ -part of $i_t^*(L_j\mathbf{u}_j)$ and b^j be the $(n-1, 1)$ -part of $i_t^*(L_j\mathbf{u}_j)$, i.e.

$$a^j = i_t^*(\partial_\phi\delta_{V_j} + \delta_{V_j}\partial_\phi)\mathbf{u}_j = i_t^*[\partial_\phi, \delta_{V_j}]\mathbf{u}_j,$$

and

$$b^j = i_t^*(\bar{\partial}\delta_{V_j} + \delta_{V_j}\bar{\partial})\mathbf{u}_j = (\bar{\partial}V_j)|_{D_t} \lrcorner u_j.$$

Then we have

$$(4.23) \quad \|\sum D_{t^j}u_j\|^2 - \sum \pi_*(c_n\{L_j\mathbf{u}_j, L_k\mathbf{u}_k\}) = -\|a\|^2 - \pi_*(c_n\{b, b\}),$$

where b is defined in (2.24) and

$$a = \sum ((D_{t^j}u_j)(t) - a^j).$$

We shall prove that:

PROPOSITION 4.4. — a is the L^2 -minimal solution of

$$(4.24) \quad \bar{\partial}^t(a) = \partial_\phi^t b + c,$$

where b and c are defined in (2.24).

Proof. — Since $i_t^*(\bar{\partial}\mathbf{u}_j) \equiv 0$ and $i_t^*(\partial_\phi\mathbf{u}_j) \equiv 0$, we have

$$(4.25) \quad \bar{\partial}^t a^j + \partial_\phi^t b^j = i_t^*(\bar{\partial}[\partial_\phi, \delta_{V_j}] + \partial_\phi[\bar{\partial}, \delta_{V_j}])\mathbf{u}_j \\ = i_t^*([\bar{\partial}, [\partial_\phi, \delta_{V_j}]] + [\partial_\phi, [\bar{\partial}, \delta_{V_j}]])\mathbf{u}_j.$$

Since

$$(4.26) \quad [\bar{\partial}, [\partial_\phi, \delta_{V_j}]] + [\partial_\phi, [\bar{\partial}, \delta_{V_j}]] + [\delta_{V_j}, [\bar{\partial}, \partial_\phi]] \equiv 0,$$

and $[\bar{\partial}, \partial_\phi] \equiv \Theta(\mathcal{L}, h)$, we get that

$$(4.27) \quad \bar{\partial}^t a^j + \partial_\phi^t b^j = -(V_j \lrcorner \Theta(\mathcal{L}, h))|_{D_t} \wedge u_j.$$

Recall that by (4.10), each $D_{t^j}u_j$ is just the Bergman projection to \mathcal{H}_t of a^j . Thus this proposition follows from (4.27). □

By Lemma 4.3, (4.22) and (4.23), we have
 (4.28)

$$\sum (\Theta_{j\bar{k}} u_j, u_k) = \sum \int_{\partial D_t} \theta_{j\bar{k}}(\rho) \langle u_j, u_k \rangle d\sigma + \sum (c_{j\bar{k}}(h) u_j, u_k) + R',$$

where

$$R' = \|c\|_{i\Theta(\mathcal{L}, h)|_{D_t}}^2 - \pi_*(c_n \{b, b\}) - \|a\|^2.$$

Now let us prove that $R' = R$. It is enough to prove that

$$-\pi_*(c_n \{b, b\}) = \|b\|_{\omega^t}^2.$$

But it follows directly from the following lemma:

LEMMA 4.5. — b is primitive with respect to $\omega^t := i\partial\bar{\partial}(-\log -\rho)|_{D_t}$.

Proof. — Recall that

$$b = \sum (\bar{\partial}V_j)|_{D_t} \lrcorner u_j.$$

Since b is an $(n - 1, 1)$ -form, by definition of primitivity, it suffices to show that

$$\omega^t \wedge b \equiv 0, \text{ on } D_t.$$

Thus it is enough to prove that

$$(4.29) \quad ((\bar{\partial}V_j) \lrcorner i\partial\bar{\partial}(-\log -\rho))|_{D_t} \equiv 0, \forall 1 \leq j \leq m.$$

By definition of V_j in our Key Lemma, $(V_j \lrcorner i\partial\bar{\partial}(-\log -\rho))|_{D_t} = 0$. Thus (4.29) is true. □

Remark. — Now we know that Theorem 2.11 is true if each u_j is a holomorphic section of \mathcal{H} . For finite rank vector bundles, the curvature operators are always pointwise defined, thus it is enough to find a curvature formula for holomorphic sections in finite rank case. One may guess that the same argument also works for the general infinite rank vector bundle. In the next subsection, we shall prove that at least the curvature operators for our bundle \mathcal{H} are pointwise defined. Thus we know that (4.28) is also true for general smooth sections of \mathcal{H} .

4.3. Curvature formula for general sections

By the above remark, we need to prove that the curvature operators $\Theta_{j\bar{k}}$ on \mathcal{H} are pointwise defined. We shall use the following two lemmas.

LEMMA 4.6. — $(\Theta_{j\bar{k}} u, v) = (u, \Theta_{k\bar{j}} v), \forall u, v \in \Gamma(\mathcal{H})$.

Proof. — By (2.6), we have

$$\begin{aligned} (u, v)_{\bar{k}j} &= \bar{\partial}_{t^k}((D_{t^j}u, v) + (u, \bar{\partial}_{t^j}v)) \\ &= (\bar{\partial}_{t^k}D_{t^j}u, v) + (D_{t^j}u, D_{t^k}v) + (\bar{\partial}_{t^k}u, \bar{\partial}_{t^j}v) + (u, D_{t^k}\bar{\partial}_{t^j}v). \end{aligned}$$

On the other hand,

$$\begin{aligned} (u, v)_{j\bar{k}} &= \partial_{t^j}((\bar{\partial}_{t^k}u, v) + (u, D_{t^k}v)) \\ &= (D_{t^j}\bar{\partial}_{t^k}u, v) + (D_{t^j}u, D_{t^k}v) + (\bar{\partial}_{t^k}u, \bar{\partial}_{t^j}v) + (u, \bar{\partial}_{t^j}D_{t^k}v). \end{aligned}$$

Since $(u, v)_{\bar{k}j} \equiv (u, v)_{j\bar{k}}$, the lemma follows by comparing the difference of the above two equality. □

LEMMA 4.7. — Assume that D satisfies (A1) and (A2). Fix $u \in \Gamma(\mathcal{H})$ and $t_0 \in \mathbb{B}$. Then $u|_{D_{t_0}}$ can be approximated by holomorphic sections of \mathcal{H} in the following sense:

For every $0 < s < 1$, there exists a holomorphic section $u^{(s)}$ of \mathcal{H} over an open neighborhood (may depend on s) of t_0 such that

$$(4.30) \quad (\eta, s) \mapsto u^{(s)}|_{D_{t_0}}(\eta), \quad 0 < s < 1, \quad (\eta, 0) \mapsto u|_{D_{t_0}}(\eta),$$

is smooth up to the boundary of $D_{t_0} \times [0, 1)$.

Proof. — Fix a sufficiently small $\varepsilon > 0$ and consider

$$D_{t_0}^s := \{\zeta \in \pi^{-1}(t_0) : \rho(t_0, \zeta) < \varepsilon s\}.$$

Let us define $u^{(s)}$ as the Bergman projection to the space of L^2 -holomorphic forms on $D_{t_0}^s$ of $u|_{D_{t_0}^s}$. By Siu’s theorem [46], for every $0 < s < 1$, $D_{t_0}^s$ has a Stein neighborhood in \mathcal{X} . Thus by Cartan’s theorem, every $u^{(s)}$ extends to a holomorphic section (also denoted by $u^{(s)}$) of \mathcal{H} over an open neighborhood of t_0 . The regularity properties of $\{u^{(s)}\}$ follows directly from Hamilton’s theorem (see Appendix A.2). □

Now let us finish the proof of Theorem 2.11. By the above two lemmas, for every $u_j \in \Gamma(\mathcal{H})$, $1 \leq j \leq m$, $t_0 \in \mathbb{B}$, we have

$$\begin{aligned} (\Theta_{j\bar{k}}u_j, u_k)(t_0) &= \lim_{s_1 \rightarrow 0} (\Theta_{j\bar{k}}u_j, u_k^{(s_1)})(t_0) = \lim_{s_1 \rightarrow 0} (u_j, \Theta_{k\bar{j}}u_k^{(s_1)})(t_0) \\ &= \lim_{s_1 \rightarrow 0} \lim_{s_2 \rightarrow 0} (u_j^{(s_2)}, \Theta_{k\bar{j}}u_k^{(s_1)})(t_0) \\ &= \lim_{s_1 \rightarrow 0} \lim_{s_2 \rightarrow 0} (\Theta_{j\bar{k}}u_j^{(s_2)}, u_k^{(s_1)})(t_0) \\ &= \lim_{s \rightarrow 0} (\Theta_{j\bar{k}}u_j^{(s)}, u_k^{(s)})(t_0). \end{aligned}$$

where $u_j^{(s)}$ are holomorphic sections of \mathcal{H} defined in Lemma 4.7. By our curvature formula for holomorphic sections, we know that $\sum(\Theta_{j\bar{k}}u_j^{(s)}, u_k^{(s)})$ can be written as

$$\sum \int_{\partial D_t} \theta_{j\bar{k}}(\rho)\langle u_j^{(s)}, u_k^{(s)}\rangle d\sigma + \sum(c_{j\bar{k}}(h)u_j^{(s)}, u_k^{(s)}) + R(s),$$

where

$$R(s) = \|c(s)\|_{i\Theta(\mathcal{L},h)|_{D_t}}^2 + \|b(s)\|_{\omega_t}^2 - \|a(s)\|_{\omega_t}^2.$$

Since $a(s)$, $b(s)$ and $c(s)$ only depend on $u_j^{(s)}|_{D_{t_0}}$, by Lemma 4.7, let $s \rightarrow 0$, we know that (2.27) is true at t_0 . Since t_0 is an arbitrary point in \mathbb{B} , the proof of Theorem 2.11 is complete.

4.4. Proof of Corollary 2.12

For any fixed $t_0 \in \mathbb{B}$, one may choose a sufficiently large positive constant A such that $\rho + A|t|^2$ is strictly plurisubharmonic in a neighborhood of the closure of $D \cap \pi^{-1}(U)$, where U is a small neighborhood of t_0 . Now for every $\varepsilon > 0$,

$$h^\varepsilon := h e^{-\varepsilon(\rho + A|t|^2)},$$

defines a smooth Hermitian metric on \mathcal{L} with positive curvature on a neighborhood of the closure of $D \cap \pi^{-1}(U)$. Denote by \mathcal{H}^ε the associated family of Hilbert spaces with respect to h^ε . Denote by $\Theta_{j\bar{k}}^\varepsilon$ the associated curvature operator on \mathcal{H}^ε . Since the total space D is Stein, we know that $\theta_{j\bar{k}}$ is semi-positive. By the construction of h^ε , we know that $c_{j\bar{k}}(h^\varepsilon)$ is positive on $D \cap \pi^{-1}(U)$. Thus our main theorem implies that \mathcal{H}^ε is Nakano positive on U . By Hamilton’s theorem, we have

$$\sum(\Theta_{j\bar{k}}u_j, u_k)(t_0) = \lim_{\varepsilon \rightarrow 0} \sum(\Theta_{j\bar{k}}^\varepsilon u_j, u_k)(t_0) \geq 0, \quad \forall u, v \in \Gamma(\mathcal{H}).$$

Thus \mathcal{H} is Nakano semi-positive at t_0 . Since t_0 is an arbitrary point in \mathbb{B} , we know that \mathcal{H} is Nakano semi-positive.

5. Curvature of the dual family

In this section, we shall prove our main application Corollary 2.15. As a direct application, we shall give a plurisubharmonicity property of the derivatives of the Bergman kernel, which can be seen as a generalization of Theorem C. In the last part of this section, based on a remarkable idea of Berndtsson and Lempert [14], we shall show how to use Corollary 2.15 to study plurisubharmonicity properties of the Bergman projection of currents with compact support.

5.1. Proof of Corollary 2.15

Let f be a holomorphic section of the dual of \mathcal{H} . By Definition 2.13, we know that there is a smooth section, say $P(f)$, of \mathcal{H} , such that

$$(5.1) \quad f^t(u^t) = (u^t, P(f)^t),$$

for every $u^t \in \mathcal{H}_t$. Moreover, by Definition 2.14, we know that

$$(5.2) \quad f(u) : t \mapsto f^t(u^t),$$

is a holomorphic function of t if u is a holomorphic section of \mathcal{H} . Thus we have

$$(5.3) \quad 0 \equiv \bar{\partial}_{tj} f(u) = (u, D_{tj} P(f)),$$

for every holomorphic section u of \mathcal{H} . By Lemma 4.7, we know that

$$(5.4) \quad D_{tj} P(f) \equiv 0,$$

which implies that

$$(5.5) \quad \partial_{tj} \bar{\partial}_{tk} (||P(f)||^2) = (\bar{\partial}_{tk} P(f), \bar{\partial}_{tj} P(f)) + (\Theta_{j\bar{k}} P(f), P(f)).$$

By Corollary 2.12, we have

$$\sum (\Theta_{j\bar{k}} (\xi_j P(f)), \xi_k P(f)) \geq 0.$$

for every $\xi \in \mathbb{C}^m$. Thus we have

$$\sum \bar{\partial}_{tk} \partial_{tj} (\log ||P(f)||^2) \xi_j \bar{\xi}_k \geq \frac{||\sum \bar{\xi}_k \bar{\partial}_{tk} P(f)||^2}{||P(f)||^2} - \frac{|(P(f), \sum \bar{\xi}_k \bar{\partial}_{tk} P(f))|^2}{||P(f)||^4},$$

on

$$(5.6) \quad U_f := \{t \in \mathbb{B} : ||P(f)^t|| > 0\}.$$

By Schwartz inequality, we have $\sum \bar{\partial}_{tk} \partial_{tj} (\log ||P(f)||^2) \xi_j \bar{\xi}_k \geq 0$ on U_f . Notice that

$$(5.7) \quad ||P(f)|| : t \mapsto ||P(f)^t|| = ||f^t||$$

is a smooth function on \mathbb{B} . Thus $\log ||P(f)|| = \log ||f||$ is plurisubharmonic on \mathbb{B} . The proof is complete.

5.2. Variation of the derivatives of the Bergman kernel

For simplicity purposes, we shall only consider the following case:

Pseudoconvex family in \mathbb{C}^n . — In this case, \mathcal{X} is $\mathbb{C}^n \times \mathbb{B}$ and π is just the natural projection to \mathbb{B} . Assume that D satisfies (A1) and (A2). One may look at $D = \{D_t\}_{t \in \mathbb{B}}$ as a smooth family of smoothly bounded strongly pseudoconvex domains in \mathbb{C}^n . Moreover, we shall assume that \mathcal{L} is a trivial line bundle over \mathcal{X} with Hermitian metric $h = e^{-\phi}$.

Variation formula of the derivatives of the Bergman kernel. Fix $\eta \in D_0$, replace \mathbb{B} by a smaller ball if necessary, one may assume that

$$(5.8) \quad \eta \in D_t, \quad \forall t \in \mathbb{B}.$$

Let us consider

$$D^\alpha \eta : f \mapsto f_\alpha(\eta) := \frac{\partial^{|\alpha|} f}{(\partial \mu^1)^{\alpha_1} \dots (\partial \mu^n)^{\alpha_n}}(\eta), \quad \forall f = f(\mu) d\mu \in \mathcal{H}_t,$$

where $\alpha \in \mathbb{N}^n$, $|\alpha| := \alpha_1 + \dots + \alpha_n$ and $d\mu$ is short for $d\mu^1 \wedge \dots \wedge d\mu^n$. By Definition 2.14, we know that every $D^\alpha \eta$ defines a holomorphic section of the dual of \mathcal{H} . Put

$$\cdot_\alpha \eta := P(D^\alpha \eta).$$

i.e., $\cdot_\alpha \eta$ is the unique smooth section of \mathcal{H} such that

$$(5.9) \quad (f, \cdot_\alpha \eta) = f_\alpha(\eta) \quad \forall f \in \mathcal{H}_t.$$

Let $K^t(\zeta, \eta) d\zeta \otimes \overline{d\eta}$ be the Bergman reproducing kernel of \mathcal{H}_t . Then (5.9) implies that

$$(5.10) \quad \cdot_0 \eta = K^t(\mu, \eta) d\mu, \quad (\cdot_\beta \eta, \cdot_\alpha \zeta) = (\cdot_\beta \eta)_\alpha(\zeta) = \overline{(\cdot_\alpha \zeta)_\beta(\eta)} = K^t_{\alpha\bar{\beta}}(\zeta, \eta),$$

where

$$K^t_{\alpha\bar{\beta}}(\zeta, \eta) = \frac{\partial^{|\alpha|+|\beta|} K^t}{(\partial \zeta^1)^{\alpha_1} \dots (\partial \zeta^n)^{\alpha_n} (\partial \bar{\eta}^1)^{\beta_1} \dots (\partial \bar{\eta}^n)^{\beta_n}}(\zeta, \eta).$$

By (5.4), we have

$$(5.11) \quad K^t_{j\bar{k}\alpha\bar{\beta}}(\zeta, \eta) = \partial_{t^j} \bar{\partial}_{t^k} (\cdot_\beta \eta, \cdot_\alpha \zeta) = ((\cdot_\beta \eta)_{\bar{k}}, (\cdot_\alpha \zeta)_{\bar{j}}) + (\Theta_{j\bar{k}}(\cdot_\beta \eta), \cdot_\alpha \zeta).$$

By (2.27), we have

$$(5.12) \quad (\Theta_{j\bar{k}}(\cdot_\beta \eta), \cdot_\alpha \zeta) = \int_{\partial D_t} \theta_{j\bar{k}}(\rho) \langle \cdot_\beta \eta, \cdot_\alpha \zeta \rangle d\sigma + (c_{j\bar{k}}(h) \cdot_\beta \eta, \cdot_\alpha \zeta) + R,$$

where

$$(5.13) \quad R = (c, c')_{i\partial\bar{\partial}\phi|_{D_t}} + (b, b')_{\omega^t} - (a, a')_{\omega^t},$$

if $i\partial\bar{\partial}\phi|_{D_t} > 0$, and

$$R = (b, b')_{\omega^t} - (a, a')_{\omega^t},$$

if $i\partial\bar{\partial}\phi \equiv 0$. Here

$$\omega^t = i\partial\bar{\partial}(-\log -\rho)|_{D_t},$$

and (a, b, c) (resp. (a', b', c')) are forms associated to $\cdot_\beta\eta$ (resp. $\cdot_\alpha\zeta$) respectively. Moreover,

$$\bar{\partial}^t a = \partial_\phi^t b + c, \quad \bar{\partial}^t a' = \partial_\phi^t b' + c'.$$

Remark. — Theorem A.5 implies that R is non-negative as a Hermitian form. Later we shall give an explicit expression of the Hörmander remaining term R in case $i\partial\bar{\partial}\phi \equiv 0$. (thus $c = c' \equiv 0$).

Hörmander remaining term for flat weight. Let us assume that $i\partial\bar{\partial}\phi \equiv 0$. By definition, then we have $c = c' \equiv 0$. Put

$$(5.14) \quad \square' = \partial_\phi^t (\partial_\phi^t)^* + (\partial_\phi^t)^* \partial_\phi^t,$$

We shall prove that:

LEMMA 5.1. — *If $i\partial\bar{\partial}\phi \equiv 0$ on D then*

$$(5.15) \quad R = (\mathbb{H}b, \mathbb{H}b')_{\omega^t},$$

where $\mathbb{H}b$ denotes the \square' -harmonic part of b .

Proof. — Since $i\partial\bar{\partial}\phi \equiv 0$ and ω^t is complete Kähler, we know that the $\bar{\partial}$ -Laplace \square'' is equal to \square' . Denote by G the associated Green operator. Let us omit ω^t in $(\cdot, \cdot)_{\omega^t}$, then we have

$$(a, a') = ((\bar{\partial}^t)^* G \partial_\phi^t b, a') = (G \partial_\phi^t b, \partial_\phi^t b').$$

Since b is primitive and $\bar{\partial}^t$ -closed, we know that b is $(\partial_\phi^t)^*$ -closed. Thus b can be written as

$$b = \mathbb{H}b + (\partial_\phi^t)^* f, \quad \partial_\phi^t f = 0.$$

Now

$$(a, a') = (G \partial_\phi^t (\partial_\phi^t)^* f, \partial_\phi^t b') = (f, \partial_\phi^t b') = (b - \mathbb{H}b, b') = (b, b') - (\mathbb{H}b, \mathbb{H}b').$$

Thus

$$R = (b, b') - (a, a') = (\mathbb{H}b, \mathbb{H}b'). \quad \square$$

Recall that

$$(5.16) \quad b = (\bar{\partial}V_j)|_{D_t} \lrcorner (\cdot_\beta\eta), \quad b' = (\bar{\partial}V_k)|_{D_t} \lrcorner (\cdot_\alpha\zeta).$$

Thus Lemma 5.1 implies that:

THEOREM 5.2 (Variation Formula of the Bergman Kernel). — *The first order variation formula of the Bergman kernel can be written as*

$$(5.17) \quad K_{j\alpha\bar{\beta}}^t(\zeta, \eta) = i^{n^2} \int_{D_t} \phi_j\{\cdot\beta\eta, \cdot\alpha\zeta\} - i^{n^2} \int_{\partial D_t} \delta_{V_j}\{\cdot\beta\eta, \cdot\alpha\zeta\},$$

Moreover, if $i\partial\bar{\partial}\phi \equiv 0$ on D then

$$(5.18) \quad K_{j\bar{k}\alpha\bar{\beta}}^t(\zeta, \eta) = ((\cdot\beta\eta)_{\bar{k}}, (\cdot\alpha\zeta)_{\bar{j}}) + \int_{\partial D_t} \theta_{j\bar{k}}(\rho)\langle\cdot\beta\eta, \cdot\alpha\zeta\rangle d\sigma + (\mathbb{H}b, \mathbb{H}b'),$$

where $b = (\bar{\partial}V_j)|_{D_t} \lrcorner (\cdot\beta\eta)$, $b' = (\bar{\partial}V_{\bar{k}})|_{D_t} \lrcorner (\cdot\alpha\zeta)$.

Proof. — Notice that (5.18) is a direct consequence of Lemma 5.1. Thus it suffices to prove (5.17). Notice that (4.3) implies that

$$K_{j\alpha\bar{\beta}}^t(\zeta, \eta) = i^{n^2} \int_{D_t} L_{V_j}^t\{\cdot\beta\eta, \cdot\alpha\zeta\}.$$

By Cartan’s formula

$$L_{V_j}^t = i_t^*(d\delta_{V_j} + \delta_{V_j}d),$$

thus we have

$$K_{j\alpha\bar{\beta}}^t(\zeta, \eta) = i^{n^2} \int_{\partial D_t} \delta_{V_j}\{\cdot\beta\eta, \cdot\alpha\zeta\} + i^{n^2} \int_{D_t} \frac{\partial}{\partial t^j}\{\cdot\beta\eta, \cdot\alpha\zeta\}.$$

By the reproducing formula,

$$i^{n^2} \int_{D_t} \frac{\partial}{\partial t^j}\{\cdot\beta\eta, \cdot\alpha\zeta\} = 2K_{j\alpha\bar{\beta}}^t(\zeta, \eta) - i^{n^2} \int_{D_t} \phi_j\{\cdot\beta\eta, \cdot\alpha\zeta\},$$

which implies (5.17). □

Remark. — If $\alpha = \beta = 0$ and $\phi \equiv 0$ then (5.17) is Komatsu’s formula (see [32]). Recently, Berndtsson [11] showed that (5.17) can be used to study the comparison principle for Bergman kernels. In fact, if D is a product then (5.17) is just [11, (2.2)].

5.3. Variation of the Bergman projection of currents

In the last section, we discussed the plurisubharmonicity properties of the Bergman projection of the derivatives of the Dirac measure. Recently, it is known that the plurisubharmonicity properties of the Bergman projection of other kind of currents are also very useful (see [14]). In this subsection, we shall show how to use Corollary 2.15 to study variation of the Bergman projection of general currents with compact support.

Smooth family of currents with compact support. Denote by A_t the space of smooth sections of $K_{D_t} + L_t$ over D_t . Put

$$\mathcal{A} = \{A_t\}_{t \in \mathbb{B}}.$$

We shall introduce the notion of the dual of \mathcal{A} by using the language of currents. Denote by A'_t the dual space of A_t , that is the space of L_t^* -valued degree $(0, n)$ -currents with compact support in D_t . Fix $f^t \in A'_t$, we shall formally write

$$f^t(u^t) = \int_{D_t} f^t \wedge u^t, \quad \forall u^t \in A_t,$$

even though the (n, n) -current $f^t \wedge u^t$ may not be integrable in general. Put

$$\mathcal{A}' = \{A'_t\}_{t \in \mathbb{B}}.$$

Denote by $\text{Supp } f^t$ the support of f^t . Denote by $K_{\mathcal{X}/\mathbb{B}}$ the relative canonical line bundle associated to π , recall that

$$(5.19) \quad K_{\mathcal{X}/\mathbb{B}} := K_{\mathcal{X}} - \pi^* K_{\mathbb{B}}, \quad K_{\mathcal{X}/\mathbb{B}}|_{D_t} \simeq K_{D_t}.$$

We shall introduce the following definition:

DEFINITION 5.3. — We call $f : t \rightarrow f^t \in A'_t$ a smooth family of currents with compact support if

$$(5.20) \quad \bigcup_{t \in K} \text{Supp } f^t \Subset D, \quad \forall K \Subset \mathbb{B},$$

and for every smooth section κ of $(K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}) \boxtimes (\overline{K_{\mathcal{X}/\mathbb{B}}} + \mathcal{L}^*)$ over

$$\mathcal{X} \times_{\pi} \mathcal{X} := \{(x, y) \in \mathcal{X} \times \mathcal{X} : \pi(x) = \pi(y)\},$$

there exists a smooth section, say $u_{f, \kappa}$, of $\overline{K_{\mathcal{X}/\mathbb{B}}} + \mathcal{L}^*$ over \mathcal{X} such that

$$(5.21) \quad f^t(\kappa^t(v^t)) = u_{f, \kappa}^t(v^t), \quad \forall v \in C^\infty(\mathcal{X}, K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}), \quad t \in \mathbb{B}.$$

Remark. — Let us explain the meaning of (5.21). The right hand side is clear, that is

$$u_{f, \kappa}^t(v^t) := \int_{D_t} u_{f, \kappa}^t \wedge v^t.$$

For the left hand side, by Assumption(A1), the restriction of π to the closure of D is proper, thus we know that

$$\kappa^t(v^t) : x \mapsto \int_{D_t} \kappa^t(x, \cdot) \wedge v^t(\cdot), \quad \forall x \in \pi^{-1}(t),$$

defines a section in A_t . Thus $f^t(\kappa^t(v^t))$ is well defined. Hence (5.21) means that the current defined by $f(\kappa)$ is smooth up to the boundary of D .

Bergman projection of smooth family of currents with compact support. We shall prove the following proposition:

PROPOSITION 5.4. — Assume that D satisfies (A1) and (A2). Let $f : t \rightarrow f^t \in A'_t$ be a smooth family of currents with compact support. Then

$$(5.22) \quad f^t : u^t \mapsto f^t(u^t), \quad \forall u^t \in \mathcal{H}_t,$$

defines a smooth section of the dual of \mathcal{H} in the sense of Definition 2.13.

Proof. — (5.20) implies that there exists a smooth real function, say χ , on D such that

$$\chi \equiv 1 \text{ on } \bigcup_{t \in \mathbb{B}} \text{Supp } f^t, \quad \text{Supp}(\chi|_{D_t}) \Subset D_t, \quad \forall t \in \mathbb{B}.$$

Denote by K^t the Bergman kernel of \mathcal{H}_t . Put

$$\chi K : (x, y) \mapsto \chi(x)K^{\pi(x)}(x, y), \quad \forall (x, y) \in D \times_{\pi} D.$$

By Hamilton’s theorem (see Appendix A.2), Assumptions (A1) and (A2) imply that χK is smooth up to the boundary, i.e., it extends to a smooth section of $(K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}) \boxtimes (\overline{K_{\mathcal{X}/\mathbb{B}}} + \mathcal{L}^*)$ over $\mathcal{X} \times_{\pi} \mathcal{X}$. By the reproducing property of K^t , we have

$$(\chi K)^t(v^t) = (\chi v)^t, \quad \forall v \in \Gamma(\mathcal{H}).$$

Thus by (5.21), we have

$$(5.23) \quad \begin{aligned} f^t(v^t) &= f^t((\chi v)^t) = f^t((\chi K)^t(v^t)) \\ &= u^t_{f, \chi K}(v^t) = \int_{D_t} u^t_{f, \chi K} \wedge v^t, \quad \forall v^t \in \mathcal{H}_t. \end{aligned}$$

Let us write

$$u^t_{f, \chi K} \wedge v^t = i^{n^2} \{v^t, P(f)^t\}, \quad \forall v^t \in C^\infty(D_t, K_{D_t} + L_t).$$

Thus we have

$$(5.24) \quad f^t((\chi K)^t(v^t)) = (v^t, P(f)^t), \quad \forall v^t \in C^\infty(D_t, K_{D_t} + L_t).$$

Since $(\chi K)^t(\mathcal{H}_t^\perp) = 0$, we have $P(f)^t \in \mathcal{H}_t$. Thus $P(f) \in \Gamma(\mathcal{H})$, and by (5.23), we have

$$(5.25) \quad f^t(v^t) = (v^t, P(f)^t), \quad \forall v \in \Gamma(\mathcal{H}).$$

By Definition 2.13, we know that f defines a smooth section of the dual of \mathcal{H} . □

Remark. — By Lemma 4.7, if D satisfies (A1) and (A2) then \mathcal{H}_t is equal to the closure of $\{u^t \in \mathcal{H}_t : u \in \Gamma(\mathcal{H})\}$. Thus (5.25) implies that

$$(5.26) \quad \|P(f)\|^2(t) = \sup\{|f^t(u)|^2 : u \in \mathcal{H}_t, i^{n^2} \int_{D_t} \{u, u\} = 1\}.$$

By this extremal property, one may generalize Corollary 2.15 to the case that the metric h on \mathcal{L} is singular.

6. Triviality and flatness

In this section, we shall prove Theorem 2.17 and use it to study triviality of holomorphic motions.

6.1. Proof of Theorem 2.17

Triviality implies flatness. By definition, if D is trivial then one may assume that the vector fields $\partial/\partial t^j$ are well defined on D , tangent to the boundary of D and can be extended to smooth vector fields on \mathcal{X} . Thus we have

$$(6.1) \quad \theta_{j\bar{k}}(\rho) \equiv 0,$$

on ∂D . Moreover, in this case $b \equiv 0$. If $\Theta(\mathcal{L}, h) \equiv 0$ then we also have $c \equiv 0$. Thus $a \equiv 0$ and $R \equiv 0$. By our main theorem, we know that \mathcal{H} is flat.

Flatness implies triviality. By Theorem 2.11 and our assumption, we have

$$\sum (\Theta_{j\bar{k}} u_j, u_k) = \sum \int_{\partial D_t} \theta_{j\bar{k}}(\rho) \langle u_j, u_k \rangle d\sigma + R,$$

where $R \geq 0$. Moreover, since D is Stein, we have

$$\sum \int_{\partial D_t} \theta_{j\bar{k}}(\rho) \langle u_j, u_k \rangle d\sigma \geq 0.$$

Thus if $\Theta_{j\bar{k}} \equiv 0$ then $R \equiv 0$ and

$$(6.2) \quad \theta_{j\bar{k}}(\rho) = \langle V_j^\rho, V_k^\rho \rangle_{i\partial\bar{\partial}\rho} \equiv 0 \text{ on } \partial D.$$

Since

$$(6.3) \quad \langle V_j, V_k \rangle_{i\partial\bar{\partial}(-\log -\rho)} = \frac{\langle V_j, V_k \rangle_{i\partial\bar{\partial}\rho}}{-\rho} + \frac{V_j(\rho)\overline{V_k(\rho)}}{\rho^2},$$

and $V_j = V_j^\rho$ on ∂D , by (6.3) and (6.2), we know that $\langle V_j, V_k \rangle_{i\partial\bar{\partial}(-\log-\rho)}$ is smooth up to the boundary of D . We shall use the following lemma, which follows from

$$(6.4) \quad V_j^\psi = \partial/\partial t^j - \sum \psi_{j\bar{\lambda}} \bar{\psi}^{\lambda\nu} \partial/\partial \mu^\nu,$$

by direct computation.

LEMMA 6.1. — *Let ψ be a smooth function on D . Assume that ψ is strictly plurisubharmonic on each fibre of D . Denote by V_j^ψ the horizontal lift of $\partial/\partial t^j$ with respect to $i\partial\bar{\partial}\psi$. Then*

$$[V_j^\psi, V_k^\psi] = 0, \quad [V_j^\psi, \overline{V_k^\psi}] = \sum c_{j\bar{k}}(\psi) \bar{\psi}^{\lambda\nu} \partial/\partial \mu^\nu - c_{j\bar{k}}(\psi) \psi^{\lambda\nu} \partial/\partial \bar{\mu}^\lambda,$$

where $c_{j\bar{k}}(\psi) := \langle V_j^\psi, V_k^\psi \rangle_{i\partial\bar{\partial}\psi}$.

Let us apply this lemma to $\psi = -\log-\rho$. Now by definition of V_j in Lemma 2.9, we have $V_j \equiv V_j^\psi$. Since $(-\log-\rho)^{\bar{\lambda}\nu} \equiv 0$ on ∂D . By (6.3) and the above lemma, we have

$$(6.5) \quad [V_j, V_k] = 0, \text{ on } D, \text{ and } [V_j, \overline{V_k}] = 0, \text{ on } \partial D.$$

on the boundary of D . Moreover, we shall prove that the following lemma is true.

LEMMA 6.2. — *Assume that D satisfies (A1) and (A2). Assume further that $K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}$ is trivial on each fibre of π and $\Theta(\mathcal{L}, h) \equiv 0$ on D . If $R \equiv 0$ then each V_j^ρ has a smooth extension, say \tilde{V}_j , that is holomorphic on fibres and smooth up to the boundary of D .*

If the above lemma is true then by (6.5), we have

$$[\tilde{V}_j, \tilde{V}_k] = [\tilde{V}_j, \overline{\tilde{V}_k}] = 0, \text{ on } \partial D.$$

Since \tilde{V}_j are holomorphic on fibres, we have

$$[\tilde{V}_j, \overline{\tilde{V}_k}] = \frac{\partial}{\partial t^j} \overline{\tilde{V}_k} - \frac{\partial}{\partial \bar{t}^k} \tilde{V}_j.$$

Thus

$$\frac{\partial}{\partial \bar{t}^k} \tilde{V}_j \equiv 0, \text{ on } \partial D.$$

Since $\frac{\partial}{\partial \bar{t}^k} \tilde{V}_j$ are holomorphic on each fibre, we have

$$\frac{\partial}{\partial \bar{t}^k} \tilde{V}_j \equiv 0, \text{ on } D.$$

Thus each \tilde{V}_j is a holomorphic vector field on D . Moreover,

$$[\tilde{V}_j, \tilde{V}_k] \equiv 0, \text{ on } \partial D.$$

Thus D is trivial. The proof of Theorem 2.17 is complete. □

Now let us prove Lemma 6.2:

Proof of Lemma 6.2. — We shall prove that $R \equiv 0$ implies that every $V_j|_{\partial D}(= V_j^\rho)$ has a holomorphic extension to D . Notice that the proof of Lemma 5.1 implies that

$$R = \|\mathbb{H}(\sum(\bar{\partial}V_j)|_{D_t} \lrcorner u_j)\|_{\omega^t}^2,$$

where $\omega^t = i\partial\bar{\partial}(-\log-\rho)|_{D_t}$. Thus $R \equiv 0$ implies that $\mathbb{H}b \equiv 0$. Since ω^t is d -bounded in the sense of Gromov (see [1, 21, 26]), and

$$b := \sum(\bar{\partial}V_j)|_{D_t} \lrcorner u_j$$

is $\bar{\partial}$ -closed, we know that there exists a smooth L_t -valued $(n-1, 0)$ -form u^t such that $\bar{\partial}^t u^t = b$ and

$$\|u^t\|_{\omega^t} \leq 2\|b\|_{\omega^t} < \infty.$$

We claim that $u : (\eta, t) \mapsto u^t(\eta)$ is smooth up to the boundary of D and $u = 0$ on ∂D . In fact, if we can show

$$(6.6) \quad \int_{D_t} \{f, b\} = 0,$$

for every ∂_ϕ^t -closed L_t -valued $(n-1, 1)$ -form f that is smooth up to the boundary of D_t , then $*b \in \text{Im}(\partial_\phi^t)^*$, where $*$ is the Hodge-de Rham operator with respect to $(i\partial\bar{\partial}\rho)|_{D_t}$ and $(\partial_\phi^t)^*$ is the adjoint of ∂_ϕ^t with respect to $(i\partial\bar{\partial}\rho)|_{D_t}$. By the regularity property of the $\bar{\partial}$ -Neumann problem (in fact, in our case, it is Dirichlet problem), one may solve

$$(\partial_\phi^t)^* v^t = *b.$$

where $v : (\eta, t) \mapsto v^t(\eta)$ is smooth up to the boundary of D . Since $(\partial_\phi^t)^* = -*\bar{\partial}^t*$, we have

$$\bar{\partial}^t(*v^t) = b.$$

Since $v^t \in \text{Dom}(\partial_\phi^t)^*$, we have $v^t = 0$ on ∂D_t . Thus $\|-*v^t\|_{\omega^t} < \infty$. Since there are no L^2 (with respect to ω^t) holomorphic L_t -valued $(n-1, 0)$ -forms on D_t , we have $u^t = -*v^t$. Thus our claim follows from (6.6).

Now let us prove (6.6). Put $\rho^t = \rho|_{D_t}$. Since b is smooth up to the boundary, we have

$$(6.7) \quad \int_{D_t} \{f, b\} = \lim_{r \rightarrow 0^-} \int_{\{\rho^t < r\}} \{f, b\} = \lim_{r \rightarrow 0^-} (-1)^n \int_{\{\rho^t = r\}} \{f, u^t\}.$$

Since

$$\omega^t \geq \frac{(i\partial\bar{\partial}\rho)|_{D_t}}{-\rho},$$

we know that $\|u^t\|_{\omega^t} < \infty$ implies that

$$\int_{D_t} \frac{|u^t|_{i\partial\bar{\partial}\rho|_{D_t}}^2}{-\rho} \frac{(i\partial\bar{\partial}\rho)^n|_{D_t}}{n!} < \infty.$$

Thus

$$\liminf_{r \rightarrow 0^-} \int_{\{\rho^t=r\}} |u^t|_{i\partial\bar{\partial}\rho|_{D_t}}^2 d\sigma = 0.$$

Since f is smooth up to the boundary, we know that

$$(6.8) \quad \lim_{r \rightarrow 0^-} (-1)^n \int_{\{\rho^t=r\}} f \wedge \bar{u}^t = 0.$$

Thus (6.6) follows from (6.7) and (6.8), and our claim is proved.

By our assumption, $K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}$ is trivial on $\pi^{-1}(t)$, thus there exists a holomorphic section, say e , of $K_{\mathcal{X}/\mathbb{B}} + \mathcal{L}$, that has no zero point in $\pi^{-1}(t)$. Now fix $t \in \mathbb{B}$, $1 \leq j \leq m$. Put

$$u_j = e, \quad u_k = 0, \quad \forall k \neq j.$$

By our claim, one may solve $\bar{\partial}^t u^t = (\bar{\partial}V_j)|_{D_t} \lrcorner e$ such that $u^t = 0$ on the boundary. Since e has no zero point in $\pi^{-1}(t)$, one may write

$$u^t = V \lrcorner e, \text{ on } D_t.$$

Thus we have

$$\bar{\partial}^t (V_j - V) = 0, \text{ on } D_t; \quad V_j - V = V_j \text{ on } \partial D_t.$$

Thus $V_j|_{\partial D_t}$ has a holomorphic extension, say $\tilde{V}_j|_{D_t}$, to D_t . The regularity property of \tilde{V}_j follows from the regularity property of u^t . The proof is complete. □

6.2. Triviality of holomorphic motions

We shall show how to use Theorem 2.17 to study triviality of holomorphic motions.

Basic notions on holomorphic motion. Recall that that if every fibre D_t is a domain in \mathbb{C} then:

∂D is Levi-flat if and only if $\theta_{j\bar{k}}(\rho) \equiv 0$.

It is known that the boundary of the total space of a holomorphic motion of a planar domain is Levi-flat. Recall that, by definition, a homeomorphism

$$(6.9) \quad F : (z, t) \mapsto (f(z, t), t),$$

from $D_0 \times \mathbb{B}$ to D is called a *holomorphic motion* (see [36]) of D_0 (with total space D) if $f(\cdot, 0)$ is the identity mapping and $f(z, \cdot)$ is holomorphic for every fixed $z \in D_0$.

Curvature formula for holomorphic motions. Assume that D_0 is a smooth domain in \mathbb{C} and F is smooth up to the boundary. Assume further that \mathcal{L} is trivial and $\phi \equiv 0$ on D . Put

$$V_j^F := F_*(\partial/\partial t^j) = \partial/\partial t^j + f_j(z, t)\partial/\partial \zeta.$$

By definition, $V_j^F(\rho) \equiv 0$ on ∂D . Thus

$$V_j^F = V_j = V_j^\rho, \text{ on } \partial D.$$

Hence we have

$$\| \sum (V_j^F - V_j) \lrcorner u_j \|_{\omega^t} < \infty,$$

which implies that

$$\sum (\Theta_{j\bar{k}} u_j, u_k) = \| \mathbb{H}(\sum (\bar{\partial} V_j)|_{D_t} \lrcorner u_j) \|^2 = \| \mathbb{H}(\sum (\bar{\partial} V_j^F)|_{D_t} \lrcorner u_j) \|^2.$$

Criterion for $\Theta_{j\bar{k}} \equiv 0$ by using the Bergman kernel. Put

$$J = f_{\bar{z}}/f_z.$$

Since

$$\partial/\partial \bar{\zeta} = z_{\bar{\zeta}}\partial/\partial z + \bar{z}_{\bar{\zeta}}\partial/\partial \bar{z},$$

and

$$\bar{z}_{\bar{\zeta}} = \frac{f_z}{|f_z|^2 - |f_{\bar{z}}|^2}, \quad z_{\bar{\zeta}} = \frac{-f_{\bar{z}}}{|f_z|^2 - |f_{\bar{z}}|^2},$$

we have

$$(6.10) \quad (\bar{\partial} V_j^F)|_{D_t} = (f_{jz}z_{\bar{\zeta}} + f_{j\bar{z}}\bar{z}_{\bar{\zeta}})d\bar{\zeta} \otimes \frac{\partial}{\partial \zeta} = \frac{(f_z)^2 J_j}{|f_z|^2(1 - |J|^2)} d\bar{\zeta} \otimes \frac{\partial}{\partial \zeta}.$$

Thus $\Theta_{j\bar{k}} \equiv 0$ is equivalent to

$$(6.11) \quad \int_{D_t} K^t(\zeta, \bar{\eta}) \left(\frac{(f_z)^2 J_j}{|f_z|^2(1 - |J|^2)} \right) (z(\zeta, t), t) \text{ id}\zeta \wedge d\bar{\zeta} = 0.$$

for every (η, t) in D and every j .

Proof of Corollary 2.18. — Since $J = a(t)$ now, by (6.11), we know that $\Theta_{j\bar{k}} \equiv 0$ is equivalent to

$$\frac{a_j}{1 - |a|^2} \int_{D_t} K^t(\zeta, \bar{\eta}) \, id\zeta \wedge d\bar{\zeta} = 0.$$

for every (η, t) in D and every j . But notice that

$$\int_{D_t} K^t(\zeta, \bar{\eta}) \, id\zeta \wedge d\bar{\zeta} \equiv 1.$$

Thus $\Theta_{j\bar{k}} \equiv 0$ is equivalent to $a_j \equiv 0$ for every j . Since $a(0) = 0$, we know that $\Theta_{j\bar{k}} \equiv 0$ is equivalent to $a \equiv 0$. □

Remark. — In [35], Ren Shan Liu showed that if $f = z + t^2\bar{z}$, then $F(\mathbb{D} \times \mathbb{D})$ is not biholomorphic equivalent to the bidisc, where \mathbb{D} denotes the unit disc. Interested readers can find more information on the holomorphic motion in [49] and [50].

Appendix

A.1. Variation of fibre integrals

Let \mathbb{B} be the unit ball in \mathbb{R}^m . Let $\{D_t\}_{t \in \mathbb{B}}$ be a family of smoothly bounded domains in \mathbb{R}^n . Put

$$D := \{(t, x) \in \mathbb{R}^{m+n} : x \in D_t, t \in \mathbb{B}\}.$$

Assume that there is a real valued function ρ on $\mathbb{B} \times \mathbb{R}^n$ such that for each t in \mathbb{B} , $\rho|_{D_t}$ is a smooth defining function of D_t .

We call $\{D_t\}_{t \in \mathbb{B}}$ a smooth family if there exists a fibre preserving diffeomorphism Φ from $\mathbb{B} \times D_0$ onto D such that for each $1 \leq j \leq m$, $\Phi_*(\partial/\partial t^j)$ extends to a smooth vector field on \mathbb{R}^n . Put

$$(A.1) \quad [D] := \bar{D} \cap (\mathbb{B} \times \mathbb{R}^n), \quad \delta D := \partial D \cap (\mathbb{B} \times \mathbb{R}^n).$$

Let $dx := dx^1 \wedge \dots \wedge dx^n$ be the Euclidean volume form on \mathbb{R}^n . Fix a smooth function f on a neighborhood of $[D]$. If $\{D_t\}_{t \in \mathbb{B}}$ is a smooth family then the fibre integrals

$$F(t) := \int_{D_t} f(t, x) dx$$

depend smoothly on $t \in \mathbb{B}$. We shall introduce a natural way to compute the derivatives of $F(t)$ (see [44] for related results). For every fixed $j \in \{1, \dots, m\}$, let

$$V_j := \frac{\partial}{\partial t^j} - \sum v_j^\lambda \frac{\partial}{\partial x^\lambda}$$

be a smooth vector field on a neighborhood of $[D]$. We shall prove that:

THEOREM A.3. — *Let $\{D_t\}_{t \in \mathbb{B}}$ be a smooth family of smoothly bounded domain in \mathbb{R}^m . Assume that $V_j(\rho) = 0$ on δD . Then we have*

$$(A.2) \quad \frac{\partial F}{\partial t^j}(t) = \int_{D_t} L_{V_j}^t(f(t, x)dx) = \int_{D_t} L_{V_j}(f(t, x)dx),$$

for every t in \mathbb{B} , where $L_{V_j}^t := i_t^*(L_{V_j})$.

Proof. — Without loss of generality, one may assume that $t = 0$ and $j = 1$. Since $V_1(\rho)$ vanishes on δD , the motion

$$\Phi : (-1, 1) \times D_0 \rightarrow \mathbb{R}^m$$

of D_0 associated to V_1 is compatible with $\{D_t\}$, i.e.

$$\Phi(a \times D_0) = D_{a\nu}, \quad \nu = (1, 0, \dots, 0) \in \mathbb{R}^m,$$

for every $a \in (-1, 1)$. Since for every fixed $a \in (-1, 1)$,

$$\Phi^a : x \mapsto \Phi(a, x)$$

is a C^∞ isomorphism from D_0 to $D_{a\nu}$, we have

$$(A.3) \quad \frac{\partial F}{\partial t^1}(0) = \lim_{0 \neq a \rightarrow 0} \int_{D_0} \frac{f(a\nu, \Phi^a(x))d\Phi^a(x) - f(0, x)dx}{a}$$

Since V_1 and f are smooth up to the boundary, we have

$$(A.4) \quad \frac{\partial F}{\partial t^1}(0) = \int_{D_0} \lim_{0 \neq a \rightarrow 0} \frac{f(a\nu, \Phi^a(x))d\Phi^a(x) - f(0, x)dx}{a}.$$

By definition of Lie derivative,

$$(A.5) \quad L_{V_1}(f(t, x)dx)(0, x) = \lim_{0 \neq a \rightarrow 0} \frac{[(\Psi^a)^*(f dx)](0, x) - f(0, x)dx}{a},$$

where

$$\Psi^a : (b\nu, \Phi^b(x)) \mapsto (b\nu + a\nu, \Phi^{b+a}(x)), \quad (b, x) \in (-1 + |a|, 1 - |a|) \times D_0.$$

Since

$$i_0^* \{[(\Psi^a)^*(f dx)](0, x) - f(a\nu, \Phi^a(x))d\Phi^a(x)\} = 0,$$

(A.2) follows from (A.4) and (A.5). □

Now assume that $m = 2$, put

$$\frac{\partial}{\partial t} := \frac{1}{2} \left(\frac{\partial}{\partial t^1} - i \frac{\partial}{\partial t^2} \right), \quad \frac{\partial}{\partial \bar{t}} := \frac{1}{2} \left(\frac{\partial}{\partial t^1} + i \frac{\partial}{\partial t^2} \right).$$

Let

$$V = \frac{\partial}{\partial t} - \sum v^\lambda \frac{\partial}{\partial x^\lambda}$$

be a smooth vector field on a neighborhood of $[D]$. If $V(\rho)$ vanishes on δD , then both $2 \operatorname{Re} V$ and $-2 \operatorname{Im} V$ satisfy the assumption of Theorem A.3. Thus we have:

COROLLARY A.4. — *If $V(\rho)$ vanishes on δD then*

$$\frac{\partial F}{\partial t}(t) = \int_{D_t} L_V^t(f(t, x)dx), \quad \frac{\partial F}{\partial \bar{t}}(t) = \int_{D_t} L_{\widetilde{V}}^t(f(t, x)\widetilde{dx}),$$

for every $t \in \mathbb{B}$.

A.2. Stability of the Bergman kernel

We shall give a short account of Hamilton’s theory on regularity properties of families of non-coercive boundary value problems. By [3, Lemma 2.1], stability of Bergman kernels follows directly from stability of solutions u^t of a family of $\bar{\partial}$ -Neumann problems $\square^t(\cdot) = f^t$. But it is not easy to prove regularity of u^t by the standard method. In fact, if we want to use

$$(A.6) \quad \|\square^t(u^t - u^s)\| = \|f^t - f^s - (\square^t - \square^s)u^s\|,$$

to estimate $\|u^t - u^s\|$ then we have to find a natural connection between the domain of \square^t and the domain of \square^s (i.e., u^s may not be in the domain of \square^t).

Hamilton [28] found a more natural way to study the regularity properties of families of non-coercive boundary value problems (not only for the $\bar{\partial}$ -Neumann problem). For reader’s convenience we give a sketch description of Hamilton’s idea.

Instead of considering \square^t (whose domain satisfies the so called $\bar{\partial}$ -Neumann condition), Hamilton considered the full Laplace operator $\widetilde{\square}^t$ (whose domain contains all forms smooth up to the boundary). Let u^t be a form smooth up to the boundary. In general, the Sobolev norm of $\widetilde{\square}^t(u^t)$ could not control the Sobolev norm of u^t . In fact, u^t has to be in the domain of \square^t (see [22]). Thus two more operators (sending forms on \bar{D}_t to forms on the boundary of D_t) are used in Hamilton’s paper, i.e., he considered the full $\bar{\partial}$ -Neumann problem

$$(A.7) \quad \mathfrak{S}^t(\cdot) := \left(\widetilde{\square}^t, (\bar{\partial}^t \rho) \vee, (\bar{\partial}^t \rho) \vee \bar{\partial}^t \right) (\cdot) = f^t,$$

where

$$(\bar{\partial}^t \rho) \vee := (\bar{\partial}^t \rho \wedge \cdot)^*.$$

Now the domain of \mathfrak{S}^t is $C_{\bullet,\bullet}^\infty(\overline{D}_t)$ for each t . Choose a C^∞ trivialization mapping

$$\mathbb{B} \times D_0 \simeq D,$$

then the domain of \mathfrak{S}^t can be seen as a fixed space $C_{\bullet,\bullet}^\infty(\overline{D}_0)$. Moreover, by [28], the constant in the basic estimates for \mathfrak{S}^t can be chosen to be independent of $t \in \mathbb{B}$. Thus (A.6) applies. The interested reader is referred to that paper for further information and a clear proof.

A.3. L^2 -estimate for $\overline{\partial}a = \partial_\phi b + c$

We shall prove a generalization of Demailly’s theorem (see [6, 19, 29]) in this section.

THEOREM A.5. — *Let (L, h) be a Hermitian line bundle over an n -dimensional complete Kähler manifold (X, ω) . Let v be a smooth $\overline{\partial}$ -closed L -valued $(n, 1)$ -form. Assume that*

$$i\Theta(L, h) > 0 \text{ on } X, \text{ (resp. } i\Theta(L, h) \equiv 0 \text{ on } X)$$

and

$$I(v) := \inf_{v=\partial_\phi b+c} \|b\|_\omega^2 + \|c\|_{i\Theta(L,h)}^2 < \infty, \text{ (resp. } I(v) := \inf_{v=\partial_\phi b} \|b\|_\omega^2 < \infty).$$

Then there exists a smooth L -valued $(n, 0)$ -form a on X such that $\overline{\partial}a = v$ and

$$(A.8) \quad \|a\|_\omega^2 \leq I(v).$$

Proof. — We shall only prove the $i\Theta(L, h) > 0$ case, since the $i\Theta(L, h) \equiv 0$ case can be proved by a similar argument. By Hörmander’s theorem and the standard density lemma for complete Kähler manifold, it suffices to prove that,

$$(A.9) \quad |(\partial_\phi b + c, g)_\omega|^2 \leq (\|b\|_\omega^2 + \|c\|_{i\Theta(L,h)}^2)(\|\overline{\partial}^* g\|_\omega^2 + \|\overline{\partial} g\|_\omega^2),$$

for every smooth L -valued $(n, 1)$ -form g with compact support in X . Notice that

$$(\partial_\phi b + c, g)_\omega = (b, \partial_\phi^* g)_\omega + (c, g)_\omega.$$

Hence

$$|(\partial_\phi b + c, g)_\omega|^2 \leq (\|b\|_\omega^2 + \|c\|_{i\Theta(L,h)}^2)(\|\partial_\phi^* g\|_\omega^2 + ([i\Theta(L, h), \Lambda_\omega]g, g)_\omega),$$

where Λ_ω denotes the adjoint of $\omega \wedge$. Thus (A.9) follows from the Bochner–Kodaira–Nakano formula. The proof is complete. \square

BIBLIOGRAPHY

- [1] B. BERNDTSSON, “The extension theorem of Ohsawa-Takegoshi and the theorem of Donnelly-Fefferman”, *Ann. Inst. Fourier* **46** (1996), no. 4, p. 1083-1094.
- [2] ———, “Prekopa’s theorem and Kiselman’s minimum principle for plurisubharmonic functions”, *Math. Ann.* **312** (1998), no. 4, p. 785-792.
- [3] ———, “Subharmonicity properties of the Bergman kernel and some other functions associated to pseudoconvex domains”, *Ann. Inst. Fourier* **56** (2006), no. 6, p. 1633-1662.
- [4] ———, “Curvature of vector bundles associated to holomorphic fibrations”, *Ann. Math.* **169** (2009), no. 2, p. 531-560.
- [5] ———, “Positivity of direct image bundles and convexity on the space of Kähler metrics”, *J. Diff. Geom.* **81** (2009), no. 3, p. 457-482.
- [6] ———, “An introduction to things $\bar{\partial}$ ”, in *Analytic and algebraic geometry*, IAS/Park City Math., vol. 17, Providence, RI, 2010, p. 7-76.
- [7] ———, “Strict and nonstrict positivity of direct image bundles”, *Math. Z.* **269** (2011), no. 3-4, p. 1201-1218.
- [8] ———, “Convexity on the space of Kähler metrics”, *Ann. Fac. Sci. Toulouse* **22** (2013), no. 4, p. 713-746.
- [9] ———, “The openness conjecture for plurisubharmonic functions”, <http://arxiv.org/abs/1305.5781>, 2013.
- [10] ———, “A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry”, *Invent. Math.* **200** (2014), no. 1, p. 149-200.
- [11] ———, “A comparison principle for Bergman kernels”, <http://arxiv.org/abs/1501.02440>, 2015.
- [12] ———, “The Openness Conjecture and Complex Brunn-Minkowski Inequalities”, in *Complex Geometry and Dynamics*, Abel Symposia, vol. 10, Springer, 2015, p. 29-44.
- [13] B. BERNDTSSON & R. J. BERMAN, “Convexity of the K-energy on the space of Kähler metrics and uniqueness of extremal metrics”, <http://arxiv.org/abs/1405.0401>, 2015.
- [14] B. BERNDTSSON & L. LEMPERT, “A proof of the Ohsawa-Takegoshi theorem with sharp estimates”, <http://arxiv.org/abs/1407.4946>, 2014.
- [15] B. BERNDTSSON & M. PAUN, “Bergman kernels and the pseudoeffectivity of relative canonical bundles”, *Duke Math. J.* **145** (2008), no. 2, p. 341-378.
- [16] H. J. BRASCAMP & E. H. LIEB, “On extensions of the Brunn-Minkowski and Prekopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation”, *J. Funct. Anal.* **22** (1976), p. 366-389.
- [17] Y. J. CHOI, “A study of variations of pseudoconvex domains via Kähler-Einstein metrics”, *Math. Z.* **281** (2015), no. 1-2, p. 299-314.
- [18] J.-P. DEMAILLY, “Complex analytic and differential geometry”, Book available from the author’s homepage <https://www-fourier.ujf-grenoble.fr/~demailly/documents.html>.
- [19] ———, “Estimations L^2 pour l’opérateur $\bar{\partial}$ d’un fibré vectoriel holomorphe semi-positif au-dessus d’une variété kählérienne complète”, *Ann. Sci. Éc. Norm. Sup.* **15** (1982), p. 457-511.
- [20] S. K. DONALDSON, “Symmetric spaces, Kähler geometry and Hamiltonian dynamics”, in *Northern California Symplectic Geometry Seminar*, Amer. Math. Soc. Transl. Ser. 2, vol. 196, Amer. Math. Soc., 1999, p. 13-33.

- [21] H. DONNELLY & C. FEFFERMAN, “ L^2 -cohomology and index theorem for the Bergman metric”, *Ann. Math.* **118** (1983), p. 593-618.
- [22] G. B. FOLLAND & J. J. KOHN, *The Neumann problem for the Cauchy-Riemann complex*, Ann. Math. Studies, vol. 75, Princeton University Press, 1972, viii+146 pages.
- [23] T. GEIGER & G. SCHUMACHER, “Curvature of higher direct image sheaves”, <http://arxiv.org/abs/1501.07070>, 2015.
- [24] R. E. GREENE & S. G. KRANTZ, “Deformation of complex structures, estimates for the $\bar{\partial}$ -equation, and stability of the Bergman kernel”, *Adv. Math.* **43** (1982), p. 1-86.
- [25] P. GRIFFITHS, *Topics in transcendental algebraic geometry*, Annals of Mathematics Studies, vol. 106, Princeton University Press, 1984, viii+316 pages.
- [26] M. GROMOV, “Kähler hyperbolicity and L_2 -Hodge theory”, *J. Diff. Geom.* **33** (1991), no. 1, p. 263-292.
- [27] R. S. HAMILTON, “Deformation of complex structures on manifolds with boundary. I. The stable case”, *J. Diff. Geom.* **12** (1977), p. 1-45.
- [28] ———, “Deformation of complex structures on manifolds with boundary. II. Families of noncoercive boundary value problems”, *J. Diff. Geom.* **14** (1979), p. 409-473.
- [29] L. HÖRMANDER, *An introduction to complex analysis in several variables*, 2nd ed., North-Holland Mathematical Library, vol. 7, Elsevier, 1973, 213 pages.
- [30] K. KODAIRA, *Complex manifolds and deformation of complex structures*, Grundlehren der Mathematischen Wissenschaften, vol. 283, Springer-Verlag, 1986, x+465 pages.
- [31] K. KODAIRA & D. C. SPENCER, “On deformations of complex analytic structures, III. Stability theorems for complex structures”, *Ann. Math.* **71** (1960), p. 43-76.
- [32] G. KOMATSU, “Hadamard’s variational formula for the Bergman kernel”, *Proc. Japan Acad.* **58** (1982), p. 345-348.
- [33] L. LEMPERT & R. SZŐKE, “Direct Images, Fields of Hilbert Spaces, and Geometric Quantization”, *Commun. Math. Phys.* **327** (2014), no. 1, p. 49-99.
- [34] K. LIU & X. YANG, “Curvatures of direct image sheaves of vector bundles and applications”, *J. Diff. Geom.* **98** (2014), no. 1, p. 117-145.
- [35] R. S. LIU, “A property of the graph of a holomorphic motion”, *J. Fudan Univ.* **47** (2008), no. 2, p. 172-176, in Chinese.
- [36] R. MÀÑÉ, P. SAD & D. SULLIVAN, “On the dynamics of rational maps”, *Ann. Éc. Norm. Supér.* **16** (1983), p. 193-217.
- [37] T. MABUCHI, “K-energy maps integrating Futaki invariants”, *Tohoku Math. J.* **38** (1986), no. 1-2, p. 575-593.
- [38] F. MAITANI, “Variations of meromorphic differentials under quasiconformal deformations”, *J. Math. Kyoto Univ.* **24** (1984), p. 49-66.
- [39] F. MAITANI & H. YAMAGUCHI, “Variation of Bergman metrics on Riemann surfaces”, *Math. Ann.* **330** (2004), no. 3, p. 477-489.
- [40] C. MOURougane & S. TAKAYAMA, “Hodge metrics and positivity of direct images”, *J. Reine Angew. Math.* **606** (2007), p. 167-178.
- [41] ———, “Hodge metrics and the curvature of higher direct images”, *Ann. Éc. Norm. Supér.* **41** (2008), no. 6, p. 905-924.
- [42] T. OHSAWA & K. TAKEGOSHI, “On the extension of L^2 holomorphic functions”, *Math. Z.* **195** (1987), p. 197-204.
- [43] A. PREKOPA, “On logarithmic concave measures and functions”, *Acad. Sci. Math.* **34** (1973), p. 335-343.
- [44] G. SCHUMACHER, “Positivity of relative canonical bundles and applications”, *Invent. Math.* **190** (2012), no. 1, p. 1-56.

- [45] S. SEMMES, “Interpolation of Banach spaces, differential geometry and differential equations”, *Rev. Mat. Iberoam.* **4** (1988), no. 1, p. 155-176.
- [46] Y. T. SIU, “Every Stein subvariety admits a Stein neighborhood”, *Invent. Math.* **38** (1976), p. 89-100.
- [47] ———, “Curvature of the Weil-Petersson metric in the moduli space of compact Kähler-Einstein manifolds of negative first Chern class”, in *Contributions to Several Complex Variables, Proc. Conf. Complex Analysis, Notre Dame/Indiana, 1984*, Aspects Math., vol. E9, Vieweg, 1986, p. 261-298.
- [48] ———, “Extension of twisted pluricanonical sections with plurisubharmonic weight and invariance of semipositively twisted plurigenera for manifolds not necessarily of general type”, in *Complex Geometry*, Springer, Berlin, 2002, p. 223-277.
- [49] Z. SŁODKOWSKI, “Holomorphic motions and polynomial hulls”, *Proc. Amer. Math. Soc.* **111** (1991), no. 2, p. 347-355.
- [50] D.-P. SULLIVAN & W.-P. THURSTON, “Extending holomorphic motions”, *Acta Math.* **157** (1986), p. 243-286.
- [51] D. V. TRAN, “Direct images as Hilbert fields and their curvatures”, PhD Thesis, Purdue University, USA, 2014.
- [52] H. TSUJII, “Variation of Bergman kernels of adjoint line bundles”, <http://arxiv.org/abs/math/0511342>, 2005.
- [53] X. WANG, “Variation of the Bergman kernels under deformations of complex structures”, <http://arxiv.org/abs/1307.5660>, 2013.

Manuscrit reçu le 6 octobre 2015,
révisé le 20 décembre 2015,
accepté le 14 juin 2016.

Xu WANG
Fudan University
School of Mathematical Sciences
Shanghai, 200433 (China)
wangxu1113@gmail.com
Current address:
Chalmers University of Technology and University
of Gothenburg
Department of Mathematical Sciences
SE-412 96 Gothenburg (Sweden)
xuwa@chalmers.se