

ANNALES DE L'I. H. P., SECTION A

VIVIANE BALADI

STEFANO ISOLA

BERNARD SCHMITT

Transfer operator for piecewise affine approximations of interval maps

Annales de l'I. H. P., section A, tome 62, n° 3 (1995), p. 251-265

http://www.numdam.org/item?id=AIHPA_1995__62_3_251_0

© Gauthier-Villars, 1995, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Transfer operator for piecewise affine approximations of interval maps

by

Viviane BALADI

ETH Zürich, CH-8092 Zürich, Switzerland
(On leave from CNRS, UMR 128, ENS Lyon, France)

Stefano ISOLA

Dipartimento di Matematica, Università degli Studi di Bologna,
I-40127 Bologna, Italy.

and

Bernard SCHMITT

Université de Bourgogne, Laboratoire de Topologie, URA 755,
21004 Dijon, France.

ABSTRACT. – We consider a natural approximation scheme for piecewise expanding, piecewise $C^{1+\text{Lipschitz}}$, mixing Markov interval maps f by piecewise affine maps. We prove that the densities of the absolutely continuous invariant probability measures of the approximations converge exponentially fast to the density of the absolutely continuous invariant probability measure of f in the uniform norm. To do this we compare the relevant transfer operators of the approximations with that of f , and use recently developed perturbation techniques.

RÉSUMÉ. – Nous considérons un algorithme naturel d'approximation par des transformations affines par morceaux d'applications de l'intervalle f dilatantes et $C^{1+\text{Lipschitz}}$ par morceaux, mélangeantes et markoviennes. Nous montrons que les densités des mesures de probabilité invariantes absolument continues des approximations convergent uniformément vers la densité de la mesure de probabilité invariante absolument continue de f , avec une vitesse exponentielle. Nous utilisons pour cela des techniques perturbatives récemment introduites qui nous permettent de comparer les opérateurs de transfert des approximations avec celui de f .

1. INTRODUCTION

A transformation f of $[0, 1]$ is called Markov if there exist disjoint open intervals I_1, \dots, I_l , the union of whose closures is $[0, 1]$, such that f restricted to each I_j is monotone and continuous, and such that the closure of each $f(I_j)$ is the closure of a union of intervals I_k . When this property holds, one may study the dynamical system generated by the iteration of f using symbolic dynamics and transfer operators which are the same as those in equilibrium statistical mechanics [13]. However, the statistical properties of the dynamical orbits of f can only be fully described in terms of Markov chains if the restriction of the Markov map f to each interval I_j is affine. Indeed, in this case, the associated transfer operator (see below) has a finite matrix representation. Thus, given a (non-linear) Markov transformation f , which we will assume to be topologically mixing, we are faced with the problem of approximating it with a sequence of piecewise affine Markov maps.

We show that the above problem can be solved constructively. More precisely, we obtain a sequence of finite Markov stochastic matrices whose normalised eigenvectors to the eigenvalue one approach the stationary probability density of f exponentially fast in the uniform norm. Our main technical tool is a non-standard perturbative argument, first used in [3] to deal with stochastic perturbations. The problem of finding the invariant probability measure by discretization of the transfer operator was first raised by Ulam [17] and has been studied by many authors. In particular, Gora and Boyarski [6] considered more general approximation schemes than ours, and did not need any Markov assumption. However, they only obtained the L^1 convergence of the invariant densities, and no estimate on the speed of this convergence (we believe that the methods from [3] used in the present article would yield another proof of the L^1 convergence of the invariant densities obtained by [6, p. 865] for piecewise affine, Markov approximations of topologically mixing piecewise monotone interval maps which are not necessarily Markov). See [11], [9, p. 328] for other approximation schemes, also with results in an L^1 framework, and [16] for some numerical results. See also [14], and references therein, for related results.

This work was made possible by a visit of the first two named authors to the Université de Bourgogne and we are grateful for the warm hospitality enjoyed in the Laboratoire de Topologie.

2. MARKOV TRANSFORMATIONS OF [0, 1] AND THEIR PIECEWISE AFFINE APPROXIMATIONS

Our assumption on the transformation $f : [0, 1] \rightarrow [0, 1]$ is that there exist points $0 = a_0 < a_1 < \dots < a_\ell = 1$ such that

1) for each $i = 1, \dots, \ell$, the restriction $f_i = f|_{]a_{i-1}, a_i[}$ is monotone, C^1 and C^1 -extends to $[a_{i-1}, a_i]$, with a Lipschitz derivative f'_i ; this extension coincides with f at least at one of the endpoints a_{i-1} or a_i ;

2) there is a number $\rho > 1$ such that $|f'_i| \geq \rho$, $i = 1, \dots, \ell$;

3) for any $1 \leq i, j \leq \ell$, if $f(]a_{i-1}, a_i[) \cap]a_{j-1}, a_j[\neq \emptyset$, then $f(]a_{i-1}, a_i[) \supset]a_{j-1}, a_j[$;

4) there exists $k_0 \geq 1$ such that f^{k_0} is onto on each branch of monotonicity (this is equivalent to a topological mixing assumption).

We set $|f'(a_i)| := \min \{ \lim_{y \uparrow a_i} |f'(y)|, \lim_{y \downarrow a_i} |f'(y)| \}$ (with the obvious modification for a_0, a_ℓ).

It is well known that under the above conditions there is a unique absolutely continuous invariant probability measure (a.c.i.p.m.) whose density h is the unique normalised positive eigenfunction with eigenvalue one of a transfer operator \mathcal{L} defined on measurable functions ϕ by

$$\mathcal{L}\phi(x) = \sum_{f(y)=x} \frac{\phi(y)}{|f'(y)|}.$$

Several interesting properties of the dynamical system generated by f are intimately related to $\sigma(\mathcal{L})$, the spectrum of \mathcal{L} (see e.g. [4]). However, the latter depends crucially on the Banach space considered. If one is interested in the unique a.c.i.p.m., one can let \mathcal{L} act on the “large” function space L^1 (Lebesgue). Then, the spectral radius of \mathcal{L} is equal to 1, which is the only element of the spectrum of \mathcal{L} on the unit circle and is a simple eigenvalue with a positive normalised eigenfunction h mentioned above, finally each $z \in \mathbb{C}$ with $|z| < 1$ is an eigenvalue of \mathcal{L} with infinite multiplicity [10].

Recall that given a Banach space of functions on the interval $(\mathcal{B}, \| \cdot \|)$ such that our transfer operator $\mathcal{L} : \mathcal{B} \rightarrow \mathcal{B}$ is bounded, we may define the essential spectral radius, $r_{\text{ess}}(\mathcal{L})(\mathcal{B})$, by:

$$r_{\text{ess}}(\mathcal{L})(\mathcal{B}) = \inf \{ r \in \mathbb{C} : z \in \sigma(\mathcal{L} : \mathcal{B} \rightarrow \mathcal{B}), |z| > r \Rightarrow z \text{ is an isolated eigenvalue of finite multiplicity} \}.$$

We define the discrete spectrum of \mathcal{L} acting on \mathcal{B} to be the set of points z in $\sigma(\mathcal{L})$ with $|z| > r_{\text{ess}}(\mathcal{L})(\mathcal{B})$. If $h \in \mathcal{B}$, $r_{\text{ess}}(\mathcal{L})(\mathcal{B}) < 1$, and if 1 is the only eigenvalue of modulus 1, then, setting

$$\tau := \max \{|\lambda| : \lambda \in \sigma(\mathcal{L}) \setminus \{1\}\} < 1,$$

for any $\phi \in \mathcal{B}$ and any $\varepsilon > 0$, the spectral decomposition of \mathcal{L} yields that

$$\left\| \mathcal{L}^k \phi - \left(\int \phi(x) dx \right) \cdot h \right\| \leq C(\tau + \varepsilon)^k \|\phi\|. \tag{2.1}$$

In other words, τ determines the rate of convergence to equilibrium, also called *rate of mixing*.

We shall consider \mathcal{L} acting on the space of function of bounded variation ($BV, \|\cdot\|$). Recall that the total variation of $\phi : [0, 1] \rightarrow \mathbb{R}$ on an interval $[a, b]$ is

$$\text{var}_{[a,b]} \phi = \sup \left\{ \sum_{i=0}^{n-1} |\phi(x_{i+1}) - \phi(x_i)| : n \geq 1, \right. \\ \left. a \leq x_0 < \dots < x_n \leq b \right\}.$$

Let then $BV := \{\phi : [0, 1] \rightarrow \mathbb{R} : \text{var}_{[0,1]} \phi < \infty\}$ and

$$\|\phi\| = \|\phi\|_\infty + \text{var}_{[0,1]} \phi. \tag{2.2}$$

The spectrum of \mathcal{L} in this setting has been studied by several authors ([18], [7], [14], [10], [2]): the spectral radius is equal to one, \mathcal{L} has 1 as an eigenvalue (and no other eigenvalues on the unit circle), and its essential spectral radius $r_{\text{ess}}(\mathcal{L})(BV)$ is

$$\theta = \lim_{k \rightarrow \infty} (\sup (1/|(f^k)'|))^{1/k} \leq 1/\rho < 1. \tag{2.3}$$

We now construct a sequence $\{f_n\}$ of piecewise affine approximations to f . For any $n \geq 1$ let $\tilde{\mathcal{A}}_n$ be the (mod 0) partition of $[0, 1]$ whose elements are the intervals of the form $I_{i_0} \cap f^{-1}(I_{i_1}) \cap \dots \cap f^{-(n-1)}(I_{i_{n-1}})$, where $I_j =]a_{j-1}, a_j[$ (i.e., the iterate f^n is monotone on each $I \in \tilde{\mathcal{A}}_n$). Let \mathcal{A}_n be the partition (in the strict sense) of $[0, 1]$ obtained by adding one or two endpoints to atoms of $\tilde{\mathcal{A}}_n$, in such a way that $f^k|_I$ is continuous for each $I \in \mathcal{A}_k$ (there might be several ways of doing this).

DEFINITION. – *The n -th piecewise affine approximation of f is the transformation f_n of $[0, 1]$ such that for any $I \in \mathcal{A}_n$ the restriction $f_{n|I}$ is affine, and for any extremal point x of $I \in \mathcal{A}_n$*

$$\lim_{y \uparrow x} f_n(y) = \lim_{y \uparrow x} f(y), \quad \lim_{y \downarrow x} f_n(y) = \lim_{y \downarrow x} f(y)$$

Again, we set $|f'_n(x)| := \min \{ \lim_{y \uparrow x} |f'_n(y)|, \lim_{y \downarrow x} |f'_n(y)| \}$ if x is an endpoint of $I \in \mathcal{A}_n$, with the obvious modification for a_0, a_ℓ .

The transfer operator associated to f_n is

$$\mathcal{L}_n \phi(x) = \sum_{f_n(y)=x} \frac{\phi(y)}{|f'_n(y)|}. \tag{2.4}$$

The results mentioned above apply, in particular 1 is a simple eigenvalue of \mathcal{L}_n acting on BV , with normalised positive eigenfunction h_n equal to the density of the unique a.c.i.p.m., and (2.3) yields a value θ_n for the essential spectral radius.

We now want to characterize the discrete spectrum and the corresponding eigenfunctions of $\mathcal{L}_n : BV \rightarrow BV$. Let Δ_n be the closed \mathcal{L}_n -invariant subspace of BV defined by

$$\Delta_n = \text{span} \{ \chi_I : I \in \mathcal{A}_n \}. \tag{2.5}$$

Then, for any $n \geq 1$, we may consider the restricted operator $\mathcal{L}_{n|\Delta_n}$, and the coinduced operator \mathcal{L}_n^Γ , acting on the quotient space $\Gamma_n = BV/\Delta_n$. We have the decomposition [5] $\sigma(\mathcal{L}_n) \subseteq \sigma(\mathcal{L}_{n|\Delta_n}) \cup \sigma(\mathcal{L}_n^\Gamma)$. Moreover, it is easy to check that in the natural basis for Δ_n , the restriction $\mathcal{L}_{n|\Delta_n}$ is given by the $\ell^n \times \ell^n$ matrix $M^{(n)}$ defined by:

$$M_{ij}^{(n)} = \frac{|J_j \cap f_n^{-1}(J_i)|}{|J_i|} = \frac{|J_j|}{\sum_n |J_k \cap f(J_k)|}, \quad J_i, J_j \in \mathcal{A}_n$$

By the mixing assumption, the matrix $M^{(n)}$ is irreducible and aperiodic, so that $\sigma(M^{(n)}) = \{1\} \cup \sigma_n$, where σ_n is strictly contained in the unit disc. On the other hand, the norm induced by (2.2) on Γ_n is $\|\phi\|^\Gamma = \|\phi - Q_n \phi\|$, where $Q_n : BV \rightarrow \Delta_n$ is given by

$$Q_n \phi(x) = \sum_{I \in \mathcal{A}_n} \alpha_I(\phi) \chi_I(x), \quad \alpha_I(\phi) = \frac{1}{|I|} \int_I \phi(s) ds,$$

so that the spectral radius of $\mathcal{L}_n^\Gamma : BV \rightarrow BV$ is easily seen to be $\leq \theta_n$ and thus $= \theta_n$ (see e.g. [1]). Putting together these observations we have:

LEMMA 2.1.

$$\begin{aligned} \{z \in \sigma(\mathcal{L}_n), |z| > \theta_n\} &= \{z \in \sigma(\mathcal{L}_n|_{\Delta_n}), |z| > \theta_n\} \\ &= \{z \in \sigma(M^{(n)}), |z| > \theta_n\}. \end{aligned}$$

In particular, the normalised fixed function h_n of \mathcal{L}_n is in Δ_n , and is a fixed vector of the matrix $M^{(n)}$.

We are now in a position to state our main result.

THEOREM. – Let f be a piecewise monotone interval map satisfying assumptions (1)-(4) from the beginning of this section, let h be the density of its unique a.c.i.p.m. and τ its rate of mixing. Let \mathcal{L}_n be the transfer operators of the n -th piecewise affine approximation of f .

Then there is a constant $C_1 > 0$ and for each $\xi^2 > \tau$ a constant $C_2 > 0$ such that for each $n \geq 1$ the normalised eigenfunction $h_n \in \Delta_n$ for the simple eigenvalue 1 of \mathcal{L}_n satisfies

$$\begin{aligned} \text{var } h_n &\leq C_1 \\ \sup_{x \in [0, 1]} |h_n(x) - h(x)| &\leq C_2 \xi^{(2/3)n}. \end{aligned}$$

Moreover, the spectrum of \mathcal{L}_n decomposes as $\{1\} \cup \Sigma_n$ with rate of mixing $\tau_n = \sup \{z \in \Sigma_n\} < 1$ and if $\tau \neq r_{\text{ess}}(\mathcal{L})$, the rates τ_n converge to the rate of mixing of f , i.e., $\lim_{n \rightarrow \infty} \tau_n = \tau$.

Remark. – We only prove convergence of the maximal eigenvector. This is related to the fact that the nature of our approximations forces the use of balanced norms (see below). It would be interesting to know whether a different approach would yields results on eigenvectors corresponding to other elements of the discrete spectrum. Note also that we do not know whether the obtained exponential rate of convergence ($\tau^{1/3}$) is optimal.

The proof of our theorem uses two lemmas. Lemma 2.2 is the analogue of the ‘‘dynamical’’ Lemma 9 in [3] and Lemma 2.3 corresponds to the ‘‘abstract functional lemmas’’ in [3]. The Markov situation considered here yields a simplification and a strenghtening of the results, because there are no ‘‘bad’’ intervals of monotonicity (in the terminology of [3]).

We first make some preliminary remarks. For each $\tilde{\theta} > \theta$, there exists $k_1 \geq 1$ so that for all $n \geq k \geq k_1$ and all $I \in \mathcal{A}_k$,

$$\sup_{x \in \bar{I}} \frac{1}{|(f_n^k)'(x)|} \leq \sup_{x \in \bar{I}} \frac{1}{|(f^k)'(x)|} \leq \tilde{\theta}^k, \tag{2.6}$$

and also (see [2, Lemma 2.3])

$$\text{var}_{\bar{I}} \frac{1}{|(f_n^k)'(x)|} \leq \text{var}_{\bar{I}} \frac{1}{|(f^k)'(x)|} \leq \tilde{\theta}^k. \tag{2.7}$$

(We may assume that k_1 is a multiple of k_0 : this will be convenient below.) Observe also that for any $\tilde{\theta} > \theta$, there exists a constant C such that for all $k \geq 1$ the maximum length of the intervals in \mathcal{A}_k is not larger than

$$\sup_{I \in \mathcal{A}_k} \sup_{x \in \bar{I}} \frac{1}{|(f^k)'(x)|} \leq C \tilde{\theta}^k. \tag{2.8}$$

It will be necessary to make use of balanced norms in BV : for $0 < \gamma \leq 1$ define

$$\|\phi\|_\gamma = \|\phi\|_\infty + \gamma \text{var}_{[0,1]} \phi.$$

LEMMA 2.2. – Let $\theta < \xi^2 < 1$. Then there exists $C_3 > 0$ such that for all $n \geq (3/2)k \geq k \geq k_1$

$$\|\mathcal{L}^k - \mathcal{L}_n^k\|_{\xi^k} \leq C_3 \xi^k.$$

Proof of Lemma 2.2. – Fix $\tilde{\theta}$ with $\theta < \tilde{\theta} < \xi^2$.

We start by preliminary computations useful to control the supremum part of the norm. For $n \geq k \geq k_1$, we denote by $(\psi_I^k)_{I \in \mathcal{A}_n}$ and $(\psi_{n,I}^k)_{I \in \mathcal{A}_n}$ the collection of the inverse branches of all the restrictions to the atoms of \mathcal{A}_n of f^k and f_n^k respectively. Since $n \geq k$, we have for $\phi \in BV$:

$$\begin{aligned} \mathcal{L}^k \phi(x) &= \sum_{I \in \mathcal{A}_n} \frac{\chi_{f^k(I)}(x) \phi(\psi_I^k(x))}{|(f^k)'(\psi_I^k(x))|}, \\ \mathcal{L}_n^k \phi(x) &= \sum_{I \in \mathcal{A}_n} \frac{\chi_{f_n^k(I)}(x) \phi(\psi_{n,I}^k(x))}{|(f_n^k)'(\psi_{n,I}^k(x))|} \end{aligned}$$

where the characteristic functions $\chi_{f^k(I)} = \chi_{f_n^k(I)}$ are the same in both sums by the definition of f_n . Therefore

$$\begin{aligned} &|\mathcal{L}^k \phi(x) - \mathcal{L}_n^k \phi(x)| \\ &\leq \sum_{I \in \mathcal{A}_n} \frac{|\phi(\psi_I^k(x)) - \phi(\psi_{n,I}^k(x))|}{|(f^k)'(\psi_I^k(x))|} \chi_{f^k(I)}(x) \\ &\quad + \sum_{I \in \mathcal{A}_n} |\phi(\psi_{n,I}^k(x))| \left| \frac{1}{(f^k)'(\psi_I^k(x))} - \frac{1}{(f_n^k)'(\psi_{n,I}^k(x))} \right| \chi_{f^k(I)}(x) \\ &= I + II. \end{aligned} \tag{2.9}$$

A straightforward calculation using (2.6) yields

$$I \leq \text{var}_{[0,1]}(\phi) \tilde{\theta}^k. \tag{2.10}$$

The second term can be estimated as follows. Let $K > 0$ be such that $\mathcal{L}^k 1(x) \leq K$ for all $k \geq 1$ (such a constant K exists because $\|\mathcal{L}^k 1\|_\infty \leq \int \mathcal{L}^k 1 dx + \text{var}_{[0,1]} \mathcal{L}^k 1 = \|\mathcal{L}^k 1\|$, and (2.1) implies that $\|\mathcal{L}^k 1\| \leq \|h\| + \text{cst}(\tau + \varepsilon)^k$) and let $C > 0$ be the constant from (2.8). Then we will prove that there is a constant \bar{C} such that for all $n \geq k$:

$$\begin{aligned} II &\leq \|\phi\|_\infty \sup_{y_1, y_2 \in I \in \mathcal{A}_n} \left| \frac{(f^k)'(y_1)}{(f_n^k)'(y_2)} - 1 \right| \\ &\quad \times \sum_{I \in \mathcal{A}_n} \left| \frac{1}{(f^k)'(\psi_I^k(x))} \right| \chi_{f^k(I)}(x) \\ &\leq \|\phi\|_\infty K \bar{C} C \tilde{\theta}^{n-k}. \end{aligned} \tag{2.11}$$

To obtain (2.11) we will use the following distortion inequality: if f is piecewise $C^{1+\text{Lipschitz}}$ and expanding (in particular, $\log|f'|$ is piecewise Lipschitz), there is a constant $\hat{C} > 0$ so that for each interval J of monotonicity of f and all $x_1, x_2 \in J$,

$$\left| \frac{f'(x_1)}{f'(x_2)} - 1 \right| \leq \hat{C} |x_1 - x_2|.$$

To apply this inequality, we first observe that for all $n \geq k \geq 1$, and all $y_1, y_2 \in I \in \mathcal{A}_n$, the mean value theorem (and the remark that $f^m(I) = f_n^m(I)$ for all $0 \leq m \leq k$), yields

$$\frac{(f^k)'(y_1)}{(f_n^k)'(y_2)} = \prod_{m=0}^{k-1} \frac{f'(f^m(y_1))}{f_n'(f_n^m(y_2))} = \prod_{m=0}^{k-1} \frac{f'(f^m(y_1))}{f'(f_n^m(y_{2,m}))}$$

where each $y_{2,m}$ is in I . We now combine the distortion inequality with a second application of the mean value theorem and obtain points

$z_m = z_m(y_2) \in I$, and a constant $\bar{C} > 0$ with

$$\begin{aligned} 0 \leq \frac{(f^k)'(y_1)}{(f_n^k)'(y_2)} &\leq \prod_{m=0}^{k-1} (1 + \hat{C} |f^m(y_1) - f^m(z_m)|) \\ &\leq \prod_{m=0}^{k-1} (1 + \hat{C} \rho^{m-k} |f^k(y_1) - f^k(z_m)|) \\ &\leq \exp \sum_{\ell=1}^k \log (1 + \hat{C} \rho^{-\ell} |f^k(y_1) - f^k(z_{k-\ell})|) \\ &\leq \sup_{y_3 \in I} (1 + \bar{C} |f^k(y_1) - f^k(y_3)|). \end{aligned}$$

Since $f^k(y_1)$ and $f^k(y_3)$ are in the same atom of \mathcal{A}_{n-k} we may apply (2.8). It then suffices to exchange numerator and denominator to get the other inequality required for (2.11). (On distortion inequalities, see e.g. [12, I.2, V.2].)

Therefore, using $\tilde{\theta} < \xi^2$, there is $D_1 > 0$ so that for all $n \geq (3/2)k \geq k \geq k_1$ and $\phi \in BV$

$$\begin{aligned} |\mathcal{L}^k \phi(x) - \mathcal{L}_n^k \phi(x)| &\leq \tilde{\theta}^k \text{var}_{[0,1]} \phi + \|\phi\|_\infty K \bar{C} \tilde{\theta}^{n-k} \\ &\leq D_1 \xi^k \|\phi\|_{\xi^k}. \end{aligned} \tag{2.12}$$

To estimate the variation, we write

$$\text{var}_{[0,1]} (\mathcal{L}^k \phi - \mathcal{L}_n^k \phi) \leq \text{var}_{[0,1]} \mathcal{L}_n^k \phi + \text{var}_{[0,1]} \mathcal{L}^k \phi$$

and bound each term of the right-hand-side separately. We consider first \mathcal{L}_n^k for $k = k_1$ where k_1 is a multiple of k_0 . We may assume that $2\tilde{\theta}^{k_1} < \xi^{k_1}$ (otherwise take k_1 to be a larger multiple of k_0). Using the fact that for each $I \in \mathcal{A}_{k_1}$ the branch $f_{|I}^{k_1}$ can be extended to a surjective function on the closure of I (and the inverse branches accordingly), we find for each $I \in \mathcal{A}_{k_1}$:

$$\begin{aligned}
& \text{var}_{[0,1]} \frac{\chi_{f^{k_1}(\bar{I})}(x) \phi(\psi_{n,\bar{I}}^{k_1}(x))}{|(f_n^{k_1})'(\psi_{n,\bar{I}}^{k_1}(x))|} \\
&= \text{var}_{[0,1]} \frac{\phi(\psi_{n,\bar{I}}^{k_1}(x))}{|(f_n^{k_1})'(\psi_{n,\bar{I}}^{k_1}(x))|} \\
&\leq \sup_{x \in \bar{I}} \frac{1}{|(f_n^{k_1})'(x)|} \text{var}_{\bar{I}} \phi + \sup_{\bar{I}} |\phi| \text{var}_{\bar{I}} \frac{1}{|(f_n^{k_1})'|} \\
&\leq \tilde{\theta}^{k_1} \text{var}_{\bar{I}} \phi + \tilde{\theta}^{k_1} \sup_{\bar{I}} |\phi| \\
&\leq 2 \tilde{\theta}^{k_1} \text{var}_{\bar{I}} \phi + \tilde{\theta}^{k_1} \inf_{\bar{I}} |\phi| \\
&\leq \xi^{k_1} \text{var}_{\bar{I}} \phi + \tilde{\theta}^{k_1} \beta(k_1)^{-1} \int_{\bar{I}} |\phi(x)| dx,
\end{aligned}$$

where $\beta(k_1) > 0$ is the minimum length of the atoms of \mathcal{A}_{k_1} . Hence taking the sum over the atoms of \mathcal{A}_{k_1} we find

$$\text{var}_{[0,1]} \mathcal{L}_n^{k_1} \phi \leq \xi^{k_1} \text{var}_{[0,1]} \phi + \tilde{\theta}^{k_1} \beta(k_1)^{-1} \int_0^1 |\phi(x)| dx.$$

Finally, applying recursively the above inequality (and using $\int |\mathcal{L}_n \phi| dx = \int |\phi| dx$) we find a constant $D_2 > 0$ so that for all $n \geq k = m \cdot k_1 + p \geq k_1$ ($p < k_1$):

$$\text{var}_{[0,1]} \mathcal{L}_n^k \phi \leq D_2 \left(\xi^k \text{var}_{[0,1]} \phi + \tilde{\theta}^{k_1} \beta(k_1)^{-1} \frac{1 - \xi^k}{1 - \xi^{k_1}} \int |\phi(x)| dx \right).$$

The estimate for \mathcal{L}^k is exactly the same:

$$\text{var}_{[0,1]} \mathcal{L}^k \phi \leq D_2 \left(\xi^k \text{var}_{[0,1]} \phi + \tilde{\theta}^{k_1} \beta(k_1)^{-1} \frac{1 - \xi^k}{1 - \xi^{k_1}} \int |\phi(x)| dx \right).$$

Putting together these estimates and using $\int |\phi| dx \leq \|\phi\|_\infty$, we find a constant $D_3 > 0$ such that for all $n \geq k \geq k_1$

$$\text{var}_{[0,1]} (\mathcal{L}^k \phi - \mathcal{L}_n^k \phi) \leq D_3 \xi^k \text{var}_{[0,1]} \phi + D_3 \|\phi\|_\infty. \quad (2.13)$$

From (2.12) and (2.13) and using again $\tilde{\theta} < \xi^2$, we obtain $C_3 > 0$ so that for all $n \geq (3/2)k \geq k \geq k_1$

$$\|\mathcal{L}^k - \mathcal{L}_n^k\|_{\xi^k} \leq C_3 \xi^k.$$

Q.E.D.

We decompose the spectrum of \mathcal{L} on BV into $\sigma(\mathcal{L}) = \Sigma_0 \cup \Sigma_1$ where $\Sigma_0 = \{1\}$, with the corresponding decomposition into generalised eigenspaces $BV = X_0 \oplus X_1 = \mathbb{C}h \oplus X_1$, and projections $\pi_0 : BV \rightarrow X_0$, $\pi_1 : BV \rightarrow X_1$ (see e.g. [8]).

LEMMA 2.3. – Consider the operators \mathcal{L}_n acting on BV and let $\theta < \xi^2 < 1$. For any ξ' so that $\tau/\xi' < \xi < \xi' < 1$, there exist $C_4 > 0$ and $k_2 \geq 1$ so that for all $n \geq (3/2)k \geq k \geq k_2$:

1) The spectrum $\sigma(\mathcal{L}_n)$ decomposes into

$$\sigma(\mathcal{L}_n) = \Sigma_0^n \cup \Sigma_1^n$$

with

$$\sup \{|z| \mid z \in \Sigma_1^n\} < \xi' < \inf \{|z| \mid z \in \Sigma_0^n\}.$$

2) Let $\pi_0^n : X_0^n \oplus X_1^n \rightarrow X_0^n$ be the projection associated with this spectral decomposition, then for each $\eta > \xi/\xi'$

$$\|\pi_0 - \pi_0^n\|_{\xi^k} < C_4 \eta^k.$$

(In particular, $\Sigma_0^n = \{1\}$ and 1 is a simple eigenvalue of \mathcal{L}_n .)

Proof of Lemma 2.3 (We essentially follow [3]). – 1) Let τ' and ξ'_0 be such that

$$\frac{\tau}{\xi'} < \frac{\tau'}{\xi'} < \xi < \xi' < \xi'_0 < 1.$$

Let $k_2 \geq k_1$ be a fixed multiple of k_0 , large enough for various purposes. In particular, we require that for $k \geq k_2$

$$\phi \in X_1 \Rightarrow \|\mathcal{L}^k \phi\| \leq (\tau')^k \|\phi\|.$$

For $n \geq (3/2)k \geq k \geq k_2$ we will show that $\lambda \notin \sigma(\mathcal{L}_n)$ for λ with $\xi' < |\lambda| < \xi'_0$ by proving that the resolvent $R(\mathcal{L}_n^k, \lambda^k) = (\mathcal{L}_n^k - \lambda^k \text{Id})^{-1}$ is a bounded operator on $(BV, \|\cdot\|_{\xi^k})$. If the resolvent exists, it can be written as:

$$R(\mathcal{L}_n^k, \lambda^k) = \sum_{m=0}^{\infty} (R(\mathcal{L}^k, \lambda^k) (\mathcal{L}_n^k - \mathcal{L}^k))^m \cdot R(\mathcal{L}^k, \lambda^k). \tag{2.14}$$

By Lemma 2.2, it is enough to show that $\|R(\mathcal{L}^k, \lambda^k)\|_{\xi^k} < (1/\xi)^k$. Since $R(\mathcal{L}^k, \lambda^k) X_i = X_i$ for $i = 0, 1$, we have for $\phi \in X$, $\|\phi\|_{\xi^k} = 1$

$$\begin{aligned} \|R(\mathcal{L}^k, \lambda^k) \phi\|_{\xi^k} &\leq \|R(\mathcal{L}^k, \lambda^k) \pi_0 \phi\|_{\xi^k} + \|R(\mathcal{L}^k, \lambda^k) \pi_1 \phi\|_{\xi^k} \\ &\leq \|R(\mathcal{L}^k, \lambda^k)|_{X_0}\|_{\xi^k} \|\pi_0\|_{\xi^k} \\ &\quad + \|R(\mathcal{L}^k, \lambda^k)|_{X_1}\|_{\xi^k} \|\pi_1\|_{\xi^k}. \end{aligned}$$

Since $\pi_0 \phi = h \int \phi(x) dx$, there exists a constant $A_0 > 0$ with

$$\|\pi_0 \phi\|_{\infty} \leq \|h\|_{\infty} \left| \int \phi(x) dx \right| \leq A_0 \|\phi\|_1 \leq A_0 \|\phi\|_{\infty}.$$

Therefore $\|\pi_0\|_{\xi^k}$ and $\|\pi_1\|_{\xi^k} \leq 1 + \|\pi_0\|_{\xi^k}$ are uniformly bounded (because $\|\pi_0 \phi\|_{\xi^k} \leq \|\pi_0 \phi\| \leq A' \|\pi_0 \phi\|_{\infty} \leq A' A_0 \|\phi\|_{\infty} \leq A' A_0 \|\phi\|_{\xi^k}$, where we have used that all norms on X_0 are equivalent). It thus suffices to bound $\|R(\mathcal{L}^k, \lambda^k)|_{X_i}\|_{\xi^k}$, $i = 0, 1$.

There exists a constant $A_1 > 0$ that for all $\phi \in X_0$

$$\|\mathcal{L}^k \phi - \lambda^k \phi\|_{\xi^k} \geq A_1 \cdot \|\phi\|_{\xi^k}.$$

For $\phi \in X_1$, we have

$$\|\mathcal{L}^k \phi\|_{\xi^k} \leq \|\mathcal{L}^k \phi\| \leq (\tau')^k \|\phi\| \leq \left(\frac{\tau'}{\xi}\right)^k \|\phi\|_{\xi^k},$$

from which it follows that there is a constant $A_2 > 0$ with

$$\|\mathcal{L}^k \phi - \lambda^k \phi\|_{\xi^k} \geq A_2 \cdot (\xi')^k \|\phi\|_{\xi^k}.$$

Therefore, there is a constant $A_3 > 0$ so that for all large enough k

$$\|R(\mathcal{L}^k, \lambda^k)\|_{\xi^k} \leq \frac{A_3}{(\xi')^k} \leq \frac{1}{\xi^k}. \tag{2.15}$$

2) Note that π_0 can be viewed as the projection associated with $(\mathcal{L}^k, (\Sigma_0)^k)$ for any k , and similarly for π_0^n . We will again consider a fixed $k \geq k_2$ and $n \geq (3/2)k$. We write B_δ for the circle of radius δ centered at 0 in \mathbb{C} . Let ξ'_0 be as in part (1) and let $\gamma = B_{\xi'_k} \cup B_{r_0^k}$ for some $\xi' < \hat{\xi} < \xi'_0$, with $\hat{\xi} < \eta(\xi')^2/\xi$ (using $\eta(\xi')^2/\xi > \xi'$) and $r_0 > 1$. Then by (1), Σ_0^k and $(\Sigma_0^n)^k$ are contained in the annular region bounded by γ , and we have

$$\pi_0 = \frac{1}{2i\pi} \int_{\gamma} R(\mathcal{L}^k, \lambda) d\lambda \quad \pi_0^n = \frac{1}{2i\pi} \int_{\gamma} R(\mathcal{L}_n^k, \lambda) d\lambda.$$

Therefore

$$\begin{aligned} \|\pi_0 - \pi_0^n\|_{\xi^k} &\leq \frac{1}{2\pi} \int_{\gamma} \|R(\mathcal{L}^k, \lambda) - R(\mathcal{L}_n^k, \lambda)\|_{\xi^k} d\lambda \\ &\leq \frac{1}{2\pi} \cdot \ell(B_{\xi^k}) \max_{\lambda \in B_{\xi^k}} \|R(\mathcal{L}^k, \lambda) - R(\mathcal{L}_n^k, \lambda)\|_{\xi^k} \\ &\quad + \text{the corresponding term for } B_{r_0^k} \\ &= III + IV. \end{aligned}$$

Using (2.14) we have

$$\|R(\mathcal{L}^k, \lambda) - R(\mathcal{L}_n^k, \lambda)\|_{\xi^k} \leq \sum_{m=1}^{\infty} \|R(\mathcal{L}^k, \lambda)\|_{\xi^k}^{m+1} \cdot \|\mathcal{L}_n^k - \mathcal{L}^k\|_{\xi^k}^m.$$

Since $\ell(B_{\xi^k}) = 2\pi\hat{\xi}^k$, and $\|R(\mathcal{L}^k, \lambda)\| \leq A_3/(\xi')^k$ for $\lambda \in B_{\xi^k}$ [by (2.15)], we obtain $A_4 > 0$ such that

$$III \leq \hat{\xi}^k \sum_{m=1}^{\infty} \left(\frac{A_3}{\xi'^k}\right)^{m+1} (\xi^k)^m \leq A_4 \hat{\xi}^k \cdot \frac{\xi^k}{(\xi'^k)^2} \leq A_4 \eta^k.$$

For IV, we use $\ell(B_{r_0^k}) = 2\pi r_0^k$, to get $A_5 > 0$ so that

$$IV \leq A_5 r_0^k \frac{\xi^k}{r_0^{2k}} \leq A_5 \eta^k.$$

The assertion in parenthesis follows from classical perturbation results, see e.g. [8].

Q.E.D.

Proof of the Theorem. – Let $\tau/\xi' < \xi < \xi' < 1$. First we show that for $n \geq (3/2)k \geq k \geq k_2$ the space X_0^n from Lemma 2.3 is the graph of some linear $S_n : X_0 \rightarrow X_1$: Let $\phi \in X_0^n$, since

$$\begin{aligned} \|\pi_1 \phi\|_{\xi^k} &= \|\phi - \pi_0 \phi\|_{\xi^k} \leq \|\pi_0^n - \pi_0\|_{\xi^k} \|\phi\|_{\xi^k}, \\ \|\pi_0 \phi\|_{\xi^k} &\geq \|\phi\|_{\xi^k} (1 - \|\pi_0 - \pi_0^n\|_{\xi^k}), \end{aligned}$$

it follows from Lemma 2.3 (2) that $\|\pi_1 \phi\|_{\xi^k} \ll \|\pi_0 \phi\|_{\xi^k}$. In particular if $\phi, \phi' \in X_0^n$ and $\pi_0 \phi = \pi_0 \phi'$, then $\pi_1 \phi = \pi_1 \phi'$ and thus $\phi = \phi'$.

We now estimate $\|S_n\|_{\xi^k}$. Since $\dim X_0 = 1$, and $\Sigma_0^n = \{1\}$, there exists $\phi_0 = \phi_0(n, k) \in X_0$, $\|\phi_0\|_{\xi^k} = 1$, such that

$$\|S_n\|_{\xi^k} = \|S_n \phi_0\|_{\xi^k} = \|\pi_1(\phi_0, S_n \phi_0)\|_{\xi^k} = \|\pi_1 \mathcal{L}_n^k(\phi_0, S_n \phi_0)\|_{\xi^k}.$$

For $\xi^2 > \tau'' > \tau$ and large enough k , Lemma 2.2 thus yields

$$\begin{aligned} \|S_n\|_{\xi^k} &= \|\pi_1 [\mathcal{L}_n^k(\phi_0, S_n \phi_0) - \mathcal{L}^k \phi_0 + \mathcal{L}^k S_n \phi_0 - \mathcal{L}^k S_n \phi_0]\|_{\xi^k} \\ &\leq \|\pi_1\|_{\xi^k} ((\tau''/\xi)^k + C_3 \xi^k) \|S_n\|_{\xi^k} + C_3 \xi^k. \end{aligned}$$

Therefore, there exist $k_3 \geq k_2$ and a constant $B > 0$ so that for all $n \geq (3/2)k \geq k \geq k_3$

$$\|S_n\|_{\xi^k} \leq B \xi^k. \quad (2.16)$$

In particular, writing $[y]$ for the integer part of y , for all $n \geq (3/2)k_3$

$$\|S_n\|_{\xi^{[(2/3)n]}} \leq B \xi^{[(2/3)n]}. \quad (2.17)$$

We need a bound on $\|S_n\|_\infty$. For $n \geq (3/2)k_3$ and $\phi \in X_0$ we have, using the constant A' from the proof of Lemma 2.3 (1),

$$\begin{aligned} \|S_n \phi\|_\infty &\leq \|S_n\|_{\xi^{[(2/3)n]}} \|\phi\|_{\xi^{[(2/3)n]}} \\ &\leq \|S_n\|_{\xi^{[(2/3)n]}} (\xi^{[(2/3)n]} \cdot A' \cdot \|\phi\|_\infty \\ &\quad + (1 - \xi^{[(2/3)n]}) \cdot \|\phi\|_\infty) \\ &\leq (A' + 1) B \xi^{[(2/3)n]} \|\phi\|_\infty. \end{aligned}$$

By definition

$$h_n = \frac{h + S_n(h)}{\int (h(x) + S_n(h)(x)) dx}.$$

Now $\left| \int (h(x) + S_n(h)(x)) dx - 1 \right| \leq \|S_n(h)\|_\infty$, which tends to zero as $n \rightarrow \infty$ by (2.18). Therefore (2.17) implies that for $n \geq (3/2)k_3$:

$$\begin{aligned} \text{var } h_n &\leq \frac{\text{var } h + \text{var } S_n h}{\int (h + S_n h) dx} \\ &\leq \frac{\text{var } h + \xi^{-[(2/3)n]} \|S_n\|_{\xi^{[(2/3)n]}} \|h\|}{1 - \|S_n(h)\|_\infty} \\ &\leq (1 + B) \|h\| (1 + 2 \|S_n(h)\|_\infty), \end{aligned}$$

which proves the existence of C_1 . We now bound $\|h_n - h\|_\infty$ using again (2.18):

$$\begin{aligned} \|h_n - h\|_\infty &\leq \frac{(1 + \|h\|_\infty) \|S_n h\|_\infty}{1 - \|S_n h\|_\infty} \\ &\leq (1 + \|h\|_\infty) (1 + 2 \|S_n h\|_\infty) (A' + 1) B \xi^{[(2/3)n]} \|h\|_\infty, \end{aligned}$$

which proves the existence of C_2 .

The statement on the convergence of τ_n follows from the property of the convergence of the discrete spectrum in [1], Corollary to the main theorem.

Q.E.D.

REFERENCES

- [1] V. BALADI, Infinite kneading matrices and weighted zeta functions of interval maps, Preprint, 1933, to appear in *J. Funct. Anal.*
- [2] V. BALADI and G. KELLER, Zeta functions and transfer operators for piecewise monotone transformations, *Comm. Math. Phys.*, Vol. **127**, 1990, pp. 459-477.
- [3] V. BALADI and L.-S. YOUNG, On the spectra of randomly perturbed expanding maps, *Comm. Math. Phys.*, Vol. **156**, 1993, pp. 355-385. (See also Erratum, *Comm. Math. Phys.*, Vol. **166**, 1994, pp. 219-220.)
- [4] P. COLLET, Some ergodic properties of maps of the interval, *Dynamical Systems and Frustrated Systems*, R. Bamon, J.-M. Gambaudo and S. Martinez Eds., 1991 (to appear).
- [5] I. ERDELYI and R. LANGE, Spectral Decompositions on Banach Spaces, *Lecture Notes in Math.*, Vol. **623**, 1977.
- [6] P. GORA and A. BOYARSKY, Compactness of invariant densities for families of expanding, piecewise monotonic transformations, *Can. J. Math.*, **XLI**, 1989, pp. 855-869.
- [7] F. HOFBAUER and G. KELLER, Ergodic properties of invariant measures for piecewise monotonic transformations, *Math. Z.*, Vol. **180**, 1982, pp. 119-140.
- [8] T. KATO, *Perturbation Theory for Linear Operators*, Second printing of second ed., Springer-Verlag, Berlin, 1984.
- [9] G. KELLER, Stochastic stability in some chaotic dynamical systems, *Mh. Math.*, Vol. **94**, 1982, pp. 313-333.
- [10] G. KELLER, On the rate of convergence to equilibrium in one-dimensional dynamics, *Comm. Math. Phys.*, Vol. **96**, 1984, pp. 181-193.
- [11] T.-Y. LI, Finite approximation for the Frobenius-Perron operator. A solution to Ulam's conjecture, *J. of Approx. Theory*, Vol. **17**, 1976, pp. 177-186.
- [12] W. DE MELO and S. VAN STRIEN, *One-Dimensional Dynamics*, Springer-Verlag, Berlin, 1993.
- [13] D. RUELLE, *Thermodynamic Formalism*, Addison-Wesley, Reading, 1978.
- [14] M. RYCHLIK, Bounded variation and invariant measures, *Studia Math.*, Vol. **LXXVI**, 1983, pp. 69-80.
- [15] M. RYCHLIK and E. SORETS, Regularity and other properties of absolutely continuous invariant measures for the quadratic family, *Comm. Math. Phys.*, Vol. **150**, 1992, pp. 217-236.
- [16] H. J. SCHOLTZ, Markov chains and discrete chaos, *Physica*, Vol. **4D**, 1982, pp. 281-286.,
- [17] S. ULAM, *Problems in modern mathematics*, Interscience Publishers, New York, 1960.
- [18] S. WONG, Some metric properties of piecewise monotonic mappings of the unit interval, *Trans. Amer. Math. Soc.*, Vol. **246**, 1978, pp. 493-500.

(Manuscript received September 9, 1994;
revised version December 20, 1994.)