

# ANNALES DE L'I. H. P., SECTION C

ANTONIO AMBROSETTI

VITTORIO COTI ZELATI

## **Non-collision orbits for a class of keplerian-like potentials**

*Annales de l'I. H. P., section C*, tome 5, n° 3 (1988), p. 287-295

[http://www.numdam.org/item?id=AIHPC\\_1988\\_\\_5\\_3\\_287\\_0](http://www.numdam.org/item?id=AIHPC_1988__5_3_287_0)

© Gauthier-Villars, 1988, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## **Non-collision orbits for a class of Keplerian-like potentials**

by

**Antonio AMBROSETTI** <sup>(1)</sup>

Scuola Normale Superiore,  
Piazza dei Cavalieri, 56100 Pisa, Italy

and

**Vittorio COTI ZELATI** <sup>(1), (2)</sup>

CEREMADE, Université de Paris-Dauphine,  
Place du M.-de-Lattre-de-Tassigny, 75775 Paris Cedex 16, France

---

**ABSTRACT.** — We prove the existence of non-collision orbits with large period for a class of Keplerian-like dynamical systems.

*Key words* : Periodic solutions, second order dynamical systems.

---

*Classification A.M.S.* : 49 A 40, 58 F 22.

<sup>(1)</sup> Supported by M.P.I., Gruppo Nazionale (40%) "Calcolo delle Variazioni...".

<sup>(2)</sup> Supported by a C.N.R. scholarship. Permanent adress S.I.S.S.A., strada Costiera 11, 34014 Trieste, Italy.

RÉSUMÉ. — Nous prouvons l'existence d'orbites de non-collision avec grand period pour une classe de systèmes dynamiques de type keplerien.

## 0. INTRODUCTION

In this paper we study the existence of  $T$ -periodic solutions for a system of ordinary differential equations of the form

$$-\ddot{y} = V'(y), \quad (1)$$

where  $V$  is a Keplerian-like potential, i. e.  $V(x)$  behaves like  $-|x|^{-\alpha}$  for  $x$  close to 0,  $\alpha$  being any real number greater than 0. We prove that, for large  $T$ , such a system has a non-collision  $T$ -periodic solution (i. e. a solution which does not cross the origin) under the only assumption that  $V$  attains its maximum on the boundary of an open set which contains the origin.

A potential of such kind arises, for example, if at  $x=0$  there are  $z$  positive charges surrounded by  $z+k$  ( $k>0$ ) negative ones uniformly distributed on a shell containing  $x=0$ . Then  $V(x) = -z/|x|$  inside the shell, while  $V(x) = k/|x|$  at infinity.

The existence of periodic solutions of (1) when  $V' \approx -|x|^{-\alpha}$  and  $\alpha \geq 2$  (or, more precisely, the case of strong forces—see [6] for a definition) has been studied in [1], [3], [6], [8], see also [2] for a review of the results in this and related fields. We notice that in such a case all the periodic solutions are non-collision orbits.

The situation is much more complicated when  $\alpha \leq 2$ . For some partial results for  $\alpha > 1$  we refer to [5] (see also [4] for a somewhat different class of potentials). In particular, the results of [5] do not cover the case  $\alpha = 1$  <sup>(3)</sup>, which is known to be quite degenerate. For example, if

---

<sup>(3)</sup> Some results are found also when  $\alpha = 1$  but under other symmetry conditions.

$V(x) = -|x|^{-1}$ , then the  $T$ -periodic solutions belong to one parameter families containing collision solutions and all the orbits of each family have the same value of the energy and the same value of the action functional [7].

Actually, in the present paper, we show that Kepler's potential is very sensitive to perturbation, at least in the sense that even a very small perturbation far from the singular set (if it goes in the "right" direction) can assure the existence of non-collision orbits.

The results proved here have been announced in the *C.R. Acad. Sci. Paris* note [0].

## 1. ASSUMPTIONS AND MAIN EXISTENCE RESULTS

We consider a potential  $V \in C^1(\mathbb{R}^N \setminus \{0\}, \mathbb{R})$  satisfying

(V1)  $V(x) \rightarrow -\infty$  as  $|x| \rightarrow 0$ ;

(V2) there exists an open, bounded set  $\Omega \subset \mathbb{R}^N$ , with smooth boundary  $\Gamma$  such that

(i)  $0 \in \Omega$  and  $\Omega$  is star-shaped with respect to 0;

(ii) letting  $b = \max \{ V(x) : x \in \mathbb{R}^N \setminus \{0\} \}$ , one has that  $b = V(\xi)$ ,  $\forall \xi \in \Gamma$ ;

(V3)  $\limsup_{|x| \rightarrow +\infty} V(x) = \beta < b$ .

Given  $T > 0$  we look for solutions of

$$-\ddot{y} = V'(y), \quad y(0) = y(T), \quad \dot{y}(0) = \dot{y}(T) \quad (P_T)$$

where  $V'$  denotes the gradient of  $V$ .

We say that a solution  $y(t)$  of (P) is a non-collision orbit if  $y(t) \neq 0$ ,  $\forall t$ .

Let  $S^1 = [0, 1] / \{0, 1\}$ ,  $H = H^1(S^1; \mathbb{R}^N)$ , and  $\Lambda = \{y \in H : y(t) \neq 0, \forall t\}$ .

We denote by  $\|u\|_1^2 = \int |\dot{u}|^2 + \int |u|^2$  <sup>(4)</sup> the norm in  $H$ .

Define  $f_T : H \rightarrow \mathbb{R} \cup \{+\infty\}$  by setting

$$f_T(u) = \frac{1}{2} \int |\dot{u}(t)|^2 dt - T^2 \int V(u(t)) dt.$$

<sup>(4)</sup> From now on we will assume that each integral is taken from 0 to 1.

[where  $V(0) = -\infty$ ].

Then  $f_T \in C^1(\Lambda; \mathbb{R})$  and, if  $u \in \Lambda$  and  $f'_T(u) = 0$  then  $y(t) = u(t/T)$  is a non-collision solution of  $(P_T)$ .

**THEOREM 1.** — *Suppose  $V \in C^1(\mathbb{R}^N \setminus \{0\}; \mathbb{R})$  satisfies (V1), (V2) and (V3). Then  $\exists T^*$  such that  $\forall T \geq T^*$  problem (P) has at least one non-collision solution  $x$  such that  $\{x(t)\} \notin \Gamma$ .*

## 2. ESTIMATES FOR THE MINIMUM OF $f_T$ ON COLLISION ORBITS

It is easy to show that it exists a function  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}$  of class  $C^1$  such that: (i)  $\psi(s) \rightarrow -\infty$  as  $s \rightarrow 0^+$ ; (ii)  $\max \psi = b$ ; (iii)  $V(x) \leq \psi(|x|)$ ,  $\forall x \in \mathbb{R}^N \setminus \{0\}$ ; (iv)  $\psi$  is not decreasing. Let  $g_T: H^1(0, 1; \mathbb{R}^+) \rightarrow \mathbb{R}$  be defined by

$$g_T(r) = \frac{1}{2} \int |r'(t)|^2 dt - T^2 \int \psi(r(t)) dt.$$

Consider now  $u \in H$ . Setting  $r(t) = |u(t)|$ , one has  $r \in H^1(0, 1; \mathbb{R}^+)$  and

$$\int |r'|^2 = \int |\langle \dot{u}, u/|u| \rangle|^2 \leq \int |\dot{u}|^2.$$

Then

$$f_T(u) \geq \frac{1}{2} \int |r'|^2 - T^2 \int V(u) \geq \frac{1}{2} \int |r'|^2 - T^2 \int \psi(r) = g_T(r).$$

Moreover, if  $u \in H \setminus \Lambda$  there exists a  $\theta \in [0, 1[$  such that  $|u(t+\theta)| \in H_0^1(0, 1; \mathbb{R}^+)$ . Hence

**LEMMA 2.** —  $m_T = \inf \{ f_T(u) : u \in H \setminus \Lambda \} \geq \inf \{ g_T(r) : r \in H_0^1(0, 1; \mathbb{R}^+) \}$ .  $\square$

**LEMMA 3.** —  $g_T$  attains its minimum on  $H_0^1(0, 1; \mathbb{R}^+)$ .

*Proof.* — Trivial since  $g_T$  is coercive and weakly lower semi continuous.  $\square$

LEMMA 4. —  $\exists c, \tau > 0$  such that  $\forall T > \tau$

$$\min \{ g_T(r) : r \in H_0^1(0, 1; \mathbb{R}^+) \} \geq cT - bT^2.$$

*Proof.* — Let  $r_T$  be such that  $g_T(r_T) = \min \{ g_T(r) : r \in H_0^1(0, 1; \mathbb{R}^+) \}$ . Then set

$$T^2 E_T \equiv \frac{1}{2} |r'_T(t)|^2 + T^2 \psi(r_T(t)).$$

From the conservation of energy it follows that  $T^2 E_T$  is a constant of the motion. Fix now  $T_0$  and correspondingly  $r_0 = r_{T_0}$  and  $E_0 = E_{T_0}$ . We claim that  $E_0 < b$ . In fact, since  $\exists t_0$  such that  $r'(t_0) = 0$  [we recall that  $r_0(0) = r_0(T) = 0$ ], then  $T_0^2 E_0 = T_0^2 \psi(r_0(t_0)) \leq T_0^2 b$ , hence  $E_0 \leq b$ . If  $E_0 = b$ , then  $\psi'(r_0(t_0)) = 0$  and  $r_0(t) = r_0(t_0) > 0, \forall t$ , contradiction which proves the claim.

Take now any  $T > T_0$ . Distinguish between: (i)  $E_T \leq E_0$  and (ii)  $E_T > E_0$ . If (i) holds, from  $T^2 \psi(r_T) \leq T^2 E_T$  and  $E_T \leq E_0$  it follows that

$$g_T(r_T) \geq -T^2 \int \psi(r_T) \geq -T^2 E_0. \quad (2)$$

Suppose now that (ii) holds. Let  $t_T$  be such that  $r'_T(t_T) = 0$  and  $r'_T(t) > 0 \forall t \in [0, t_T[$ . Set  $\rho_T = r_T(t_T)$ . From  $E_T > E_0$  and the monotonicity of  $\psi$  it follows that  $\rho_T \geq \rho_0 \equiv r_0(t_0)$ . Since  $r'_T(t) > 0, \forall t \in [0, t_T[$ , we can solve  $r_T(t) = \rho$  in  $[0, t_T[$  to get  $t = \tau_T(\rho)$  such that  $r_T(\tau_T(\rho)) = \rho, \forall \rho \in [0, \rho_T[$ . From the conservation of energy we get, since  $E_T > E_0$

$$(2T^2)^{-1} r'_T(\tau_T(\rho))^2 > (2T_0^2)^{-1} r'_0(\tau_0(\rho))^2, \quad \forall \rho \in [0, \rho_0[.$$

We can now evaluate

$$\begin{aligned} & \frac{1}{2} \int_0^1 |r'_T(t)|^2 dt \geq \frac{1}{2} \int_0^{t_T} |r'_T(t)|^2 dt \\ & = \frac{1}{2} \int_0^{\rho_T} r'_T(\tau_T(\rho)) d\rho \geq \frac{1}{2} \frac{T}{T_0} \int_0^{\rho_0} r'_0(\tau_0(\rho)) d\rho \end{aligned} \quad (3)$$

Set

$$c = \frac{1}{2T_0} \int_0^{\rho_0} r'_0(\tau_0(\rho)) d\rho.$$

Then  $c > 0$  and from (3) and  $\psi(r) \leq b$  it follows

$$g_T(r_T) \geq cT - bT^2, \quad \forall T > T_0 \quad (4)$$

(2) and  $E_0 < b$  jointly with (4) prove the Lemma.  $\square$

### 3. PROOF OF THE THEOREM

We start by showing

LEMMA 5. —  $\forall \varepsilon > 0$ ,  $f_T$  satisfies the PS condition in the set

$$\{x \in \Lambda: f_T(u) \leq \alpha_T - \varepsilon\},$$

where  $\alpha_T = \min\{m_T, -\beta T^2\}$ .

*Proof.* — Let  $(u_n) \subset \Lambda$  be such that

$$f_T(u_n) \leq \alpha_T - \varepsilon, \quad f'_T(u_n) \rightarrow 0.$$

Then  $\frac{1}{2} \int |u'_n|^2 \leq \text{Const.}$ , hence, setting  $w_n = u_n - \int u_n$ ,  $w_n \rightarrow w$  in  $C^0(S^1; \mathbb{R}^N)$ .

Suppose, by contradiction, that  $\xi_n = \int u_n \rightarrow +\infty$ . Then  $|u_n(t)| \rightarrow +\infty$  uniformly and using (V3)

$$f_T(u_n) \geq -\beta T^2 - \varepsilon/2 \quad \text{for } n \text{ sufficiently large,}$$

contradiction which proves the boundedness of  $\|u_n\|_1$ . We immediately deduce that  $u_n \rightarrow u$  strongly in  $C^0(S^1; \mathbb{R}^N)$  and weakly in  $H^1(S^1; \mathbb{R}^N)$ . Moreover, from the weakly lower semi-continuity of  $f_T$ , we deduce

$$f_T(u) \leq \liminf_{n \rightarrow +\infty} f_T(u_n) \leq \alpha_T - \varepsilon < m_T,$$

hence  $u \in \Lambda$ . Usual arguments then prove that  $u_n \rightarrow u$  in  $H^1$ .  $\square$

*Proof of Theorem 1.* — Consider the set of functions  $\Sigma = \{x \in H \text{ such that } x(t) = \xi \cos(2\pi t) + \eta \sin(2\pi t) + x_0,$

$$\xi, \eta, x_0 \in \mathbb{R}^N, |\xi| = |\eta| \leq 1,$$

$$\langle \xi, \eta \rangle = \langle \xi, x_0 \rangle = \langle \eta, x_0 \rangle = 0, |x_0|^2 = 1 - |\xi|^2 - |\eta|^2\}.$$

Then,  $\forall x \in \Sigma, |x(t)| = 1, \forall t$ .

If  $\Phi: \mathbb{S}^{N-1} \rightarrow \Gamma$  is the radial projection [which is a diffeomorphism by (V2)], set

$$\Sigma' = \{ \Phi(x) \text{ such that } x \in \Sigma \}.$$

We have that:

$$\begin{aligned} f_T(u) &= \frac{1}{2} \int |\Phi'(u(t))|^2 |\dot{u}(t)|^2 - T^2 b \\ &\leq c_1 \int |\dot{u}(t)|^2 - T^2 b \quad (c_1 = \sup_{x \in \mathbb{S}^{N-1}} |\Phi'(x)|^2) \\ &\leq 4\pi^2 c_1 - T^2 b. \end{aligned} \quad (5)$$

By Lemmas 2, 3 and 4

$$m_T \geq cT - bT^2, \quad \forall T > \tau. \quad (6)$$

From (5) and (6) it follows:  $\exists \tau_1$  such that

$$f_T(\Sigma') < m_T, \quad \forall T \geq \tau_1. \quad (7)$$

Moreover, since  $\beta < b$ , one has

$$f_T(\Sigma') < -\beta T^2,$$

and hence

$$f_T(\Sigma') < \alpha_T = \min \{ m_T, -\beta T^2 \}. \quad (8)$$

One can now proceed as in the proof of the theorem by Lyusternik and Fet on the existence of one closed geodesic on a compact Riemannian manifold (see [9], Theorem A.1.5). In fact, letting  $\varepsilon > 0$  be such that  $f_T(\Sigma') < \alpha_T - \varepsilon$ , we can work in the set  $\{ u: f_T(u) \leq \alpha_T - \varepsilon \}$ , where the PS condition holds according to Lemma 5. Since the minimum on such a set is achieved on  $\{ x \in \mathbb{R}^N: V(x) = b \}$ , set which is homeomorphic (through  $\Phi$ ) to  $\mathbb{S}^{N-1}$ , the existence of a critical point  $u$  such that  $-T^2 b < f_T(u) < \alpha_T$  follows. Lastly, if such a critical point is such that  $u(t) \in \Gamma, \forall t$ , then for the corresponding solution  $y(t) = u(t/T)$  one would find  $y(t) = y_0$ . Hence one finds  $f_T(u) = -T^2 b$ , a contradiction. This completes the proof.  $\square$

## 4. FINAL REMARKS

PROPOSITION 6. — Let  $N=2$  and  $V \in C^1(\mathbb{R}^N \setminus \{0\}, \mathbb{R})$  satisfy (V1-2). Then

(i)  $\exists T^*$  such that  $\forall T > T^*$ ,  $(P_T)$  has a non constant solution  $y(t)$  which is a non-collision orbit;

(ii) if  $\Omega$  is convex, then  $y(t) \in \Omega, \forall t$ .

*Proof.* — The only point where (V3) has been used is in proving Lemma 5. If  $N=2$ , this can be avoided using Lemmas 2-4 jointly with the arguments of [6]. We will be sketchy here. Let  $\Lambda_0 = \{u \in \Lambda: u \text{ is non-contractible to a constant in } \Lambda\}$ . It is possible to show that  $f_T$  is (bounded from below on  $H$  and) coercive on  $\Lambda_0$ . Since  $\Sigma \subset \Lambda_0$ , (7) implies that  $\inf \{f_T(u): u \in \Lambda_0\} < m_T$  for  $T$  large. Then it follows that  $\exists u_0 \in \Lambda_0: f_T(u_0) = \min \{f_T(u): u \in \Lambda_0\}$ . This proves (i).

As for (ii), consider

$$U(x) = \begin{cases} V(x), & \forall x \in \Omega \\ b, & \forall x \in \mathbb{R}^2 \setminus \Omega. \end{cases}$$

$U$  is of class  $C^1$ . Applying (i) above we find a  $T$ -periodic solution of

$$-\ddot{y} = U'(y)$$

with  $y \in \Lambda_0$ . It follows easily that such a solution must be contained in  $\Omega$  for every  $t$  (in fact, if it hits the boundary it must be a straight line in the past or in the future, so that it cannot be periodic).  $\square$

Remark 7. — By a suitable modification of Lemma 5, it would be possible to show that Proposition 6 (ii) holds even if  $N > 2$ .

PROPOSITION 8. — Let the assumptions of Theorem 5 be satisfied. Then for every compact set  $K \subset \Omega$ ,  $\exists T_0: \forall T \geq T_0$ ,  $(P_T)$  has a solution  $y_T$  with  $\{y_T(t)\} \not\subset K$ .

*Proof.* — From the proof of Theorem 5 we know that for  $T$  large enough  $f_T$  has a critical point  $u_T$  such that

$$f_T(u_T) < 4\pi^2 c_1 - bT^2. \quad (9)$$

If there is a compact set  $K \subset \Omega$  such that  $u_T(t) \in K, \forall t$ , one would have

$$f_T(u_T) \geq -T^2 \int V(u_T) \geq -T^2 \max_K V.$$

Since  $\max_k V < b$ , this is in contradiction with (9) for  $T$  large.  $\square$

## ACKNOWLEDGEMENTS

Part of this work was done while the Authors were visiting the C.R.M., Université de Montreal. They want to thank for the kind hospitality.

## REFERENCES

- [0] A. AMBROSETTI and V. COTI ZELATI, Solutions périodiques sans collision pour une classe de potentiels de type Keplerien, *C.R. Acad. Sci. Paris*, **305**, 1987, pp. 813-815.
- [1] A. AMBROSETTI and V. COTI ZELATI, Critical Points with Lack of Compactness and Singular Dynamical Systems, *Annali di Matematica Pura ed Applicata* (to appear).
- [2] A. AMBROSETTI and V. COTI ZELATI, Periodic Solutions of Singular Dynamical Systems, *Proceed. NATO ARW "Periodic Solutions of Hamiltonian Systems and Related Topics"*, Il Ciocco, Italy, 1986 (to appear).
- [3] A. CAPOZZI, C. GRECO and A. SALVATORE, *Lagrangian Systems in Presence of Singularities*, preprint Università di Bari, 1985.
- [4] C. COTI ZELATI, Remarks on Dynamical Systems with Weak-Forces, *Manuscripta Math.*, Vol. **57**, 1987, pp. 417-424.
- [5] M. DEGIOVANNI, F. GIANNONI and A. MARINO, Dynamical Systems with Newtonian Potentials, *Proceed. NATO ARW "Periodic Solutions of Hamiltonian Systems and Related Topics"*, Il Ciocco, Italy, 1986 (to appear).
- [6] W. GORDON, Conservative Dynamical Systems Involving Strong Forces, *Trans. Am. Math. Soc.*, Vol. **204**, 1975, pp. 113-135.
- [7] W. GORDON, A Minimizing Property of Keplerian Orbits, *Am. Journ. of Math.*, Vol. **99**, 1977, pp. 961-971.
- [8] C. GRECO, *Periodic Solutions of a Class of Singular Hamiltonian Systems*, preprint, Università di Bari, 1986.
- [9] W. KLINGENBERG, *Lectures on Closed Geodesics*, Springer-Verlag, Berlin, 1978.

(Manuscrit reçu le 7 juillet 1987.)