

ANNALES DE L'I. H. P., SECTION C

A. AMBROSETTI

V. COTI ZELATI

Addendum to closed orbits of fixed energy for a class of N-body problems

Annales de l'I. H. P., section C, tome 9, n° 3 (1992), p. 337-338

http://www.numdam.org/item?id=AIHPC_1992__9_3_337_0

© Gauthier-Villars, 1992, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Addendum to
Closed orbits of fixed energy for a class of N-body
problems

by

A. AMBROSETTI

Scuola Normale Superiore, Pisa

and

V. COTI ZELATI

University of Napoli, Faculty of Architecture, Napoli

In Lemma of 6 the paper *Closed Orbits of fixed energy for a class of N-body problems*, published on this Journal (Vol. 9, No. 2, 1992, pp. 187-200) it is claimed that critical points of f_ε are found by applying the Mountain-Pass Theorem. Actually, such a Theorem needs to be slightly changed, in order to make it applicable in Lemma 6. We indicate below the required modification.

Let $\Sigma_\rho = \{u \in \Lambda_0 : \|u\| \geq \rho\}$, $\Gamma_\rho = \{p \in C([0, 1], \Sigma_\rho) : \|p(0)\| = \rho, p(1) = u_1\}$, and

$$c = \inf_{p \in \Gamma_\rho} \max_{0 \leq \xi \leq 1} f_\varepsilon(p(\xi)).$$

Hereafter, $0 < \varepsilon \leq \varepsilon_0$. By Lemma 2 (i), $a \geq \beta > 0$. Suppose c is not a critical level for f_ε . Since (PS^+) holds, then there exist $m > 0$ and $k \in]0, \beta/2[$ such that $\|f'_\varepsilon\| \geq m$, for all $u \in \Lambda_0 \cap \{c - k \leq f_\varepsilon(u) \leq c + k\}$. Let $\eta(s, u)$, $\eta : [0, \tau_u[\times \Lambda_0 \rightarrow E$ denote the steepest descent flow satisfying

$$\frac{d\eta}{ds} = -X(\eta), \quad \eta(0, u) = u,$$

where X is, as usual, a pseudo-gradient vector field for f_ε , such that (i) $X(u) = 0$ if $f_\varepsilon(u) \leq c - 2k$ or $f_\varepsilon(u) \geq c + 2k$, (ii) $X(u) = f'_\varepsilon$ if $c - k \leq f_\varepsilon(u) \leq c + k$, and (iii) $f_c(\eta(s, u)) \leq f_\varepsilon(u)$, $\forall 0 \leq s < \tau_u$ (see the Deformation lemma in [2]).

Since, as a consequence of (2.3), $f_\varepsilon(u) \rightarrow +\infty$ whenever $u \rightarrow v \in \partial\Lambda_0 - \{0\}$, then (iii) above readily implies that $\eta(s, u) \in \Lambda_0$ whenever $u \in \Lambda_0$ and $\eta(s, u) \neq 0$.

Let us show that $\tau_u \geq \rho/2$, for all $u \in \Sigma_\rho$. According to the preceding remark, this follows in the usual way if $\|\eta(s, u)\| > \rho/2$, $\forall s < \tau_u$ (indeed, in such a case $\tau_u = +\infty$). Otherwise, let $S \in]0, \tau_u[$ be such that $\|\eta(S, u)\| = \rho/2$, for some $u \in \Sigma_\rho$. Then

$$\frac{\rho}{2} \leq \|\eta(S, u) - u\| \leq \int_0^S \|X(\eta)\| \leq S,$$

proving the claim.

Note also that the same arguments used to prove Lemma 2 (i) yield that $(f'_\varepsilon(u) | u) > 0$, $\forall u \in \Lambda_0$, $\|u\| = \rho$. Therefore $\|\eta(s, u)\| < \rho$, whenever $s \geq 0$ and $u \in \Lambda_0$, $\|u\| = \rho$.

After these preliminaries, let $k' < \min(k, \rho/4m)$ and let $p \in \Gamma_\rho$ be such that $\max_\xi f_\varepsilon(p(\xi)) < c + k'$. Consider the path $p_1(\xi) = \eta\left(\frac{\rho}{2}, p(\xi)\right)$. Using the properties of the vector field X , one shows in the standard way that $\max_\xi f_\varepsilon(p_1(\xi)) < c - k'$. Moreover, $p_1(1) = u_1$, whereas, as noted before, $\|p_1(0)\| < \rho$. Let $\xi_0 = \min\{\chi \in [0, 1] : \|p_1(\xi)\| > \rho, \forall \xi \in]\chi, 1]\}$. Setting $q(\xi) = p_1(\xi_0 + \xi(1 - \xi_0))$, it follows that $q \in \Gamma_\rho$ as well as $\max_\xi f_\varepsilon(q(\xi)) < c - k'$, a contradiction which shows that c is a critical level for f_ε .