

ANNALES DE L'I. H. P., SECTION C

J.-P. DIAS

M. FIGUEIRA

The Cauchy problem for a nonlinear Wheeler-DeWitt equation

Annales de l'I. H. P., section C, tome 10, n° 1 (1993), p. 99-107

http://www.numdam.org/item?id=AIHPC_1993__10_1_99_0

© Gauthier-Villars, 1993, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

The Cauchy problem for a nonlinear Wheeler-DeWitt equation

by

J.-P. DIAS and M. FIGUEIRA

CMAF, Av. Prof. Gama Pinto, 2
1699 Lisboa Codex, Portugal

ABSTRACT. — In this paper we consider a nonlinear version of the simplified Wheeler-DeWitt equation which describes the minisuperspace model for the wave function ψ of a closed universe (*cf.* [3], [2], [1]). Following [1], where the linear case has been solved, we study this equation as an evolution equation in the scalar field $y \in \mathbf{R}$ with a scale factor $x \in]0, \mathbf{R}[$. We solve the Cauchy problem for the initial data $\psi(x, 0)$ and $\frac{\partial \psi}{\partial y}(x, 0)$ and we study some decay properties and blow-up situations. A particular nonlinear version has been proposed in [6].

RÉSUMÉ. — Dans cet article nous considérons une version non linéaire de l'équation de Wheeler-DeWitt simplifiée qui décrit le modèle de minisuperspace pour la fonction d'onde ψ d'un univers fermé (*cf.* [3], [2], [1]). Dans l'esprit de [1], où nous avons résolu le cas linéaire, nous étudions cette équation comme une équation d'évolution dans le champ scalaire $y \in \mathbf{R}$ avec le facteur d'échelle $x \in]0, \mathbf{R}[$. Nous résolvons le problème de Cauchy pour des données initiales $\psi(x, 0)$ et $\frac{\partial \psi}{\partial y}(x, 0)$ et nous étudions des propriétés de décroissance à l'infini et des situations d'explosion de la solution. Une version non linéaire particulière a été proposée dans [6].

Research partially supported by JNICT under contract n° PMCT/C/CEN/18/90.
Classification A.M.S. : 35 L 70.

1. INTRODUCTION

We consider the following nonlinear model for the simplified Wheeler-DeWitt equation:

$$\frac{1}{x^2} \frac{\partial^2 \psi}{\partial y^2} - \frac{\partial^2 \psi}{\partial x^2} - \frac{p}{x} \frac{\partial \psi}{\partial x} + x^2 \psi - k^2 x^4 \psi + g x^{q-2} |\psi|^r \psi = 0 \quad (1.1)$$

where $y \in \mathbf{R}$ is the scalar field, $x \in]0, \mathbf{R}[$ ($\mathbf{R} > 0$) is a scale factor, $p \in \mathbf{R}$, $k^2 > 0$, $g \in \mathbf{R}$, $r \geq 1$ and $q \geq \frac{1}{2} r p$ are given constants (p reflects the factor-ordering ambiguity and k^2 is a cosmological constant) and $\psi:]0, \mathbf{R}[\times \mathbf{R} \rightarrow \mathbf{C}$ is the wave function of the universe for the minisuper-space model. The equation (1.1) is equivalent, in the sense of distributions, to the following one:

$$\frac{\partial^2 \psi}{\partial y^2} - x^2 \frac{\partial^2 \psi}{\partial x^2} - p x \frac{\partial \psi}{\partial x} + x^4 \psi - k^2 x^6 \psi + g x^q |\psi|^r \psi = 0 \quad (1.2)$$

which can be considered as an evolution equation in $y \in \mathbf{R}$.

Assuming that $\psi(x, 0)$ and $\frac{\partial \psi}{\partial y}(x, 0)$, belong to some suitable weighted Sobolev spaces (*cf.* [1]) we first prove the existence of a unique local solution for the correspondent Cauchy problem. Under the hypothesis

$$g > 0 \quad \text{and} \quad k^2 \mathbf{R}^2 \leq 1 \quad (\text{that is } x^4 - k^2 x^6 \geq 0), \quad (1.3)$$

we will prove that this solution is a global solution.

Furthermore, if $k^2 \mathbf{R}^2 \leq \frac{2}{3}$ (that is $\frac{d}{dx}(x^4 - k^2 x^6) \geq 0$), we obtain the decay property

$$\int_y^{y+1} \int_0^{\mathbf{R}} x^{q+p-2} |\psi(x, s)|^{r+2} dx ds \rightarrow 0 \quad (y \rightarrow +\infty).$$

Finally we point out that, if $g < 0$ and $k^2 \mathbf{R}^2 \leq 1$, and for special initial data, there exists $y_0 \in \mathbf{R}_+$ such that

$$\int_0^{\mathbf{R}} x^{p-2} |\psi(x, y)|^2 dx \rightarrow +\infty \quad (y \rightarrow y_0^-).$$

We are indebted to O. Bertolami, T. Cazenave, A. B. Henriques and J. M. Mourão for stimulating discussions and useful suggestions.

2. SOME PROPERTIES OF THE ASSOCIATED LINEAR OPERATOR

Following [1] let us denote by $H_{p,0}^1$ the closure of $\mathcal{D}([0, R])$ in the space $\left\{ u \in L_p^2 \mid \frac{du}{dx} \in L_p^2 \right\}$ where $L_p^2([0, R])$ is the L^2 space for the measure $d\mu = x^p dx$. We recall that we have $H_{p,0}^1 \subset L_{p-2}^2$ if $p \neq 1$ and $H_{p,0}^1 \cap L_{p-2}^2 = \{ u \mid x^{p/2} u \in H_0^1 \}$ where $H_0^1 = H_{0,0}^1$ is the usual Sobolev space.

Let us put $H = H_{p,0}^1 \times L_{p-2}^2$,

$$B\psi = x^2 \frac{\partial^2 \psi}{\partial x^2} + px \frac{\partial \psi}{\partial x}, \quad V\psi = -x^4 \psi + k^2 x^6 \psi$$

$$D(A) = D(B) \times (H_{p,0}^1 \cap L_{p-2}^2) \quad \text{with} \quad D(B) = \{ u \in H_{p,0}^1 \mid Bu \in L_{p-2}^2 \}.$$

Now, with $\psi_1 = \frac{\partial \psi}{\partial y}$, the linear equation associated to (1.2) can be written as follows

$$\frac{\partial}{\partial y} \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ B & 0 \end{pmatrix} \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ V & 0 \end{pmatrix} \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} \tag{2.1}$$

In [1] we proved that $A : D(A) \rightarrow H$ defined by

$$A \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ B & 0 \end{pmatrix} \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix}$$

is skew-self-adjoint in H and that $D : H \rightarrow H$ defined by

$$D \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ V & 0 \end{pmatrix} \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix}$$

is continuous. Furthermore

$$D(B) \cap L_{p-2}^2 = \{ u \in H_{p,0}^1 \cap L_{p-2}^2 \mid x^{p/2+1} u \in H^2 \} \\ = \{ u \mid x^{p/2} u \in H_0^1 \text{ and } x^{p/2+1} u \in H^2 \} \quad (\text{cf. [1]}).$$

We denote by $S(y)$, $y \in \mathbf{R}$, the strongly continuous group of operators in H with infinitesimal generator $T = A + D : D(A) \rightarrow H$.

We need the following

PROPOSITION 2.1. — Assume $p = 1$ (that is $H = H_{1,0}^1 \times L_{-1}^2$) and put

$$H_1 = (H_{1,0}^1 \cap L_{-1}^2) \times L_{-1}^2 \subset H, \quad S_1(y)_{|H_1}, \quad T_1 = T_{|D(T) \cap H_1}.$$

Then $S_1(y)$, $y \in \mathbf{R}$, is a strongly continuous group of operators in H_1 with infinitesimal generator T_1 .

Proof. — Let $(u_0, v_0) \in H_1 \cap D(T)$, $(u(y), v(y)) = S(y)(u_0, v_0)$.

Since $\frac{\partial}{\partial y}|u|^2 = 2 \Re e u \bar{v}$ we obtain, for $\varepsilon > 0$,

$$\begin{aligned} \int_0^{\mathbf{R}} x^{-1+\varepsilon} |u(y)|^2 dx &\leq \int_0^{\mathbf{R}} x^{-1+\varepsilon} |u_0|^2 dx + 2 \int_0^y \int_0^{\mathbf{R}} \frac{|u(s)||v(s)|}{x^{(1/2)-(\varepsilon/2)} x^{(1/2)-(\varepsilon/2)}} dx ds \\ &\leq \mathbf{R}^\varepsilon \int_0^{\mathbf{R}} x^{-1} |u_0|^2 dx + 2 \mathbf{R}^{\varepsilon/2} \int_0^y \|u(s)\|_{L^2_{-1+\varepsilon}} \|v(s)\|_{L^2_{-1}} ds. \end{aligned}$$

Hence, by applying Gronwall's inequality and Fatou's lemma ($\varepsilon \rightarrow 0$), we get $u(y) \in L^2_{-1}$ and

$$\|u(y)\|_{L^2_{-1}} \leq \|u_0\|_{L^2_{-1}} + \int_0^y \|v(s)\|_{L^2_{-1}} ds \quad (y \geq 0)$$

and this inequality can be extended, by density, to $(u_0, v_0) \in H_1$. Since for $(u_0, v_0) \in H_1$ we have

$$\|v(s)\|_{L^2_{-1}} \leq \|S(s)(u_0, v_0)\|_{\mathbf{H}} \leq M e^{\omega s} \|(u_0, v_0)\|_{\mathbf{H}}$$

with $M > 0$, $\omega \geq 0$. Hence for $(u_0, v_0) \in H_1$ we get

$$\|u(y)\|_{L^2_{-1}} \leq c(y) (\|u_0\|_{L^2_{-1}} + \|u_0\|_{H^1_{1,0}} + \|v_0\|_{L^2_{-1}}).$$

We also have $\|u(y) - u_0\|_{L^2_{-1}} \leq \int_0^y \|v(s)\|_{L^2_{-1}} ds$.

Now, let \tilde{T} be the infinitesimal generator of S_1 . It is easy to see that $D(\tilde{T}) \subset D(T) \cap H_1$ and that, in $D(\tilde{T})$, $\tilde{T} = T_1$. Let $(u_0, v_0) \in D(T) \cap H_1$.

We have $\frac{1}{y}(u(y) - u_0) \rightarrow v_0$ ($y \rightarrow 0^+$) in $H^1_{1,0}$.

We want to prove that $\frac{1}{y}(u(y) - u_0) \rightarrow v_0$ ($y \rightarrow 0^+$) in L^2_{-1} .

If $y_n \rightarrow 0^+$, we obtain

$$\limsup_{n \rightarrow \infty} \left\| \frac{1}{y_n}(u(y_n) - u_0) \right\|_{L^2_{-1}} \leq \lim_{n \rightarrow \infty} \frac{1}{y_n} \int_0^{y_n} \|v(s)\|_{L^2_{-1}} ds = \|v_0\|_{L^2_{-1}}.$$

Therefore there exists a subsequence y_{n_k} such that

$$\frac{1}{y_{n_k}}(u(y_{n_k}) - u_0) \rightharpoonup \eta \in L^2_{-1} \quad (k \rightarrow \infty), \quad \text{weakly in } L^2_{-1}.$$

But, for $\varepsilon > 0$, $L^2_{-1} \subset L^2_{-1+\varepsilon}$, $H^1_{1,0} \subset H^1_{1+\varepsilon,0} \subset L^2_{-1+\varepsilon}$, and so $\eta = v_0$.

Hence $\frac{1}{y_{n_k}}(u(y_{n_k}) - u_0) \rightarrow v_0$ ($k \rightarrow \infty$) in L^2_{-1} .

We conclude that $D(T) \cap H_1 \subset D(\tilde{T})$. \square

3. THE CAUCHY PROBLEM FOR THE NONLINEAR EQUATION

In the remaining of the paper we put

$$X = \begin{cases} H = H_{p,0}^1 \times L_{p-2}^2, & \text{if } p \neq 1 \\ H_1 = (H_{1,0}^1 \cap L_{-1}^2) \times L_{-1}^2, & \text{if } p = 1 \end{cases} \quad (3.1)$$

$$U = \begin{cases} T = A + D, \text{ with domain } D(A), & \text{if } p \neq 1 \\ T_1 = T_{|H_1}, & \text{if } p = 1 \end{cases} \quad (3.2)$$

and we will denote by $G(y)$, $y \in \mathbf{R}$, the strongly continuous group of operators generated by U in X .

Let

$$F \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} = \begin{pmatrix} 0 \\ -gx^q |\psi|^r \psi \end{pmatrix}, \quad \psi_1 = \frac{\partial \psi}{\partial y} \quad (3.3)$$

The nonlinear equation (1.2) can be written as follows

$$\frac{\partial}{\partial y} \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} = U \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} + F \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} \quad (3.4)$$

and, since $H_{p,0}^1 \cap L_{p-2}^2 = \{u \mid x^{p/2} u \in H_0^1\}$ and $H_0^1 \hookrightarrow L^\infty$, it is easy to prove that if $r \geq 1$ and $q \geq r \frac{p}{2}$, then F is a locally Lipschitz continuous function from X to X .

Furthermore X is an Hilbert space. Hence, by theorem 1.6 in [4] ch. 6, if we take

$$\begin{pmatrix} \psi_0 \\ \psi_{1,0} \end{pmatrix} \in D(U), \quad \text{that is } \psi_0 \in D(B) \cap L_{p-2}^2, \quad \psi_{1,0} \in H_{1,0}^1 \cap L_{p-2}^2,$$

then there exists a unique local mild solution $\begin{pmatrix} \psi \\ \psi_1 \end{pmatrix}$ of the corresponding Cauchy problem for the equation (3.4) which is a local strong solution, that is

$$\left. \begin{aligned} & \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} \in \mathcal{C}]-\varepsilon, \varepsilon[; D(U) \cap \mathcal{C}^1]-\varepsilon, \varepsilon[; X, \\ & \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} \text{ satisfies (3.4) for } y \in]-\varepsilon, \varepsilon[\quad \text{and} \quad \begin{pmatrix} \psi \\ \psi_1 \end{pmatrix} (0) = \begin{pmatrix} \psi_0 \\ \psi_{1,0} \end{pmatrix}. \end{aligned} \right\} (3.5)$$

In order to obtain a result on the global existence of solution we need the following

LEMMA 3.1. — Assume $\begin{pmatrix} \psi_0 \\ \psi_{1,0} \end{pmatrix} \in D(U)$ and let $\begin{pmatrix} \psi \\ \psi_1 \end{pmatrix}$ be the corresponding local solution for the Cauchy problem. We have

$$E(y) = \frac{1}{2} \int_0^{\mathbf{R}} x^{p-2} \left| \frac{\partial \psi}{\partial y} \right|^2 dx + \frac{1}{2} \int_0^{\mathbf{R}} x^p \left| \frac{\partial \psi}{\partial x} \right|^2 dx + \frac{1}{2} \int_0^{\mathbf{R}} x^{p+2} |\psi|^2 dx - \frac{k^2}{2} \int_0^{\mathbf{R}} x^{p+4} |\psi|^2 dx + \frac{g}{r+2} \int_0^{\mathbf{R}} x^{p+q-2} |\psi|^{r+2} dx \equiv E(0) = E, \quad y \in]-\varepsilon, \varepsilon[. \quad (3.6)$$

Proof. — If we multiply the equation (1.2) by $x^{p-2} \frac{\partial \bar{\psi}}{\partial y}$, take the real part and integrate in x (over $]0, \mathbf{R}[$) we obtain $\frac{d}{dy} E(y) = 0$, since

$$\left(x^p \frac{\partial \psi}{\partial x} \frac{\partial \bar{\psi}}{\partial y} \right)(y) \in \mathbf{W}_0^{1,1}. \quad \square$$

Hence, we have

THEOREM 3.1. — Assume (1.3) and $\begin{pmatrix} \psi_0 \\ \psi_{1,0} \end{pmatrix} \in D(U)$.

Then there exists a unique global solution $\begin{pmatrix} \psi \\ \psi_1 \end{pmatrix}$ of the corresponding Cauchy problem for the equation (1.2) in the sense of (3.5) (for $y \in]-\infty, +\infty[$).

Proof. — By (3.6) we have

$$\int_0^{\mathbf{R}} x^{p-2} |\psi_1|^2 dx + \int_0^{\mathbf{R}} x^p \left| \frac{\partial \psi}{\partial x} \right|^2 dx \leq E.$$

In the special case $p = 1$, this implies, reasoning as in the first part of the proof of proposition 2.1, that

$$\|\psi(y)\|_{L^2_1} \leq \|\psi_0\|_{L^2_1} + \int_0^y \|\psi_1(s)\|_{L^2_1} ds \leq \|\psi_0\|_{L^2_1} + E y (y \geq 0). \quad \square$$

Remark. — The theorem 3.1 can be extended to more general situations, for example, if gx^q is replaced by a function $g(x, y) \geq 0$, g continuous, $\frac{\partial g}{\partial y}$ continuous and such that

$$g(x, y) \leq c(y) x^q, \quad \left| \frac{\partial g}{\partial y}(x, y) \right| \leq \min(c_1(y) x^q, c_2(y) g(x, y)),$$

with $c, c_1 \in L^\infty_{\text{loc}}(\mathbf{R})$, $c_2 \in L^1_{\text{loc}}(\mathbf{R} \setminus \{0\})$, $c, c_1, c_2 \geq 0$. Furthermore, the condition $k^2 \mathbf{R}^2 \leq 1$ can be replaced by $\mathbf{R}^4 - k^2 \mathbf{R}^6 + \frac{1}{4}(p-1)^2 > 0$ if $p \neq 1$.

4. DECAY PROPERTIES AND A BLOW-UP CASE

THEOREM 4.1. — Let us assume $g > 0$ and $k^2 R^2 \leq \frac{2}{3}$ and let

$$\begin{pmatrix} \psi_0 \\ \psi_{1,0} \end{pmatrix} \in D(U) \text{ and}$$

$$\psi \in \mathcal{C}(\mathbf{R}; D(\mathbf{B}) \cap L^2_{p-2}) \cap \mathcal{C}^1(\mathbf{R}; H^1_{p,0} \cap L^2_{p-2}) \cap \mathcal{C}^2(\mathbf{R}; L^2_{p-2})$$

be the unique global solution of the corresponding Cauchy problem for the equation (1.2). Then we have

$$(2 - 3k^2 R^2) \int_0^y \int_0^{\mathbf{R}} x^{p+2} |\psi|^2 dx ds + g \frac{2q - rp + r}{2(r+2)} \int_0^y \int_0^{\mathbf{R}} x^{p+q-2} |\psi|^{r+2} dx ds \leq 4E, \quad (4.1)$$

$\forall y \in \mathbf{R}_+$, where E is defined by (3.6).

Proof. — Let $v = x^{p/2} \psi$. We have $v \in \mathcal{C}^1(\mathbf{R}; H^1_0) \cap \mathcal{C}^2(\mathbf{R}; L^2_{-2})$, $xv \in \mathcal{C}(\mathbf{R}; H^2)$ and the equation (1.2) takes the form

$$\frac{\partial^2 v}{\partial y^2} - x^2 \frac{\partial^2 v}{\partial x^2} + \frac{p}{4}(p-2)v + x^4 v - k^2 x^6 v + gx^{q-(1/2)pr} |v|^r v = 0. \quad (4.2)$$

Let us multiply the equation (4.2) by $\frac{1}{2} \frac{\bar{v}}{x^2} - \frac{1}{x} \frac{\partial \bar{v}}{\partial x}$ take the real part and integrate over $]0, \mathbf{R}[$. Taking in account that

$$\Re \int_0^{\mathbf{R}} \frac{\partial^2 v}{\partial y^2} \frac{\bar{v}}{x^2} dx = \Re \frac{d}{dy} \int_0^{\mathbf{R}} \frac{\partial v}{\partial y} \frac{\bar{v}}{x^2} dx - \int_0^{\mathbf{R}} \frac{1}{x^2} \left| \frac{\partial v}{\partial y} \right|^2 dx$$

and

$$\Re \int_0^{\mathbf{R}} \frac{\partial^2 v}{\partial y^2} \frac{\partial \bar{v}}{\partial x} \frac{1}{x} dx = \Re \frac{d}{dy} \int_0^{\mathbf{R}} \frac{\partial v}{\partial y} \frac{\partial \bar{v}}{\partial x} \frac{1}{x} dx - \frac{1}{2} \int_0^{\mathbf{R}} \frac{1}{x} \frac{\partial}{\partial x} \left| \frac{\partial v}{\partial y} \right|^2 dx$$

and integrating by parts we obtain (recall that if $u \in H^1$ and $u(0) = 0$ then $x^{-(1/2)} u \rightarrow 0 (x \rightarrow 0^+)$):

$$\begin{aligned} 0 &= \frac{1}{2} \frac{d}{dy} \Re \int_0^{\mathbf{R}} \frac{1}{x^2} \frac{\partial v}{\partial y} \bar{v} dx - \frac{d}{dy} \Re \int_0^{\mathbf{R}} \frac{1}{x} \frac{\partial v}{\partial y} \frac{\partial \bar{v}}{\partial x} dx \\ &+ \int_0^{\mathbf{R}} (2x^2 - 3k^2 x^4) |v|^2 dx + g \frac{2q - rp + r}{2(r+q)} \int_0^{\mathbf{R}} x^{q-(1/2)pr-2} |v|^{r+2} dx \\ &+ \frac{1}{2} \mathbf{R} \left| \frac{\partial v}{\partial x} \right|^2(\mathbf{R}) - \frac{1}{2} \lim_{x \rightarrow 0^+} \left(x \left| \frac{\partial v}{\partial x} \right|^2 \right) + \frac{1}{4} \lim_{x \rightarrow 0^+} \frac{\partial |v|^2}{\partial x}. \quad (4.3) \end{aligned}$$

But

$$x \frac{\partial v}{\partial x} \in H^1 \quad \text{and} \quad x \frac{\partial v}{\partial x} = \frac{\partial}{\partial x}(xv) - \frac{xv}{x} \rightarrow 0 (x \rightarrow 0^+).$$

Hence

$$x \left| \frac{\partial v}{\partial x} \right|^2 \rightarrow 0 (x \rightarrow 0^+)$$

and

$$\frac{\partial |v|^2}{\partial x} = 2 \Re e \left[(x^{-1/2} \bar{v}) \left(x^{1/2} \frac{\partial v}{\partial x} \right) \right] \rightarrow 0 (x \rightarrow 0^+)$$

From (4.3) we obtain for $\psi = x^{-(p/2)} v$ and with $k^2 R^2 \leq \frac{2}{3}$:

$$\begin{aligned} \theta &= (2 - 3k^2 R^2) \int_0^R x^{p+2} |\psi|^2 dx + g \frac{2q - rp + r}{2(r+2)} \int_0^R x^{p+q-2} |\psi|^{r+2} dx \\ &\leq -\frac{1}{2} \frac{d}{dy} \Re e \int_0^R x^{p-2} \frac{\partial \psi}{\partial y} \bar{\psi} dx + \frac{d}{dy} \Re e \int_0^R x^{p-1} \frac{\partial \psi}{\partial y} \frac{\partial \bar{\psi}}{\partial x} dx \\ &\quad + \frac{d}{dy} \Re e \frac{p}{2} \int_0^R x^{p-2} \frac{\partial \psi}{\partial x} \bar{\psi} dx. \end{aligned}$$

Hence, for $y > 0$,

$$\begin{aligned} &\int_0^y \theta(s) ds \\ &\leq \left[\int_0^R x^{p-2} \left| \frac{\partial \psi}{\partial y} \right|^2 dx + \frac{1}{2} \int_0^R x^p \left| \frac{\partial \psi}{\partial x} \right|^2 dx + \frac{1}{2} \frac{(p-1)^2}{4} \int_0^R x^{p-2} |\psi|^2 dx \right] (y) \\ &\quad + \left[\int_0^R x^{p-2} |\psi_{1,0}|^2 dx + \frac{1}{2} \int_0^R x^p \left| \frac{\partial \psi_0}{\partial x} \right|^2 dx \right. \\ &\quad \left. + \frac{1}{2} \frac{(p-1)^2}{4} \int_0^R x^{p-2} |\psi_0|^2 dx \right] \leq 4E, \end{aligned}$$

since for $p \neq 1$ we have, by Hardy's inequality (cf. (2.1) in [1]),

$$\frac{(p-1)^2}{4} \int_0^R x^{p-2} |\psi|^2 dx \leq \int_0^R x^p \left| \frac{\partial \psi}{\partial x} \right|^2 dx. \quad \square$$

COROLLARY. — Under the hypothesis of theorem 4.1 we have

$$\int_y^{y+1} \int_0^R x^{p+q-2} |\psi|^{2+r} dx ds \rightarrow 0 (y \rightarrow +\infty).$$

Remark. — By the reversibility in y we have similar results for $y \in \mathbf{R}_-$.

Finally we can point out a blow-up situation. The proof of the following result is similar to the one in the example of X. 13 in [5] for the nonlinear wave equation and so we omit it.

PROPOSITION 4.1. — Assume $g < 0$, $k^2 R^2 \leq 1$ and let $\begin{pmatrix} \psi_0 \\ \psi_{1,0} \end{pmatrix} \in D(U)$ be real and such that $E \leq 0$ and $\int_0^R x^{p-2} \psi_0 \psi_{1,0} dx > 0$. Let ψ be the local solution of the corresponding Cauchy problem for the equation (1.2). Then there exists $y_0 \in \mathbf{R}_+$ such that

$$\lim_{y \rightarrow y_0} \int_0^R x^{p-2} \psi^2(x, y) dx = +\infty.$$

Remark. — Like in the quoted example for the nonlinear wave equation, it is easy to check that $y_0 \leq \frac{2}{r} \left(\int_0^R x^{p-2} \psi_0^2 dx \right) \left(\int_0^R x^{p-2} \psi_0 \psi_{1,0} dx \right)^{-1}$.

REFERENCES

- [1] J. P. DIAS and M. FIGUEIRA, The Simplified Wheeler-DeWitt Equation: the Cauchy Problem and Some Spectral Properties, *Ann. Inst. H. Poincaré*, Physique Théorique, Vol. **54**, 1991, pp. 17-26.
- [2] G. W. GIBBONS and L. P. GRISHCHUK, What is a Typical Wave Function for the Universe?, *Nucl. Phys. B*, Vol. **313**, 1989, pp. 736-748.
- [3] J. B. HARTLE and S. W. HAWKING, Wave Function of the Universe, *Phys. Rev. D*, Vol. **28**, 1983, pp. 2960-2975.
- [4] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer, 1983.
- [5] M. REED and B. SIMON, *Methods of Modern Mathematical Physics*, Vol. II: *Fourier Analysis, Self-Adjointness*, Academic Press, 1975.
- [6] L. SUSSKIND, *Lectures at the Trieste School on String Theories and Quantum Gravity*, April 1990.

(Manuscript received January 25 1991;
revised October 1st 1991.)