

ANNALES DE L'I. H. P., SECTION C

FENG ZHOU

QING ZHOU

A remark on multiplicity of solutions for the Ginzburg-Landau equation

Annales de l'I. H. P., section C, tome 16, n° 2 (1999), p. 255-267

http://www.numdam.org/item?id=AIHPC_1999__16_2_255_0

© Gauthier-Villars, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A remark on multiplicity of solutions for the Ginzburg-Landau equation

by

Feng ZHOU and Qing ZHOU

Department of Mathematics,
East China Normal University, Shanghai 200062, China
e-mail: fzhov@euler.math.ecnu.edu.cn
qzhou@euler.math.ecnu.edu.cn

ABSTRACT. – In this paper we study the structure of certain level set of the Ginzburg-Landau functional which has similar topology with the configuration space. As an application, we generalize Almeida-Bethuel's result on multiplicity of solutions for the Ginzburg-Landau equation. © Elsevier, Paris

Key words: Ginzburg-Landau equation, Ljusternik-Schnirelman theory, renormalized energy

RÉSUMÉ. – On étudie la structure de certains ensembles de niveau de la fonctionnelle du type Ginzburg-Landau qui ont des topologies similaires à celles de l'espace de configuration. Comme application, on généralise le résultat d'Almeida-Bethuel sur la multiplicité des solutions des équations de G-L. © Elsevier, Paris

1. INTRODUCTION

Let $\Omega \subset \mathbb{C}$ be a smooth, bounded and simply connected domain. Let $g : \partial\Omega \rightarrow \mathbb{C}$ be a prescribed smooth map with $|g(x)| = 1$, for all $x \in \partial\Omega$.

1991 Mathematics Subject Classification. 35 J 20.

Both authors are supported partially by NNSF and SEDC of China. The second author is also supported by HYTEF.

The Ginzburg-Landau functional, for any $\varepsilon > 0$, is given by

$$E_\varepsilon(u) = \frac{1}{2} \int_\Omega \left\{ |\nabla u|^2 + \frac{1}{2\varepsilon^2} (1 - |u|^2)^2 \right\} \quad (1.1)$$

which is defined on the Hilbert space

$$H_g^1(\Omega, \mathbb{C}) = \{u \in H^1(\Omega, \mathbb{C}); u = g \text{ on } \partial\Omega\}.$$

It is easy to verify that E_ε is a positive, C^2 -functional satisfying the Palais-Smale condition. So

$$\mu_\varepsilon = \min_{u \in H_g^1(\Omega, \mathbb{C})} E_\varepsilon(u)$$

is achieved by some $u_\varepsilon \in H_g^1(\Omega, \mathbb{C})$ and these minimizers satisfy the following Ginzburg-Landau equation:

$$\begin{cases} -\Delta u = \frac{1}{\varepsilon^2} u(1 - |u|^2) & \text{in } \Omega \\ u = g & \text{on } \partial\Omega. \end{cases} \quad (1.2)$$

The Ginzburg-Landau equation (1.2) has been extensively studied by F. Bethuel, H. Brezis and F. Hélein [BBH1, 2] and many others. A complete characterization of asymptotic behavior (as $\varepsilon \rightarrow 0^+$) for minimizing solutions of (1.2) is given. It has been shown that the degree of g , denoted by $k = \deg(g, \partial\Omega)$, plays a crucial role in the asymptotic analysis of the minimizers. Without loss of generality, we will always assume $k \geq 0$ throughout this paper.

In this paper, we will study the multiplicity of the solutions for the Ginzburg-Landau equation (1.2), many such results have been given for special domains and/or boundary values (see for instance Almeida and Bethuel [AB1], Felmer and Del Pino [FP], F.H. Lin [Li]). The motivation of our paper comes from the recent work of Almeida-Bethuel [AB2, 3] concerning the existence of non-minimizing solutions of (1.2). They showed that if $k \geq 2$, the Ginzburg-Landau equation (1.2) has at least three distinct solutions, among which at least one is not minimizing. Based on topological arguments directly inspired by Almeida-Bethuel's work, we obtain our main result as follows

THEOREM 1. – *Assume that $k \geq 2$, there is some $\varepsilon_0 > 0$ (depending on Ω and g only) such that if $\varepsilon < \varepsilon_0$, the equation (1.2) has at least $k + 1$ distinct solutions.*

To prove Theorem 1, we will apply the standard Ljusternik-Schnirelman theory to a suitable covering space of a level set

$$E_\varepsilon^a = \{u \in H_g^1(\Omega, \mathbb{C}); E_\varepsilon(u) < a\},$$

for an a of the form

$$a = \mu_\varepsilon + \lambda \tag{1.3}$$

where λ is a fixed positive constant to be determined later. The proof is strongly related to the topological similarities between E_ε^a and the configuration space $\Sigma_k(\Omega)$ of k distinct points in Ω . As in [AB3], we need to use a map $\tilde{\Phi}$ from E_ε^a into $\Sigma_k(\Omega)$. More precisely, We may assign to each function u in E_ε^a , a set of k distinct points $\{a_1, \dots, a_k\}$, called the vortices of u , where each vortex has the topological degree $+1$. The map $\tilde{\Phi} : E_\varepsilon^a \rightarrow \Sigma_k(\Omega)$ is not continuous. However this difficulty can be overcome by applying the notion of η -almost continuity given in [AB3]. The topological similarity between E_ε^a and $\Sigma_k(\Omega)$ allows us to define a covering space \tilde{E}_ε^a of E_ε^a corresponding to the covering $F_k(\Omega) \rightarrow \Sigma_k(\Omega)$, where $F_k(\Omega)$ is the configuration space of ordered k distinct points in Ω . Again we have topological similarity between these two spaces, and we than can prove that the category of \tilde{E}_ε^a is at least k . The Ljusternik-Schnirelman minimax theorem concludes that the functional \tilde{E}_ε on \tilde{E}_ε^a , which is the composition of E_ε and the covering projection either has at least k distinct critical values or the dimension of the critical set is at least 1. These imply that E_ε has at least k critical points on E_ε^a . Finally, the fact that $E_\varepsilon^\infty = H_g^1(\Omega, \mathbb{C})$ is an affine space guarantee that E_ε^a has at least another critical point outside of E_ε^a , if $k \geq 2$.

This paper is organized as follows: In the next section we will recall some preliminary results about the configuration space and the construction of the map $\tilde{\Phi}$ in [AB3] and Theorem 1 will be proved in Section 3.

2. PRELIMINARIES

Our proof of Theorem 1 relies essentially on the properties of the map $\tilde{\Phi} : E_\varepsilon^a \rightarrow \Sigma_k(\Omega)$ described by Almeida and Bethuel [AB3]. With a such map, they showed that the fundamental group $\pi_1(E_\varepsilon^a)$ is non trivial for some suitable value a of the form (1.3) when ε is sufficiently small. We review here some basic facts about the configuration space and the construction of the map $\tilde{\Phi}$.

We study the configuration space and renormalized energy first. Let the metric on \mathbb{C}^k be defined by the following norm

$$\|(z_1, \dots, z_k)\| = \sum_{i=1}^k |z_i|. \quad (2.1)$$

The configuration space of the ordered k distinct points in Ω

$$F_k(\Omega) = \{(a_1, \dots, a_k) \in \Omega^k; a_i \neq a_j \text{ for all } i \neq j\} \subset \mathbb{C}^k$$

with the inherited metric (2.1) on \mathbb{C}^k is a smooth manifold. The cohomology ring $H^*(F_k(\Omega)) = H^*(F_k(\Omega), \mathbb{R})$ of the space $F_k(\Omega)$ has been determined by Arnol'd in 1969 (see [Ar]), which is generated by elements $\omega_{ij} \in H^1(F_k(\Omega))$, $1 \leq i < j \leq k$ and subject to the following defining relations

$$\omega_{ij}\omega_{jl} + \omega_{jl}\omega_{il} + \omega_{il}\omega_{ij} = 0.$$

Arnol'd also showed that the p th Betti number B_p of $F_k(\Omega)$ is the coefficient of t^p in the polynomial

$$(1+t)(1+2t) \cdots (1+(k-1)t).$$

In particular, $B_{k-1} = (k-1)! \neq 0$, and this concludes that

LEMMA 2. – *The cuplength of $F_k(\Omega)$ is $k-1$.*

The cuplength of a space X is the largest integer n such that there are n elements $\varphi_j \in H^{p_j}(X)$, $p_j > 0$, $1 \leq j \leq n$ and $\varphi_1 \cup \cdots \cup \varphi_n \neq 0$.

The symmetric group S_k on $\{1, \dots, k\}$ acts isometrically on $F_k(\Omega)$ by permuting coordinates, i.e., for all $\sigma \in S_k$,

$$\sigma(a_1, \dots, a_k) = (a_{\sigma(1)}, \dots, a_{\sigma(k)}).$$

This action is free, and the quotient space $F_k(\Omega)/S_k$ is called the configuration space of k distinct point in Ω and it will be denoted by $\Sigma_k(\Omega)$.

On $\Sigma_k(\Omega)$, we have a natural metric such that the quotient map $\pi : F_k(\Omega) \rightarrow \Sigma_k(\Omega)$ is a Riemannian regular covering. This metric on $\Sigma_k(\Omega)$ is the same as the length of minimal connection introduced by Brezis, Coron and Lieb in [BCL], i.e., for $a = \{a_1, \dots, a_k\}$, $a' = \{a'_1, \dots, a'_k\} \in \Sigma_k(\Omega)$,

$$\|a - a'\| = L(a, a') = \inf_{\sigma \in S_k} \sum_{i=1}^k |a_i - a'_{\sigma(i)}|.$$

We now define the renormalized energy W_g on $\Sigma_k(\Omega)$ which is introduced by Bethuel-Brezis-Hélein in [BBH2] as follows, for $a = \{a_1, \dots, a_k\} \in \Sigma_k(\Omega)$,

$$W_g(a_1, \dots, a_k) = -\pi \sum_{i \neq j} \log |a_i - a_j| + \frac{1}{2} \int_{\partial\Omega} \phi \cdot (g \times g_\tau) - \pi \sum_{j=1}^k R(a_j)$$

where ϕ is the solution of

$$\begin{cases} \Delta\phi = 2\pi \sum_{i=1}^k \delta_{a_i} & \text{in } \Omega \\ \frac{\partial\phi}{\partial\nu} = g \times g_\tau & \text{on } \partial\Omega \\ \int_{\partial\Omega} \phi = 0. \end{cases}$$

Here ν denotes the unit outer normal to $\partial\Omega$ and τ is unit tangent to $\partial\Omega$ oriented so that $\nu \times \tau = 1$. And the function R is the regular part of ϕ , i.e.,

$$R(z) = \phi(z) - \sum_{i=1}^k \log |z - a_i|.$$

It is clear that $W_g(a) \rightarrow +\infty$ if $\text{dist}(a_j, \partial\Omega) \rightarrow 0$ for some i or if $|a_i - a_j| \rightarrow 0$ for some $i \neq j$. It has been proved in [BBH2] that, as $\varepsilon \rightarrow 0$, we have

$$\mu_\varepsilon = k\pi |\log \varepsilon| + W_g(a_1^*, \dots, a_k^*) + k\nu_0 + o(1),$$

where $o(1) \rightarrow 0$ as $\varepsilon \rightarrow 0$, ν_0 is a universal constant, and (a_1^*, \dots, a_k^*) is a global minimum of the function W_g .

Next we will turn to the construction of the map $\tilde{\Phi}$. We will use a regularization technique, that is, for any $u \in E_\varepsilon^a$, we can associate a map u^h , which is a minimizer (not necessarily to be unique) of the following minimization problem

$$\inf_{v \in H_g^1(\Omega, \mathbb{C})} \left\{ E_\varepsilon(v) + \int_\Omega \frac{|u - v|^2}{2h^2} \right\} \tag{2.2}$$

where $h = \varepsilon^{\frac{2}{4k+1}} > 0$. We denote $u^h = T(u)$ where $T : H_g^1(\Omega, \mathbb{C}) \rightarrow H_g^1(\Omega, \mathbb{C})$. Clearly we have $u^h \in E_\varepsilon^a$ and it satisfies an equation similar to the Ginzburg-Landau equation (1.2). One of the main observations in [AB3]

is that we can describe the “vortex structure” not only for the solutions of the Ginzburg-Landau equation, but also for such maps u^h . To be more precise, let us collect some of results of [AB3].

THEOREM 3 [AB3]. – *Assume that a is of the form (1.3) for some constant $\lambda > 0$. Then there is a constant $0 < \varepsilon'_0 < 1$ depending only on Ω , g and λ , such that if $\varepsilon < \varepsilon'_0$, then for $u \in E_\varepsilon^a$, $|u| \leq 1$ on Ω , there is a point $a = \{a_1, \dots, a_k\}$ in $\Sigma_k(\Omega)$ such that*

$$|u^h(x)| \geq \frac{1}{2}, \quad \forall x \in \Omega \setminus \bigcup_{i=1}^k B(a_i, \rho)$$

where ρ satisfies $\varepsilon^\chi \leq \rho \leq \varepsilon^{\bar{\chi}}$, for some constants $\chi, \bar{\chi} \in]0, 1[$ independent of ε .

$$\deg(u^h, a_i) = \deg\left(\frac{u^h}{|u^h|}, \partial B(a_i, \rho)\right) = +1, \text{ for all } 1 \leq i \leq k.$$

Moreover, there exists some constant $\beta > 0$ depending only on Ω , g and λ such that $\text{dist}(a_i, \partial\Omega) \geq \beta$, for all $1 \leq i \leq k$ and $|a_i - a_j| \geq \beta$, for all $1 \leq i \neq j \leq k$.

Thus we can see that the properties of maps u^h are very close to that of minimizers of (1.2) as in [BBH], and it allows us to define vortices $\{a_1, \dots, a_k\}$ for u^h and each of the vortices has topological degree +1. That defines a map Ψ from $\text{Im}(T(P(E_\varepsilon^a)))$ to $\Sigma_k(\Omega)$, by $\Psi(u^h) = \{a_1, \dots, a_k\}$, where the map $P : H_g^1(\Omega, \mathbb{C}) \rightarrow H_g^1(\Omega, \mathbb{C})$ defined by

$$\begin{cases} Pu(x) = u(x) & \text{if } |u(x)| \leq 1 \\ Pu(x) = \frac{u(x)}{|u(x)|} & \text{if } |u(x)| \geq 1 \end{cases}$$

is continuous. Composing P, T and Ψ , we define $\tilde{\Phi} : E_\varepsilon^a \rightarrow \Sigma_k(\Omega)$;

$$\tilde{\Phi}(u) = \Psi(T(Pu)).$$

As already noticed in [AB3], the minimizer u^h to the problem (2.2) may not be unique and moving slightly the points a_i 's, the new positions would still match the requirements of Theorem 2. Hence the assignment of u^h and the vortices for u^h require some choices, so we can not expect the map $\tilde{\Phi}$ to be continuous. However the freedom in these choices are not too wild, and we can say that $\tilde{\Phi}$ is “almost” a continuous map from E_ε^a to $\Sigma_k(\Omega)$. More precisely, we have

PROPOSITION 4 [AB3]. – Assume that $a, \varepsilon'_0, \bar{\chi}$ are as in Theorem 3. Then for all $\varepsilon < \varepsilon'_0, u, v \in E_\varepsilon^a$ we have

$$\|\tilde{\Phi}(u) - \tilde{\Phi}(v)\| \leq C_1 \left(|\log \varepsilon| \varepsilon^{\frac{2}{4k+1}} + \varepsilon^{\bar{\chi}} + \|u - v\|_{H_0^1(\Omega, \mathbb{C})} \right)$$

where C_1 is a constant depending only on Ω and g .

Remark. – In [AB3], Almeida-Bethuel studied the more general configuration space corresponding to the “vortices” of the map u^h for $u \in E_\varepsilon^a$, where a is of the form

$$\mu_\varepsilon \leq a \leq K_1(|\log \varepsilon| + 1),$$

and the map $\tilde{\Phi}$ from E_ε^a to the configuration space. We refer reader to [AB3] for the details.

Here is the notion of η -almost continuity introduced in [AB3]: A map $\Phi : X \rightarrow Y$ from a metric space X to a metric space Y is said to be η -almost continuous, if for all $x \in X$ and $\varepsilon > 0$, there is a δ , such that for all x' with $d_X(x, x') < \delta$, we have $d_Y(\Phi(x), \Phi(x')) \leq \eta + \varepsilon$. Proposition 4 says that the map $\tilde{\Phi}$ is actually η -almost equi-continuous for $\eta = C_1(|\log \varepsilon| \varepsilon^{\frac{2}{4k+1}} + \varepsilon^{\bar{\chi}})$.

By Theorem 3, the image of $\tilde{\Phi}$ lies in the set

$$\Sigma_{k,\beta}(\Omega) = \{ \{a_1, \dots, a_k\} \in \Sigma_k(\Omega); \text{dist}(a_i, \partial\Omega) \geq \beta, \text{ and } |a_i - a_j| \geq \beta \text{ for } i \neq j \}$$

which is compact in Σ_k . So we have

PROPOSITION 5 [AB3]. – We have an η_0 which only depends on β , such that for any $\eta \leq \eta_0$ and compact set $W \in H_g^1(\Omega, \mathbb{C})$, if $\tilde{\Phi}$ is η -almost continuous and $\tilde{\Phi}(W) \subset \Sigma_{k,\beta}(\Omega)$, then there exists a continuous map $\Phi : W \rightarrow \Sigma_k(\Omega)$ such that

$$\|\tilde{\Phi}(u) - \Phi(u)\| \leq 3\eta \quad \text{for all } u \in W.$$

3. PROOF OF THEOREM 1

In this section, we are going to prove Theorem 1 which is stated in §1.

Let $K \subset \Sigma_k(\Omega)$ be a compact core, i.e., K is compact and the natural inclusion $i : K \rightarrow \Sigma_k(\Omega)$ is a homotopy equivalence. Actually, $\Sigma_{k,\beta}(\Omega)$

is a compact core for sufficiently small β . We start with a construction of maps $f_\varepsilon : K \rightarrow E_\varepsilon^a$.

LEMMA 6. – *There are constants $\varepsilon_0'' > 0$, λ and C_2 such that for all $\varepsilon \leq \varepsilon_0''$, we can define $f_\varepsilon : K \rightarrow E_\varepsilon^a$, where $a = k\pi |\log \varepsilon| + \lambda$ such that*

$$\|\tilde{\Phi} \cdot f_\varepsilon - \text{id}\| \leq \eta$$

on K , where η is given by

$$\eta = C_2 \left(|\log \varepsilon| \varepsilon^{\frac{2}{4k+1}} + \varepsilon^{\bar{\chi}} \right).$$

Proof. – Since K is compact, we can pick $\eta_K > 0$ such that for any $\{a_1, \dots, a_k\} \in K$, the balls $B(a_i, 4\eta_K) \subset \Omega$ and are pairwise disjoint. Now once $\varepsilon \leq 4\eta_K$, we can construct a map $f_\varepsilon : \Sigma_k(\Omega) \rightarrow H_g^1(\Omega, \mathbb{C})$ as follows: for any $a = \{a_1, \dots, a_k\} \in \Sigma_k(\Omega)$, let

$$\Omega_{\varepsilon,a} = \Omega \setminus \bigcup_{i=1}^k B(a_i, \varepsilon),$$

then on $\Omega_{\varepsilon,a}$, $f_\varepsilon(a)$ is defined by

$$f_\varepsilon(a)(z) = e^{i\varphi_{\varepsilon,a}(z)} \prod_{j=1}^k \frac{z - a_j}{|z - a_j|}$$

where the function $\varphi_{\varepsilon,a}$ is defined on Ω by the following equation

$$\begin{cases} \Delta\varphi_{\varepsilon,a}(z) = 0 & \text{in } \Omega \\ e^{i\varphi_{\varepsilon,a}(z)} \prod_{j=1}^k \frac{z - a_j}{|z - a_j|} = g & \text{on } \partial\Omega. \end{cases}$$

Notice that for a given a the map $\varphi_{\varepsilon,a}$ is uniquely defined, up to an integer multiple of 2π . In fact, we can choose this constant such that the map $a \rightarrow e^{i\varphi_{\varepsilon,a}}$ is continuous by the standard lifting argument. On each $B(a_i, \varepsilon)$, $f_\varepsilon(a)$ is defined by

$$\begin{cases} \Delta f_\varepsilon(a) = 0 & \text{in } B(a_i, \varepsilon) \\ f_\varepsilon(a)(z) = e^{i\varphi_{\varepsilon,a}(z)} \prod_{j=1}^k \frac{z - a_j}{|z - a_j|} & \text{on } \partial B(a_i, \varepsilon). \end{cases}$$

It is then easy to check that f_ε is a continuous map from $\Sigma_k(\Omega)$ to $H_g^1(\Omega, \mathbb{C})$.

Moreover we can estimate the energy $E_\varepsilon(f_\varepsilon(a))$. Using the same analysis as in [Section I, BBH2], we have a constant C which depends on Ω and g only, such that

$$E_\varepsilon(f_\varepsilon(a)) \leq W_g(a_1, \dots, a_k) + k\pi|\log \varepsilon| + C.$$

Let

$$\lambda' = \sup_{a \in K} W_g(a_1, \dots, a_k),$$

it is finite by the compactness of K . So

$$E_\varepsilon(f_\varepsilon(a)) \leq k\pi|\log \varepsilon| + \lambda' + C.$$

Hence there is an $\varepsilon_1 > 0$ such that for all $\varepsilon \leq \varepsilon_1$, $f_\varepsilon(a) \in E_\varepsilon^a$, for $a = \mu_\varepsilon + \lambda$ provided λ is chosen large enough (but independent of ε).

Now suppose that $\varepsilon \leq \varepsilon_0'' = \min\{\varepsilon_0', \varepsilon_1, 4\eta_K\}$, and denote $f_\varepsilon(a)$ by $f_{\varepsilon,a}$ for simplicity. Let $a = \{a_1, \dots, a_k\}$ be given in K , and $a' = \{a'_1, \dots, a'_k\}$ be the vortices for $(f_{\varepsilon,a})^h$, i.e., $\Phi(f_{\varepsilon,a}) = \{a'_1, \dots, a'_k\}$. According to Theorem 3, on $\Omega_{\rho,a'} = \Omega \setminus \bigcup_{i=1}^k B(a'_i, \rho)$, we have

$$|f_{\varepsilon,a}^h(x)| \geq \frac{1}{2}, \text{ for all } x \in \Omega_{\rho,a'},$$

where $\varepsilon^x \leq \rho \leq \varepsilon^{\bar{x}}$. We may therefore consider on $\tilde{\Omega} = \Omega_{\rho,a'} \setminus \bigcup_{i=1}^k B(a_i, \varepsilon)$, the map $\xi = \frac{f_{\varepsilon,a}^h}{|f_{\varepsilon,a}^h|} f_{\varepsilon,a}^{-1}$. ξ takes its values in S^1 and satisfies $\xi \equiv 1$ on $\partial\Omega$. Moreover we have

$$|\xi - 1| \leq 4|f_{\varepsilon,a}^h - f_{\varepsilon,a}|.$$

This yields

$$\begin{aligned} \int_{\tilde{\Omega}} |\xi - 1|^2 &\leq 16 \int_{\tilde{\Omega}} |f_{\varepsilon,a}^h - f_{\varepsilon,a}|^2 \leq 32\varepsilon^{\frac{4}{4k+1}} (E_\varepsilon(f_{\varepsilon,a}) - E_\varepsilon(f_{\varepsilon,a}^h)) \\ &\leq C|\log \varepsilon| \varepsilon^{\frac{4}{4k+1}} \end{aligned} \tag{3.1}$$

for some constant C depending only on g, K and Ω .

On the other hand, for any $1 \leq i \leq k$, we have

$$\deg(\xi, \partial B(a'_i, \rho)) = -\deg(\xi, \partial B(a_i, \varepsilon)) = 1.$$

So for any regular value $y \in S^1$ of ξ and $y \neq 1$, $\xi^{-1}(y)$ is a connection between balls $B(a_i, \varepsilon)$ and $B(a'_i, \rho)$. By the definition of length of minimal connection L given in (2.2), we get

$$L(a', a) - k(\rho + \varepsilon) \leq \mathcal{H}^1(\xi^{-1}(y)) \text{ for almost every } y \in S^1.$$

Let

$$N = \left\{ y \in S^1, \frac{1}{8} \leq |y - 1| \leq \frac{1}{4} \right\}$$

and take $A = \xi^{-1}(N)$, using the coarea formula of Federer-Fleming, we obtain

$$\begin{aligned} \int_N \mathcal{H}^1(\xi^{-1}(y)) dy &= \int_A |\nabla \xi| \\ &\leq \left(\int_A |\nabla \xi|^2 \right)^{1/2} (\text{meas } A)^{1/2}. \end{aligned} \tag{3.2}$$

By (3.1), we have

$$(\text{meas } A) \leq 64 \int_{\tilde{\Omega}} |\xi - 1|^2 \leq C |\log \varepsilon| \varepsilon^{\frac{4}{4k+1}};$$

On the other hand

$$\int_{\tilde{\Omega}} |\nabla \xi|^2 \leq 8 \left(\int_{\Omega} |\nabla f_{\varepsilon,a}^h|^2 + |\nabla f_{\varepsilon,a}|^2 \right) \leq C |\log \varepsilon|.$$

Together with (3.2) we get that

$$L(a, a') - k(\rho + \varepsilon) \leq \frac{1}{(\text{meas } N)} \int_N \mathcal{H}^1(\xi^{-1}(y)) dy \leq C |\log \varepsilon| \varepsilon^{\frac{2}{4k+1}},$$

that is the conclusion we required. □

For any $a \in K$, the ball $B(a, 4\eta_K) \subset \Sigma_k(\Omega)$ with radius $4\eta_K$, where η_K is the constant in the proof of Lemma 6, is in fact isometric to a standard ball in \mathbb{C}^k . To see this, let $\hat{K} = \pi^{-1}(K) \subset F_k(\Omega)$, which is also a compact core of $F_k(\Omega)$, and for any $\tilde{a} \in \pi^{-1}(a)$, the condition that $B(a_i, 4\eta_K)$'s are pairwise disjoint implies that the ball $B(\tilde{a}, 4\eta_K) \subset \mathbb{C}^k$ is contained in $F_k(\Omega)$ entirely, and $B(a, 4\eta_K)$ is isometric to $B(\tilde{a}, 4\eta_K)$.

LEMMA 7. – *There is an ε_0 , such that for any $\varepsilon \leq \varepsilon_0$, the map f_ε induces an injection*

$$f_{\varepsilon*} : \pi_1(K) \rightarrow \pi_1(E_\varepsilon^a),$$

where a is chosen by Lemma 6.

Proof. – The constant $\varepsilon_0 \leq \varepsilon_0''$ is chosen such that

$$\max(C_1, C_2) \left(|\log \varepsilon_0| \varepsilon_0^{\frac{2}{4k+1}} + \varepsilon_0^{\bar{\lambda}} \right) \leq \min\{\eta_0, \eta_K\},$$

where C_1 is as in Proposition 4, η_0 as in Proposition 5 and ε''_0, C_2 and η_K as in Lemma 6.

For each element $\alpha \in \pi_1(K)$, we can choose a closed path $c : S^1 \rightarrow K$ which representing α . Now for $\varepsilon \leq \varepsilon_0$, if $f_\varepsilon \cdot c : S^1 \rightarrow E_\varepsilon^a$ is null homotopic, we get a map $\tilde{f} : D^2 \rightarrow E_\varepsilon^a$, such that $\tilde{f}|_{\partial D^2} = f_\varepsilon \cdot c$. By Proposition 5, on the compact set $\tilde{f}(D^2) \subset E_\varepsilon^a$, we can define a continuous map $\Phi : \tilde{f}(D^2) \rightarrow \Sigma_k(\Omega)$, such that for any $u \in \tilde{f}(D^2)$, $\|\Phi(u) - \tilde{\Phi}(u)\| < 3\eta_K$. The map $\Phi \cdot \tilde{f}|_{\partial D^2} = \Phi \cdot f_\varepsilon \cdot c : S^1 \rightarrow \Sigma_k(\Omega)$ is null homotopic. On the other hand, by Lemma 6,

$$\|\Phi \cdot f_\varepsilon \cdot c(t) - c(t)\| \leq \|\Phi \cdot f_\varepsilon \cdot c(t) - \tilde{\Phi} \cdot f_\varepsilon \cdot c(t)\| + \|\tilde{\Phi} \cdot f_\varepsilon \cdot c(t) - c(t)\| < 4\eta_K.$$

Then we can find a unique minimum geodesic in $\Sigma_k(\Omega)$ connecting $\Phi \cdot f_\varepsilon \cdot c(t)$ and $c(t)$. This implies that $\Phi \cdot f_\varepsilon \cdot c$ is homotopic to c . So α is a trivial element in $\pi_1(K)$, and this means that f_{ε^*} is injective. \square

Since $\pi_* : \pi_1(\tilde{K}) \rightarrow \pi_1(K)$ and $f_{\varepsilon^*} : \pi_1(K) \rightarrow \pi_1(E_\varepsilon^a)$ are injective, so is $f_{\varepsilon^*} \cdot \pi_* : \pi_1(\tilde{K}) \rightarrow \pi_1(E_\varepsilon^a)$. Consider a covering space $p : \tilde{E}_\varepsilon^a \rightarrow E_\varepsilon^a$ corresponding to the group $f_{\varepsilon^*} \cdot \pi_*(\pi_1(\tilde{K})) \subset \pi_1(E_\varepsilon^a)$, the map $f_\varepsilon \cdot \pi : \tilde{K} \rightarrow E_\varepsilon^a$ can be lift to a map $\tilde{f} : \tilde{K} \rightarrow \tilde{E}_\varepsilon^a$ such that the following diagram commutes

$$\begin{array}{ccc} \tilde{K} & \longrightarrow & \tilde{E}_\varepsilon^a \\ \downarrow & & \downarrow \\ K & \longrightarrow & E_\varepsilon^a. \end{array}$$

LEMMA 8. – *The map \tilde{f} induces maps $\tilde{f}_* : H_p(\tilde{K}) \rightarrow H_p(\tilde{E}_\varepsilon^a)$ on the homology groups which are injective for all p .*

Proof. – The argument here goes in the same fashion as the proof of Lemma 7. Consider a singular cycle $c \in Z_p(\tilde{K})$ such that $\tilde{f}_*([c]) = 0$ in $H_p(\tilde{E}_\varepsilon^a)$. This means that we have a $p + 1$ -chain $c' \in C_{p+1}(\tilde{E}_\varepsilon^a)$ and $\partial c' = \tilde{f}_*(c)$. The set $W = \tilde{f}(\tilde{K}) \cup \text{support}(c')$ is compact in \tilde{E}_ε^a . Then we define a continues map $\Phi_1 : p(W) \rightarrow \Sigma_k(\Omega)$ such that for any $u \in p(W)$, $\|\Phi_1(u) - \tilde{\Phi}(u)\| < 3\eta_K$.

Notice that $\|\Phi_1 \cdot f_\varepsilon - \text{id}\| < 4\eta_K$, as before, we have $\Phi_{1*} \cdot f_{\varepsilon^*} = \text{id}$. This implies that $\Phi_{1*} \cdot p_*(\pi_1(W)) \subset \Phi_{1*} \cdot f_{\varepsilon^*} \cdot \pi_*(\pi_1(\tilde{K})) = \pi_*(\pi_1(\tilde{K}))$. So we can lift $\Phi_1 \cdot p : W \rightarrow \Sigma_k(\Omega)$ to $\tilde{\Phi}_1 : W \rightarrow \tilde{F}_k(\Omega)$.

In fact, we can make $\|\tilde{\Phi}_1 \cdot \tilde{f} - \text{id}\| < 4\eta_K$. Since $\|\Phi_1 \cdot p \cdot \tilde{f} - \pi\| < 4\eta_K$, there is a homotopy H_t such that $H_0 = \pi$ and $H_1 = \Phi_1 \cdot p \cdot \tilde{f}$. Lift this homotopy to a homotopy \tilde{H}_t with $\|\tilde{H}_0 - \tilde{H}_1\| < 4\eta_K$ and $\tilde{H}_0 = \text{id}_{\tilde{K}}$. Define $\tilde{\Phi}_2 : \tilde{f}(\tilde{K}) \rightarrow \tilde{F}_k(\Omega)$ by $\tilde{\Phi}_2(\tilde{f}(a)) = \tilde{H}_1(a)$. Note that

$$\pi \cdot \tilde{\Phi}_2 = \Phi_1 \cdot p = \pi \cdot \tilde{\Phi}_1|_{\tilde{f}(\tilde{K})}.$$

$\tilde{\Phi}_2$ and $\tilde{\Phi}_1$ differ by a deck transformation, i.e., there is an elements $\sigma \in S^k$, such that

$$\tilde{\Phi}_2 = \sigma \cdot \tilde{\Phi}_1|_{\tilde{f}(\tilde{K})}.$$

Replace $\tilde{\Phi}_1$ by $\sigma \cdot \tilde{\Phi}_1$, which is also a lifting of $\Phi_1 \cdot p : W \rightarrow \Sigma_k(\Omega)$ and $\|\sigma \cdot \tilde{\Phi}_1 \cdot \tilde{f} - \text{id}\| < 4\eta_K$. The new lifting will still denoted by $\tilde{\Phi}_1$.

Now $\tilde{\Phi}_1$ maps the chain c' into a chain in $C_{p+1}(F_k(\Omega))$, and $\partial\tilde{\Phi}_1(c') = \tilde{\Phi}_1(\partial c') = \tilde{\Phi}_1 \cdot \tilde{f}_*(c)$. We get that $\tilde{\Phi}_1 \cdot \tilde{f}_*(c)$ is a boundary in $C_p(F_k(\Omega))$. On the other hand, $\tilde{\Phi}_1 \cdot \tilde{f}$ is homotopic to the natural inclusion $i : \tilde{K} \rightarrow F_k(\Omega)$. So c is homologous to $\tilde{\Phi}_1 \cdot \tilde{f}_*(c)$, and c is null homologous as well. This shows that \tilde{f}_* is injective. □

The lemma allows us to estimate the category of \tilde{E}_ε^a .

COROLLARY 9. – *The category $\text{cat}(\tilde{E}_\varepsilon^a)$ of \tilde{E}_ε^a is at least k .*

Proof. – By Lemma 8, the map $f^* : H^*(\tilde{E}_\varepsilon^a) \rightarrow H^*(\tilde{K})$ between cohomology rings are surjective, and this implies that the cuplength of E_ε^a is at least the cuplength of \tilde{K} , which is the same as the cuplength of $F_k(\Omega)$. By Lemma 2, the cuplength of \tilde{E}_ε^a is at least $k - 1$. Finally, according to [BG], the category $\text{cat}(\tilde{E}_\varepsilon^a)$ of \tilde{E}_ε^a is at least the cuplength of \tilde{E}_ε^a plus one. This completes the proof. □

Now we are in the position to complete the proof of Theorem 1. The Lusternik-Schnirelman minimax theorem we will use is the following

THEOREM 10. – *Suppose F is a C^2 non-negative functional defined on a smooth Hilbert manifold M such that*

- i) the backwards gradient flow is complete;*
- ii) F satisfies the following weak Palais-Smale condition: if we have a sequence $\{u_n\}$ in M such that $F(u_n) \rightarrow c$ and $\|\nabla F(u_n)\| \rightarrow 0$ as $n \rightarrow \infty$, then c is a critical value;*
- iii) $\text{cat}M = k$.*

Then we have either F has at least k distinct critical values in $[0, a]$ or the dimension of the critical set of F is at least 1.

The proof is standard, we refer reader to [Pa].

Proof of Theorem 1. – Now we want to apply Theorem 10 to the positive functional $\tilde{E}_\varepsilon = E_\varepsilon \cdot p : \tilde{E}_\varepsilon^a \rightarrow \mathbb{R}$. Notice that \tilde{E}_ε and E_ε have the same critical values and critical sets of the two functionals have the same dimension. If all three conditions in the theorem hold, both conclusions will imply that E_ε has at least k critical points on E_ε^a .

We now check the three conditions in Theorem 10. First, the backwards gradient flow of \tilde{E}_ε is a lift of the backwards flow of E_ε , so it is

complete. Second, let $\{u_n\}$ be a sequence in \tilde{E}_ε^a such that $\tilde{E}_\varepsilon(u_n) \rightarrow c$ and $\|\nabla \tilde{E}_\varepsilon(u_n)\| \rightarrow 0$ as $n \rightarrow \infty$, then $E_\varepsilon(p(u_n)) \rightarrow c$ and $\|\nabla E_\varepsilon(p(u_n))\| \rightarrow 0$. We know that E_ε satisfies Palais-Smale condition, so $p(u_n)$ has a subsequence converges to a critical point. This shows that c is a critical value of E_ε and then it is a critical value of \tilde{E}_ε as well. Finally, $\text{cat} \tilde{E}_\varepsilon^a \geq k$ is the conclusion of Corollary 9. So we now can conclude that E_ε has at least k critical points on E_ε^a .

Outside of E_ε^a , E_ε has at least another critical point, since $H_g^1(\Omega, \mathbb{C})$ is contractible, but E_ε^a is not (if $k \geq 2$). So totally E_ε will have at least $k + 1$ critical points on $H_g^1(\Omega, \mathbb{C})$. \square

REFERENCES

- [AB1] L. ALMEIDA and F. BETHUEL, Multiplicity results for the Ginzburg-Landau equation in presence of symmetries, to appear in *Houston Math. J.*
- [AB2] L. ALMEIDA and F. BETHUEL, Méthodes topologiques pour l'équation de Ginzburg-Landau, *C. R. Acad. Sci. Paris*, Vol. **320**, 1996, pp. 935-938.
- [AB3] L. ALMEIDA and F. BETHUEL, Topological methods for the Ginzburg-Landau equations, to appear in *J. Math. Pures et Appl.*, 1996.
- [Ar] V. I. ARNOL'D, The cohomology ring of the colored braid group (English, Russian original), *Math. Notes*, Vol. **5**, 1969, pp. 138-140 ; translation from, *Mat. Zametki*, Vol. **5**, 1969, pp. 227-231.
- [BBH1] F. BETHUEL, H. BREZIS and F. HÉLEIN, Asymptotics for the minimization of a Ginzburg-Landau functional, *Calc. of Vari. and PDE's*, Vol. **1**, 1993, pp. 123-148.
- [BBH2] F. BETHUEL, H. BREZIS and F. HÉLEIN, Ginzburg-Landau Vortices, Birkhäuser, 1994.
- [BCL] H. BREZIS, J. M. CORON and E. H. LIEB, Harmonic maps with defects, *Comm. Math. Phys.*, Vol. **107**, 1986, pp. 649-705.
- [BG] I. BERNSTEIN and T. GANEA, Homotopical nilpotency, *Illinois J. Math.*, Vol. **10**, 1961, pp. 99-130.
- [FP] P. FELMER and M. DEL PINO, Local minimizers for the Ginzburg-Landau energy, Preprint.
- [Li] F. H. LIN, Solutions of Ginzburg-Landau equations and critical points of the renormalized energy, *Ann. IHP, Analyse Non Linéaire*, Vol. **12**, 1995, pp. 599-622.
- [Pa] R. S. PALAIS, Ljusternik-Schnirelman theory on Banach manifolds, *Topology*, Vol. **5**, 1966, pp. 115-132.

(Manuscript received April 30, 1997;
revised October 27, 1997.)