

ANNALES DE L'I. H. P., SECTION C

ETIENNE SANDIER

SYLVIA SERFATY

Global minimizers for the Ginzburg-Landau functional below the first critical magnetic field

Annales de l'I. H. P., section C, tome 17, n° 1 (2000), p. 119-145

http://www.numdam.org/item?id=AIHPC_2000__17_1_119_0

© Gauthier-Villars, 2000, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Global minimizers for the Ginzburg–Landau functional below the first critical magnetic field

by

Etienne SANDIER^{a,1}, Sylvia SERFATY^{b,c,2}

^a Université François Rabelais, Département de Mathématiques, Parc Grandmont,
37200 Tours, France

^b CMLA, Ecole Normale Supérieure de Cachan, 61 avenue du Président Wilson,
94235 Cachan Cedex, France

^c Laboratoire d'Analyse Numérique et EDP, Bâtiment 425, Université de Paris-Sud,
91405 Orsay, France

Manuscript received 5 October 1998

ABSTRACT. – We prove that the global minimizer of the Ginzburg–Landau functional of superconductors in an external magnetic field is, below the first critical field, the vortex-less solution found in (S. Serfaty, to appear). © 2000 Éditions scientifiques et médicales Elsevier SAS

RÉSUMÉ. – On montre que le minimiseur global de la fonctionnelle de Ginzburg–Landau en supraconductivité, avec champ magnétique extérieur inférieur au premier champ critique est la solution sans vortex trouvée dans (S. Serfaty, à paraître). © 2000 Éditions scientifiques et médicales Elsevier SAS

¹ E-mail: sandier@univ-tours.fr.

² E-mail: serfaty@cmla.ens-cachan.fr.

1. INTRODUCTION

We are interested in this paper in describing the minimizers of the Ginzburg–Landau functional

$$J(u, A) = \frac{1}{2} \int_{\Omega} |\nabla_A u|^2 + |h - h_{ex}|^2 - h_{ex}^2 + \frac{\kappa^2}{2} (1 - |u|^2)^2, \quad (1.1)$$

that corresponds to the free energy of a superconductor in a prescribed, constant magnetic field h_{ex} . Here, $\Omega \subset \mathbb{R}^2$ is the smooth, bounded, and simply connected section of the superconductor; and the unknowns are the complex-valued order parameter $u \in H^1(\Omega, \mathbb{C})$ and the $U(1)$ connection $A \in H^1(\Omega, \mathbb{R}^2)$. The *induced* magnetic field h is defined by $h = \text{curl } A$. The order parameter u indicates the local state of the material: $|u|$ is the density of superconducting electron pairs, so that, where $|u| \simeq 1$, the material is in its superconducting state, whereas where $|u| \simeq 0$, it is in its normal state. Finally $\kappa = 1/\varepsilon > 0$ is the Ginzburg–Landau parameter depending on the material, and we are interested in the case of superconductors with high kappa. We stress that no boundary conditions are imposed on (u, A) , the characteristics of the solutions to the minimization problem are governed solely by h_{ex} .

Minimizers of $J(u, A)$ solve the associated Euler equations,

$$\begin{cases} -\nabla_A^2 u = \kappa^2 u(1 - |u|^2), \\ -*dh = (iu, d_A u). \end{cases} \quad (\text{G.L.})$$

It turns out that a key physical feature of solutions to (G.L.) is the existence, for a certain range of values of h_{ex} , of *vortices*, i.e., isolated zeros of u at which u has a nonzero *degree*: $u/|u|$ restricted to a small circle around the zero has a non-zero winding number as a map to the unit circle. Away from these zeros, $|u| \approx 1$. Describing solutions to (G.L.) then typically consists in describing the *vortex structure* of the solutions, i.e., to determine the number, degree and position of vortices.

The difficulty here is that without boundary conditions, there is no a-priori bound on the number of vortices. Even if boundary conditions are imposed, the problem of defining mathematically and describing the vortex structure of solutions is not an easy one. This was done in [3] for

the functional

$$F(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \frac{1}{2\varepsilon^2} (1 - |u|^2)^2 \quad (1.2)$$

supplemented by a Dirichlet boundary condition $g : \partial\Omega \rightarrow S^1$. There, a vortex structure is shown to exist for minimizers (and even critical points) of $F(u)$ when $\varepsilon \rightarrow 0$. Moreover it is proved that there are exactly $d = |\deg(g)|$ vortices, and their position is determined. In [5] the functional (1.1) is studied with h_{ex} set to zero and replaced by a gauge invariant variant of the Dirichlet condition. There again the vortex structure is shown to exist when ε is small, and results similar to those in [3] are obtained about the number and position of the vortices.

In [13,14] and [15], the second author studied minimizers of (1.1) without boundary condition. For more details on the notations and physical description of superconductors, we refer to [13] and the references therein. Let us just say that it is observed that for small values of the applied field h_{ex} , the material is superconducting everywhere (there are no vortices), the magnetic field does not penetrate it and approximately satisfies the London equation

$$\begin{cases} -\Delta h + h = 0 & \text{in } \Omega, \\ h = h_{ex} & \text{on } \partial\Omega. \end{cases} \quad (1.3)$$

This state is called the Meissner state, corresponding to vortex-less solutions in the terminology of [13] and [14], the corresponding solution (u, A) to (G.L.) being called the Meissner solution. For h_{ex} higher than some critical value H_{c_1} , the vortex-less solution is no longer energy minimizing.

In [13], to replace the absence of boundary conditions and thus the lack of a-priori estimates on the number of vortices, the functional $J(u, A)$ was minimized on a subdomain of $H^1(\Omega, \mathbb{C}) \times H^1(\Omega, \mathbb{R}^2)$. More precisely, choosing some (large) number $\mathcal{M} > 0$, the minimization was performed on

$$D = \left\{ (u, A) \mid \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \frac{1}{2\varepsilon^2} (1 - |u|^2)^2 < \mathcal{M} |\log \varepsilon| \right\}, \quad (1.4)$$

and the following theorem was proved:

THEOREM 1 [13]. – *There exist $k_1 = 1/(2 \max |\xi_0|)$, $k_2^\varepsilon = O_\varepsilon(1)$ and $k_3^\varepsilon = o_\varepsilon(1)$, such that*

$$H_{c_1} = k_1 |\log \varepsilon| + k_2^\varepsilon,$$

and $\varepsilon_0(\mathcal{M})$ such that for $\varepsilon < \varepsilon_0$, the following holds:

- if $h_{ex} \leq H_{c_1}$, a solution of (G.L.) that is minimizing in D exists, and satisfies $\frac{1}{2} \leq |u| \leq 1$;
- if $H_{c_1} + k_3^\varepsilon \leq h_{ex} \leq H_{c_1} + O_\varepsilon(1)$, a solution of (G.L.) that is minimizing in D exists, it has a bounded positive number of vortices a_i^ε of degree one, such that $\text{dist}(a_i^\varepsilon, \Lambda) \rightarrow 0$ where

$$\Lambda = \{x \in \Omega \mid |\xi_0(x)| = \max |\xi_0|\},$$

and $\exists C > 0$, $\text{dist}(a_i^\varepsilon, a_j^\varepsilon) \geq C$ for $i \neq j$, i.e., the a_i 's tend to distinct points $\in \Lambda$.

In addition, it is proved in [15] that the Meissner solution found for $h_{ex} \leq H_{c_1}$ still exists for $h_{ex} \geq H_{c_1}$ (even though it is then only locally minimizing in D), and is unique among vortex-less solutions.

H_{c_1} is known as the first critical field. It is the value of h_{ex} for which the energy of the Meissner solution becomes equal to the energy of a single-vortex configuration. Here, we wish to know whether or not, for $h_{ex} \leq H_{c_1}$, the Meissner solution is a global minimizer of the energy in addition to being a minimizer in D .

This question arises naturally for all the vortex solutions found in [13] and [14] that are minimizers in D , but that are all likely to be global minimizers. However, the proof in [13] and [14] uses repeatedly the a-priori bound on the number of vortices given by (1.4). Here, without this upper bound, we are still able to prove a result about vortex-less solutions:

THEOREM 1. – *There exists a value $H'_{c_1} \simeq H_{c_1}$ (more precisely $H'_{c_1} = H_{c_1} + O(|\log |\log \varepsilon||)$), such that, for sufficiently small ε , if $h_{ex} \leq H'_{c_1}$, a globally minimizing solution of (G.L.) satisfies $|u| \geq 3/4$ on Ω , and coincides with the solution found in Theorem 1 of [13].*

Thus, we answer positively the question, though we have an imprecision on H_{c_1} that we were not able to avoid.

In order to prove this theorem, we consider a minimizing solution of (G.L.) and assume it has possible vortices. We use a technique of R. Jerrard [7] to construct balls $B_i = B(a_i, r)$ of size $r \simeq 1/|\log \varepsilon|^\alpha$ with sufficiently high α , such that

$$|u| \geq \frac{3}{4} \quad \text{on } \Omega \setminus \bigcup_i B_i, \tag{1.5}$$

$$d_i = \text{deg}(u, \partial B_i), \tag{1.6}$$

$$F_\varepsilon(u, B_i) > \pi |d_i| \log \frac{r}{\varepsilon}, \tag{1.7}$$

where we use the notation

$$F_\varepsilon(u, V) = \frac{1}{2} \int_V |\nabla u|^2 + \frac{1}{2\varepsilon^2} (1 - |u|^2)^2.$$

The lower bound (1.7) was known to be true, see [3], *under some assumptions on the restriction of u on ∂B_i* . Here, adapting slightly the techniques in [7], we are able to avoid making these assumptions and to construct these balls even though, in contrast to [13,14] and [15], their number is not bounded a priori independently of ε .

Then the key argument of the proof is to split conveniently the energy in a way that is similar to, but slightly different from the one used in [13], in order to obtain a lower bound of the energy on $\Omega \setminus \bigcup_i B_i$. Precisely, we find that for a minimizer (u, A) ,

$$J(u, A) \geq J_0 + 2\pi h_{ex} \sum_i d_i \xi_0(a_i) + \sum_i F_\varepsilon(u, B_i) + o(1), \tag{1.8}$$

as $\varepsilon \rightarrow 0$. In the above expansion J_0 is roughly the minimal energy of a vortex-less configuration; and ξ_0 is a negative function depending only on the domain Ω (see [13] or Section 2). Putting together (1.7) and (1.8), and using the fact that the energy of a minimizer must be no greater than J_0 allows to conclude that vortices are not present if h_{ex} is less than some number

$$H'_{c_1} \approx \frac{1}{2 \max |\xi_0|} |\log \varepsilon|,$$

the right-hand side is precisely the H_{c_1} computed in [13].

The idea is the same as in [13]: a vortex of degree d “costs” almost $\pi |d| |\log \varepsilon|$ to make (see (1.7)), while it can decrease $J(u, A)$ by at most $\pi d h_{ex} \xi_0(a_i)$ (see (1.8)). Note that the choice of $r = 1/|\log \varepsilon|^\alpha$ for the size of the balls B_i is dictated by the fact that it is the largest radius for which we can prove that the expansion (1.8) is valid.

Of course, the structure of global minimizers for $h_{ex} > H_{c_1}$ is still open. We study them in [11], give an estimate on their energy, and

we expect that the number of vortices is equivalent to h_{ex} as soon as $h_{ex} \gg H_{c_1}$.

2. PROOF OF THE THEOREM

2.1. Preliminary results and notations

We consider (u, A) such that

$$J(u, A) = \inf\{J(v, B) \mid (v, B) \in H^1(\Omega, \mathbb{C}) \times H^1(\Omega, \mathbb{R}^2)\}.$$

By a standard argument, this infimum is achieved and yields a solution of the Ginzburg–Landau equations. We recall that, as in [13], for a suitable choice of gauge, $\operatorname{div} A = 0$, and there is a function $\xi \in H^2(\Omega, \mathbb{R})$ such that

$$A = \nabla^\perp \xi = (-\xi_{x_2}, \xi_{x_1}), \quad \xi = 0 \text{ on } \partial\Omega, \tag{2.1}$$

thus

$$h = \Delta \xi, \tag{2.2}$$

and

$$|\nabla_A u|^2 = |\nabla u|^2 - 2(iu, \xi_{x_2} u_{x_1} - \xi_{x_1} u_{x_2}) + |u|^2 |A|^2.$$

Our solution (u, A) is easily seen to satisfy

$$\begin{cases} \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega, \\ h = h_{ex} & \text{on } \partial\Omega. \end{cases}$$

In the sequel, C denotes any positive constant independent from ε .

Since the value of H_{c_1} computed in [13] is of the order of $|\log \varepsilon|$, we will assume from now on that

$$h_{ex} \leq C |\log \varepsilon|. \tag{2.3}$$

Considering the London equation (1.3), we are led as in [13] to introduce ξ_0 , the solution of

$$\begin{cases} -\Delta^2 \xi_0 + \Delta \xi_0 = 0 & \text{in } \Omega, \\ \Delta \xi_0 = 1 & \text{on } \partial\Omega, \\ \xi_0 = 0 & \text{on } \partial\Omega. \end{cases} \tag{2.4}$$

The approximate minimal vortex-less configuration is $(u_0 \equiv 1, A_0 = h_{ex} \nabla^\perp \xi_0)$ and, as in [13], we let

$$J_0 = J(u_0, A_0) = -\frac{h_{ex}^2}{2} \int_{\Omega} |\nabla \xi_0|^2 + |\Delta \xi_0|^2. \tag{2.5}$$

Note that (u_0, A_0) is only a solution to the second (G.L.) equation and not to the first one, therefore it is *not* the Meissner solution. However, it is proved in [13] that the infimum of the energy among vortex-less configurations in D is $J_0 + o(1)$ as $\varepsilon \rightarrow 0$.

As in [13], we decompose ξ as

$$\xi = h_{ex} \xi_0 + \zeta \tag{2.6}$$

so that

$$\zeta = 0, \quad \Delta \zeta = 0 \quad \text{on } \partial\Omega.$$

We state some results borrowed from [13] that are going to be useful in the sequel.

LEMMA 2.1. – *Let (u, A) be a solution of (G.L.). The following holds:*

$$\|\nabla u\|_{L^\infty(\Omega)} \leq \frac{C}{\varepsilon}, \tag{2.7}$$

$$\|\nabla \xi\|_{L^\infty(\Omega)} \leq Ch_{ex}. \tag{2.8}$$

If (u, A) is in addition a minimizer of the energy, then

$$J(u, A) \leq J_0 \leq Ch_{ex}^2, \tag{2.9}$$

$$\|\nabla u\|_{L^2(\Omega)} \leq Ch_{ex}. \tag{2.10}$$

Proof. – All the assertions have been proved in [13] except the last one. In view of (1.1) and (2.9),

$$\int_{\Omega} |\nabla_A u|^2 \leq Ch_{ex}^2,$$

which is equivalent to

$$\int_{\Omega} |\nabla u|^2 - 2(iu, \xi_{x_2} u_{x_1} - \xi_{x_1} u_{x_2}) + |u|^2 |A|^2 \leq Ch_{ex}^2.$$

Hence,

$$\int_{\Omega} |\nabla u|^2 \leq Ch_{ex}^2 + C \|\nabla \xi\|_{L^\infty(\Omega)}^2 + C \|\nabla \xi\|_{L^\infty(\Omega)} \|\nabla u\|_{L^2(\Omega)}.$$

Thus, in view of (2.8),

$$\|\nabla u\|_{L^2(\Omega)} \leq Ch_{ex}. \quad \square$$

We then need to define the vortices of u with their degrees, by defining balls $(B_i)_{i \in I}$, such that $|u| \geq 3/4$ on $\Omega \setminus \bigcup_{i \in I} B_i$, and $d_i = \text{deg}(u, \partial B_i)$. As already mentioned, we achieve this by adjusting a result of Jerrard [7], to obtain the following proposition, the proof of which is deferred to Section 3.

PROPOSITION 2.1. – *Let $u \in H^1(\Omega, \mathbb{C})$ such that $\|\nabla u\|_{L^\infty} \leq C/\varepsilon$, and $F(u) \leq Ch_{ex}^2$. Then, for any $\alpha > 0$, there is an $\varepsilon_0 > 0$ such that $\forall \varepsilon < \varepsilon_0$ there exists a finite family of disjoint balls $(B_i)_{i \in I} = (B(a_i, r_i))_{i \in I}$ such that*

$$\left\{ x / |u(x)| < \frac{3}{4} \right\} \subset \bigcup_{i \in I} B_i, \tag{2.11}$$

$$F_\varepsilon(u, B_i) \geq \pi |d_i| (|\log \varepsilon| - O(|\log |\log \varepsilon||)), \tag{2.12}$$

where $d_i = \text{deg}(u, \partial B_i)$ if $\overline{B_i} \subset \Omega$, and $d_i = 0$ otherwise,

$$r_i \leq C |\log \varepsilon|^{-\alpha}, \tag{2.13}$$

$$\text{Card } I \leq Ch_{ex}^2. \tag{2.14}$$

Proof. – See Proposition 3.2. \square

2.2. Splitting of the energy

Let

$$\tilde{\Omega} = \Omega \setminus \bigcup_{i \in I} B_i,$$

where $\{B_i\}_{i \in I}$ is the family of balls given by Proposition 2.1. Recall that they have radii r_i less than $C |\log \varepsilon|^{-\alpha}$, where α is to be chosen below.

LEMMA 2.2. – We have following identity:

$$\begin{aligned} \frac{1}{2} \int_{\Omega} |\nabla_{Au}|^2 &= \int_{\Omega} \left(\frac{1}{2} |\nabla u - i \nabla^{\perp} \zeta u|^2 + \frac{1}{2} h_{ex}^2 |\nabla \xi_0|^2 + h_{ex} \nabla \xi_0 \cdot \nabla \zeta \right) \\ &\quad + 2\pi h_{ex} \sum_{i \in I} d_i \xi_0(a_i) + o(1). \end{aligned}$$

Proof. – From (2.2) and (2.6),

$$\begin{aligned} |\nabla u - i Au|^2 &= |\nabla u - i h_{ex} \nabla^{\perp} \xi_0 u - i \nabla^{\perp} \zeta u|^2 \\ &= |\nabla u - i \nabla^{\perp} \zeta u|^2 + h_{ex}^2 |u|^2 |\nabla \xi_0|^2 \\ &\quad + 2(\nabla u - i \nabla^{\perp} \zeta u, -i h_{ex} \nabla^{\perp} \xi_0 u). \end{aligned} \tag{2.15}$$

Moreover,

$$\begin{aligned} &\int_{\Omega} (\nabla u - i \nabla^{\perp} \zeta u, -i h_{ex} \nabla^{\perp} \xi_0 u) \\ &= \int_{\Omega} (\nabla u, -i h_{ex} \nabla^{\perp} \xi_0 u) + h_{ex} \int_{\Omega} |u|^2 \nabla \xi_0 \cdot \nabla \zeta. \end{aligned} \tag{2.16}$$

To finish the proof of Lemma 2.2 we need the \square

LEMMA 2.3. – If $\alpha > 5$,

$$\int_{\Omega} (\nabla u, -i h_{ex} \nabla^{\perp} \xi_0 u) = 2\pi h_{ex} \sum_{i \in I} d_i \xi_0(a_i) + o(1). \tag{2.17}$$

Proof. – We start with the same method as in [13]. First,

$$\left| \int_{\bigcup B_i} (\nabla u, -i h_{ex} \nabla^{\perp} \xi_0 u) \right| \leq (\text{Card } I) h_{ex} \|\nabla u\|_{L^2} \max_{i \in I} r_i \leq C |\log \varepsilon|^{4-\alpha},$$

where we used (2.3), (2.13), (2.14), and (2.10). It is here that the size of the balls B_i is important. Then, letting $\tilde{\Omega} = \Omega \setminus \bigcup_i B_i$,

$$\int_{\tilde{\Omega}} (\nabla u, -i h_{ex} \nabla^{\perp} \xi_0 u) = h_{ex} \int_{\tilde{\Omega}} (i u, (\xi_0)_{x_2} u_{x_1} - (\xi_0)_{x_1} u_{x_2}).$$

Setting $v = u/|u|$ and integrating by parts, we find, exactly as in the proof of Lemma IV.3 of [13], that

$$\begin{aligned}
 & \int_{\tilde{\Omega}} (iu, (\xi_0)_{x_2} u_{x_1} - (\xi_0)_{x_1} u_{x_2}) \\
 &= \int_{\tilde{\Omega}} (iv, dv \wedge d\xi_0) + o(|\log \varepsilon|^{-1}) \\
 &= \sum_{i \in I} \int_{\partial B_i} \xi_0 \left(iv, \frac{\partial v}{\partial \tau} \right) + o(|\log \varepsilon|^{-1}). \tag{2.18}
 \end{aligned}$$

We claim that, letting $\mathcal{J} = \{i \in I / \overline{B_i} \subset \Omega\}$,

$$\forall i \in \mathcal{J}, \quad h_{ex} \int_{\partial B_i} \xi_0 \left(iv, \frac{\partial v}{\partial \tau} \right) = 2\pi h_{ex} d_i \xi_0(a_i) + o(|\log \varepsilon|^{-2}). \tag{2.19}$$

To prove this claim, we use the same proof as in [13] (Lemma IV.3), different from that of [5] which does not adjust to a possible divergent number of balls. Let

$$U_i = \left\{ x \in B_i / |u| \leq \frac{1}{2} \right\}.$$

U_i does not intersect ∂B_i and by Stokes' theorem

$$\begin{aligned}
 & \left| \int_{\partial B_i} (\xi_0 - \xi_0(a_i)) \left(iv, \frac{\partial v}{\partial \tau} \right) - \int_{\partial U_i} (\xi_0 - \xi_0(a_i)) \left(iv, \frac{\partial v}{\partial \tau} \right) \right| \\
 &= \left| \int_{B_i \setminus U_i} d\xi_0 \wedge (iv, dv) \right| \leq C \|\nabla \xi_0\|_{L^\infty} r_i \left(\int_{B_i \setminus U_i} |\nabla v|^2 \right)^{1/2} \\
 &\leq C |\log \varepsilon|^{1-\alpha}.
 \end{aligned}$$

Hence, as $\alpha > 5$,

$$\begin{aligned}
 & h_{ex} \int_{\partial B_i} (\xi_0 - \xi_0(a_i))(iv, dv) \\
 &= h_{ex} \int_{\partial U_i} (\xi_0 - \xi_0(a_i))(iv, dv) + o(|\log \varepsilon|^{-2}). \tag{2.20}
 \end{aligned}$$

On the other hand,

$$h_{ex} \int_{\partial U_i} (\xi_0 - \xi_0(a_i))(iv, dv) = h_{ex} \int_{\partial U_i} (\xi_0 - \xi_0(a_i)) \frac{(iu, du)}{|u|^2}.$$

But, by definition of U_i , $|u| = 1/2$ on ∂U_i , hence

$$\begin{aligned}
 & \left| h_{ex} \int_{\partial U_i} (\xi_0 - \xi_0(a_i))(iv, dv) \right| \\
 &= 4h_{ex} \left| \int_{\partial U_i} (\xi_0 - \xi_0(a_i))(iu, du) \right| \\
 &= 4h_{ex} \left| \int_{U_i} d\xi_0 \wedge (iu, du) + (\xi_0 - \xi_0(a_i))(idu, du) \right| \\
 &\leq Ch_{ex}r_i \|\nabla u\|_{L^2} + Ch_{ex}r_i \|\nabla \xi_0\|_{L^\infty} \int_{\Omega} |\nabla u|^2 \\
 &\leq C|\log \varepsilon|^{3-\alpha} \leq o(|\log \varepsilon|^{-2}).
 \end{aligned} \tag{2.21}$$

Combining (2.20) and (2.21), we conclude that

$$\begin{aligned}
 h_{ex} \int_{\partial B_i} \xi_0(iv, dv) &= h_{ex} \int_{\partial B_i} \xi_0(a_i)(iv, dv) + o(|\log \varepsilon|^{-2}) \\
 &= 2\pi h_{ex} d_i \xi_0(a_i) + o(|\log \varepsilon|^{-2}),
 \end{aligned}$$

and (2.19) is proved.

We now deal with the balls that intersect $\partial\Omega$. We claim that

$$\forall i \in I \setminus \mathcal{J}, \quad h_{ex} \int_{\partial B_i \cap \Omega} \xi_0(iv, dv) = o(|\log \varepsilon|^{-2}).$$

The proof of this claim is almost the same as that of (2.19). Indeed, since $\xi_0 = 0$ on $\partial\Omega$, letting $U_i = B_i \cap \{|u| \leq 1/2\}$,

$$\begin{aligned}
 & h_{ex} \int_{\partial B_i \cap \Omega} \xi_0(iv, dv) \\
 &= h_{ex} \int_{\partial U_i \cap \Omega} \xi_0(iv, dv) + o(|\log \varepsilon|^{-2}) \\
 &= 4h_{ex} \left| \int_{U_i \cap \Omega} d\xi_0 \wedge (iu, du) + \xi_0(idu, du) \right| = o(|\log \varepsilon|^{-2}).
 \end{aligned}$$

Using (2.18) and (2.19), the above claim and the fact that $\text{Card } I \leq Ch_{ex}^2$ prove the lemma. \square

Proof of Lemma 2.2 completed. – Combining (2.15), (2.16) and (2.17), we are led to

$$\begin{aligned} \frac{1}{2} \int_{\Omega} |\nabla_A u|^2 &= \frac{1}{2} \int_{\Omega} |\nabla u - i \nabla^\perp \zeta u|^2 + \frac{1}{2} h_{ex}^2 \int_{\Omega} |u|^2 |\nabla \xi_0|^2 \\ &\quad + h_{ex} \int_{\Omega} |u|^2 \nabla \xi_0 \cdot \nabla \zeta + 2\pi h_{ex} \sum_{i \in I} d_i \xi_0(a_i) + o(1). \end{aligned} \quad (2.22)$$

From the upper bound (2.9), we know that

$$\frac{1}{\varepsilon^2} \int_{\Omega} (1 - |u|^2)^2 \leq Ch_{ex}^2, \quad (2.23)$$

hence

$$\begin{aligned} h_{ex}^2 \int_{\Omega} (1 - |u|^2) |\nabla \xi_0|^2 &\leq Ch_{ex}^2 \left(\int_{\Omega} (1 - |u|^2)^2 \right)^{1/2} \\ &\leq C\varepsilon h_{ex}^3 \leq o(1), \end{aligned}$$

and similarly

$$h_{ex}^2 \left| \int_{\Omega} (1 - |u|^2) \nabla \xi_0 \cdot \nabla \zeta \right| \leq C\varepsilon h_{ex}^3 \leq o(1).$$

With (2.22), the lemma is proved. \square

LEMMA 2.4. – *We have the following identity:*

$$\begin{aligned} &\frac{1}{2} \int_{\Omega} (|h - h_{ex}|^2 - h_{ex}^2) + \frac{1}{2} h_{ex}^2 \int_{\Omega} |\nabla \xi_0|^2 + h_{ex} \int_{\Omega} \nabla \xi_0 \cdot \nabla \zeta \\ &= J_0 + \int_{\Omega} |\Delta \zeta|^2. \end{aligned}$$

Proof. – Using the decomposition (2.6),

$$\begin{aligned} &\int_{\Omega} (|h - h_{ex}|^2 - h_{ex}^2) \\ &= \int_{\Omega} |h_{ex} \Delta \xi_0 + \Delta \zeta|^2 - 2h_{ex} (h_{ex} \Delta \xi_0 + \Delta \zeta) \end{aligned}$$

$$\begin{aligned}
 &= h_{ex}^2 \int_{\Omega} |\Delta \xi_0|^2 + \int_{\Omega} |\Delta \zeta|^2 - 2h_{ex}^2 \int_{\Omega} \Delta \xi_0 + 2h_{ex} \int_{\Omega} \Delta \xi_0 \Delta \zeta \\
 &\quad - 2h_{ex} \int_{\Omega} \Delta \zeta.
 \end{aligned}$$

But, from [13], Section 4 (using (2.4)),

$$\int_{\Omega} |\nabla \xi_0|^2 + |\Delta \xi_0|^2 = \int_{\Omega} \Delta \xi_0,$$

and

$$\int_{\Omega} \nabla \zeta \cdot \nabla \xi_0 + \Delta \zeta \Delta \xi_0 = \int_{\Omega} \Delta \zeta.$$

Therefore,

$$\begin{aligned}
 &\frac{1}{2} \int_{\Omega} (|h - h_{ex}|^2 - h_{ex}^2) + \frac{1}{2} h_{ex}^2 \int_{\Omega} |\nabla \xi_0|^2 + h_{ex} \int_{\Omega} \nabla \xi_0 \cdot \nabla \zeta \\
 &= -\frac{h_{ex}^2}{2} \int_{\Omega} |\nabla \xi_0|^2 + |\Delta \xi_0|^2 + \int_{\Omega} |\Delta \zeta|^2 \\
 &= J_0 + \int_{\Omega} |\Delta \zeta|^2.
 \end{aligned}$$

This completes the proof. \square

Combining Lemmas 2.2 and 2.4, we obtain the following expansion of the energy:

$$\begin{aligned}
 J(u, A) &= \frac{1}{2} \int_{\Omega} |\nabla u - iu \nabla^{\perp} \zeta|^2 + \frac{1}{2\epsilon^2} (1 - |u|^2)^2 + \int_{\Omega} |\Delta \zeta|^2 \\
 &\quad + J_0 + 2\pi h_{ex} \sum_{i \in I} d_i \xi_0(a_i) + o(1). \tag{2.24}
 \end{aligned}$$

Notice that this expansion is quite similar to that of [13], but the terms in ζ are treated differently and gathered in positive expressions.

We need a last lower bound:

LEMMA 2.5. –

$$\int_{\Omega} |\nabla u - iu \nabla^{\perp} \zeta|^2 \geq \sum_{i \in I} \int_{B_i} |\nabla u|^2 + o(1).$$

Proof. – Indeed,

$$\begin{aligned} \int_{\Omega} |\nabla u - iu \nabla^\perp \zeta|^2 &\geq \int_{\bigcup_{i \in I} B_i} |\nabla u - iu \nabla^\perp \zeta|^2 \\ &\geq \int_{\bigcup_{i \in I} B_i} |\nabla u|^2 + |u \nabla^\perp \zeta|^2 - 2(\nabla u, iu \nabla^\perp \zeta). \end{aligned}$$

But,

$$\begin{aligned} \left| \int_{\bigcup_{i \in I} B_i} (\nabla u, iu \nabla^\perp \zeta) \right| &\leq (\text{Card } I) \|\nabla \zeta\|_{L^\infty} \|\nabla u\|_{L^2} \max_i r_i \\ &\leq C |\log \varepsilon|^{4-\alpha} = o(1), \end{aligned}$$

similarly to the beginning of the proof of Lemma 2.3. Hence,

$$\int_{\Omega} |\nabla u - iu \nabla^\perp \zeta|^2 \geq \int_{\bigcup_{i \in I} B_i} |\nabla u|^2 + o(1). \quad \square$$

From this lemma, we deduce that

$$\begin{aligned} &\frac{1}{2} \int_{\Omega} |\nabla u - iu \nabla^\perp \zeta|^2 + \frac{1}{2\varepsilon^2} (1 - |u|^2)^2 \\ &\geq \sum_{i \in I} \int_{B_i} \frac{1}{2} |\nabla u|^2 + \frac{1}{4\varepsilon^2} (1 - |u|^2)^2 \\ &\geq \sum_{i \in I} F(u, B_i), \end{aligned} \tag{2.25}$$

and this last expression can be bounded from below by (2.12).

2.3. End of the proof of the theorem

Considering our minimizing solution, we deduce from (2.24), Lemma 2.5, and (2.12), that

$$\begin{aligned} J(u, A) &\geq J_0 + 2\pi h_{ex} \sum_{i \in I} d_i \xi_0(a_i) \\ &\quad + \pi \sum_{i \in I} |d_i| (|\log \varepsilon| + O(|\log |\log \varepsilon||)). \end{aligned} \tag{2.26}$$

On the other hand, by minimality,

$$J(u, A) \leq J_0,$$

thus, as ξ_0 is negative,

$$\begin{aligned} \pi \sum_{i \in I} |d_i| (|\log \varepsilon| + O(|\log |\log \varepsilon||)) &\leq 2\pi h_{ex} \sum_{i \in I} d_i |\xi_0(a_i)| \\ &\leq 2\pi h_{ex} \left(\sum_{i \in I} |d_i| \right) \max |\xi_0|. \end{aligned}$$

If $\sum_{i \in I} |d_i| \neq 0$, we deduce that

$$h_{ex} \geq \frac{1}{2 \max |\xi_0|} (|\log \varepsilon| + O(|\log |\log \varepsilon||)) := H'_{c_1}.$$

But, in [13], $H_{c_1} = k_1 |\log \varepsilon| + O(1)$ with

$$k_1 = \frac{1}{2 \max |\xi_0|},$$

thus

$$H'_{c_1} = H_{c_1} + O(|\log |\log \varepsilon||). \tag{2.27}$$

Consequently, if $h_{ex} < H'_{c_1}$, we must have $d_i = 0, \forall i \in I$. Then, with (2.24) and Lemma 2.5,

$$J_0 \geq J(u, A) \geq \sum_{i \in I} F_\varepsilon(u, B_i) + J_0 + o(1),$$

implying

$$\sum_{i \in I} F_\varepsilon(u, B_i) \leq o(1). \tag{2.28}$$

We conclude that $|u| \geq 3/4$ in Ω . Indeed, it is well known from [3] that if $|u(x_0)| < 3/4$, there exist constants $\lambda, \mu > 0$ such that

$$\forall \varepsilon > 0, \quad \frac{1}{\varepsilon^2} \int_{B(x_0, \lambda \varepsilon)} (1 - |u|^2)^2 > \mu > 0, \tag{2.29}$$

contradicting (2.28).

Knowing that u is vortex-less, one may re-use the computations of [13] to find that

$$J(u, A) = J_0 + F(u) + \tilde{V}(\zeta) + o(1),$$

where $\tilde{V}(\zeta) \geq 0$. Hence $F(u) \leq o(1)$, and by definition $(u, A) \in D$. This proves that for $h_{ex} < H'_{c_1}$, (u, A) coincides with the unique Meissner solution found in [13] and [15].

The theorem is proved.

3. CONSTRUCTION OF THE BALLS

In this section, we use the method of R. Jerrard introduced in [7], in order to construct balls containing all the zeros of u , on which we have a suitable lower bound on F_ε of the order $\pi|d||\log \varepsilon|$. The size of the balls has to be large enough so that most of the energy F_ε is concentrated in these balls, but it has to be smaller than $|\log \varepsilon|^{-5}$ as we saw in Section 2. We follow almost readily the proofs of [7].

3.1. Main steps

First, we include the set $\{x/|u(x)| < 3/4\}$ in well-chosen disjoint “small” balls B_i of radii $r_i > \varepsilon$ such that

$$F_\varepsilon(u, B_i) \geq \frac{C_1 r_i}{\varepsilon},$$

where C_1 is a constant. This is possible according to the following lemma, adjusted from [7]:

LEMMA 3.1. – *Let $u : \Omega \rightarrow \mathbb{C}$ be such that $|\nabla u| < C/\varepsilon$. Then there exist disjoint balls B_1, \dots, B_k of radii r_i such that*

- (1) $\forall 1 \leq i \leq k, r_i > \varepsilon,$
- (2) $\{|u| < 3/4\} \subset \bigcup_i B_i$ and $\forall 1 \leq i \leq k, B_i \cap \{|u| < 3/4\} \neq \emptyset,$
- (3) $\forall 1 \leq i \leq k,$

$$F_\varepsilon(u, B_i \cap \Omega) \geq \frac{C_1 r_i}{\varepsilon}.$$

Then the proof involves dilating the balls B_i into balls B'_i . A lower bound for $F_\varepsilon(u, B'_i)$ is obtained by combining the lower bound for $F_\varepsilon(u, B_i)$ and a lower bound of the energy on the annulus $B'_i \setminus \bar{B}_i$:

LEMMA 3.2. – $\forall r > s > \varepsilon$, if B_r and B_s are two concentric balls of respective radii r and s , and if $u : B_r \setminus \overline{B_s} \rightarrow \mathbb{C}$ is such that $|u| > 3/4$, $d = \text{deg}(u, \partial B_r)$, then

$$F_\varepsilon(u, B_r \setminus \overline{B_s}) \geq |d| \left(\Lambda_\varepsilon \left(\frac{r}{|d|} \right) - \Lambda_\varepsilon \left(\frac{s}{|d|} \right) \right), \tag{3.1}$$

where Λ_ε is a function that satisfies the following properties:

- (1) $\Lambda_\varepsilon(s)/s$ is decreasing on \mathbb{R}_+ ,
- (2) $\sup_{s \in \mathbb{R}_+} \Lambda_\varepsilon(s)/s \leq C_1/\varepsilon$,
- (3) there exist $\varepsilon_0, t_0 > 0$ such that if $\varepsilon < \varepsilon_0$ and $\varepsilon < t < t_0$ then

$$\left| \Lambda_\varepsilon(t) - \pi \log \frac{t}{\varepsilon} \right| \leq C.$$

Thanks to this lemma, if

$$F_\varepsilon(u, B) \geq \pi |d| \log \frac{r}{\varepsilon}, \tag{3.2}$$

where $d = \text{deg}(u, \partial B)$, and r is the radius of B , if B' is the dilated ball, and if $|u| \geq 3/4$ on $B'_i \setminus \overline{B_i}$, then an estimate of the type (3.2) is still true on B' .

Thus we start with the balls B_i given by Lemma 3.1, then make them grow progressively. Say the growth rate is governed by a parameter s , we thus construct a family $\mathcal{B}(s)$ of disjoint balls. To keep this family of balls disjoint, when some of them intersect, we merge them into a larger ball of radius equal to the sum of the radii of the merged balls. If the growth-rates of the balls have been properly synchronized, then the lower bounds for the energy on each of the merged balls add up nicely so that a lower bound of the type (3.2) is still true for the larger ball. We then resume the dilation, etc., until we reach the size of balls that we wish.

The following proposition sums up the whole growth process:

PROPOSITION 3.1. – Let $u : \Omega \rightarrow \mathbb{C}$ be such that $|\nabla u| \leq C/\varepsilon$; and $\{B_i\}_i$ be a family of balls of radii r_i satisfying the results of Lemma 3.1. Let

$$d_i = \begin{cases} \text{deg}(u, \partial B_i) & \text{if } \overline{B_i} \subset \Omega, \\ 0 & \text{otherwise.} \end{cases}$$

Let also

$$s_0 = \min_{\{i|d_i \neq 0\}} \frac{r_i}{|d_i|}.$$

Then, for every $s \geq s_0$, there exists a family $\mathcal{B}(s)$ of disjoint balls $B_1(s), \dots, B_{k(s)}(s)$ of radii $r_i(s)$ such that

- (1) the family of balls is monotone, i.e., if $s < t$ then

$$\bigcup_i B_i(s) \subset \bigcup_i B_i(t),$$

- (2) for every i , $F_\varepsilon(u, B_i(s)) \geq r_i(s) \Lambda_\varepsilon(s)/s$, where Λ_ε is defined in Lemma 3.2,
- (3) if $\overline{B_i(s)} \subset \Omega$ and $d_i(s) = \deg(u, \partial B_i(s))$, then $r_i(s) \geq s|d_i(s)|$.

We then get as a consequence the following proposition, that was stated as Proposition 2.1 in Section 2, and which yields the final balls that we needed:

PROPOSITION 3.2. – Let $u : \Omega \rightarrow \mathbb{C}$ be such that $|\nabla u| \leq C/\varepsilon$ and that $F_\varepsilon(u) \leq C|\log \varepsilon|^2$. Then, for any $\alpha > 0$ there exist disjoint balls $(B_i)_{i \in I}$ of radii r_i such that, for sufficiently small ε ,

- (1) $\{x/|u(x)| < 3/4\} \subset \bigcup_{i \in I} B_i$,
- (2) $\text{Card } I \leq C|\log \varepsilon|^2$,
- (3) $r_i \leq C/|\log \varepsilon|^\alpha$,
- (4) if $\overline{B_i} \subset \Omega$ and $d_i = \deg(u, \partial B_i)$, then

$$F_\varepsilon(u, B_i) \geq \pi|d_i|(|\log \varepsilon| - O(|\log |\log \varepsilon||)). \quad (3.3)$$

Proof. – We first consider the balls given by Lemma 3.1, and then apply Proposition 3.1 to get bigger balls. We need to check that s_0 is small enough to be able to apply Proposition 3.1 for s large enough. Indeed, $s_0 = \min_{\{i|d_i \neq 0\}}(r_i/d_i)$, but from assertion (3) of Lemma 3.1,

$$C_1 r_i < \varepsilon F_\varepsilon(u, B_i \cap \Omega) \leq C\varepsilon |\log \varepsilon|^2,$$

so that $s_0 \leq C\varepsilon |\log \varepsilon|^2$. We can thus apply Proposition 3.1 for all $s \geq C\varepsilon |\log \varepsilon|^2$. We choose in particular

$$s_1 = \frac{1}{|\log \varepsilon|^{\alpha+1}}. \quad (3.4)$$

Proposition 3.1 yields final balls $\mathcal{B}(s_1)$ such that

$$\forall i, \text{ if } \overline{B_i(s_1)} \subset \Omega, \quad F_\varepsilon(u, B_i(s_1)) \geq \frac{\Lambda_\varepsilon(s_1)}{s_1} r_i(s_1), \quad (3.5)$$

with

$$r_i(s_1) \geq s_1 |d_i(s_1)|. \quad (3.6)$$

Therefore,

$$F_\varepsilon(u, B_i(s_1)) \geq \Lambda_\varepsilon(s_1) |d_i(s_1)|,$$

and from Lemma 3.2 (assertion (3)),

$$\begin{aligned} F_\varepsilon(u, B_i(s_1)) &\geq |d_i(s_1)| \left(\pi \log \frac{s_1}{\varepsilon} - C \right) \\ &\geq \pi |d_i(s_1)| (|\log \varepsilon| - O(|\log |\log \varepsilon||)). \end{aligned}$$

We thus have the lower bound (4) on F_ε .

We then show that (3) is true. We know that $F_\varepsilon(u) \leq Ch_{ex}^2 \leq C|\log \varepsilon|^2$. Combining this with (3.5), we get

$$r_i(s_1) \frac{\Lambda_\varepsilon(s_1)}{s_1} \leq C|\log \varepsilon|^2.$$

But, as $\Lambda_\varepsilon(s) \simeq \pi \log(s/\varepsilon)$, in view of (3.4),

$$\Lambda_\varepsilon(s_1) \geq \pi |\log \varepsilon| - O(|\log |\log \varepsilon||).$$

Hence, if ε is sufficiently small,

$$r_i(s_1) \leq C \frac{s_1}{\Lambda_\varepsilon(s_1)} |\log \varepsilon|^2 \leq \frac{C}{|\log \varepsilon|^{\alpha+1}} \frac{|\log \varepsilon|^2}{|\log \varepsilon|} \leq \frac{C}{|\log \varepsilon|^\alpha},$$

which is the desired estimate.

There only remains to show that (2) is true. This is easy since in Lemma 3.1, each ball satisfies $F_\varepsilon(u, B_i \cap \Omega) \geq C_1 r_i / \varepsilon$, with $r_i > \varepsilon$, hence carries an energy that is bounded from below by a constant independent from ε . As $F_\varepsilon \leq C|\log \varepsilon|^2$, we see that the number of these balls has to be bounded by $C|\log \varepsilon|^2$. Then, the procedure of Proposition 3.1 does not increase the number of balls, so that property (2) is true. \square

3.2. Proof of Lemma 3.1

We use the notation $S(x, r)$ for the circle in \mathbb{R}^2 of center x and radius r . We begin with the following lemma, taken from [7]:

LEMMA 3.3. – *Let $u : S_t \rightarrow \mathbb{C}$, where S_t is a circle of radius t in \mathbb{R}^2 such that $t > \varepsilon$. Let $m = \min_{S_t} |u|$ and, if $m \neq 0$, $d = \deg(u, S_t)$. If $m = 0$, let $d = 0$. Then, assuming $0 \leq m \leq 1$,*

$$\int_{S_t} \frac{1}{2} |\nabla u|^2 + \frac{1}{4\varepsilon^2} (1 - |u|^2)^2 \geq m^2 \frac{\pi d^2}{t} + \frac{(1 - m)^{C_2}}{C_2 \varepsilon}, \tag{3.7}$$

where C_2 is an absolute constant.

Proof. – See [7], Theorem 2.1. \square

We also have the following variant of Lemma 3.3 (Lemma 2.4 in [7]):

LEMMA 3.4. – *If $u : \Omega \rightarrow \mathbb{C}$, there exists $\rho(\Omega), C(\Omega) > 0$ such that $\forall x \in \Omega, \forall \varepsilon < r < \rho$, letting $m = \min_{S_r \cap \Omega} |u|$,*

$$F_\varepsilon(u, S_r) \geq \frac{(1 - m)^C}{C\varepsilon}. \tag{3.8}$$

Proof. – See [7]. \square

We then divide the proof of Lemma 3.1 in five steps.

Step 1. We wish to include $\{|u| < 3/4\}$ in balls. Let S_1, \dots, S_k be the connected components of $\{|u| < 4/5\}$ that intersect $\{|u| < 3/4\}$, and $x_1 \in S_1, \dots, x_k \in S_k$ be such that $|u(x_i)| < 3/4$. Also, for every i let

$$r_i = \sup \{r > 0 \mid S(x_i, r) \cap \{|u| < 4/5\} \neq \emptyset\}.$$

We claim that

$$F_\varepsilon(u, B(x_i, r_i) \cap \Omega) \geq \frac{C_1 r_i}{\varepsilon}. \tag{3.9}$$

Indeed from the hypothesis $|\nabla u| < C/\varepsilon$, we get as in (2.29) that $F_\varepsilon(u, B(x_i, r_i) \cap \Omega) \geq C$. Therefore if $r_i \leq 2\varepsilon$, the claim is true.

If $r_i > 2\varepsilon$ then, from Lemma 3.4, as $\forall t \in [\varepsilon, r_i], \min_{S(x_i, t)} |u| < 4/5$ by definition of r_i ,

$$\forall t \in [\varepsilon, r_i], \quad F_\varepsilon(u, S(x_i, t) \cap \Omega) \geq \frac{C}{\varepsilon}.$$

We deduce that

$$\begin{aligned}
 F_\varepsilon(u, \Omega \cap B(x_i, r_i)) &\geq \int_\varepsilon^{r_i} F_\varepsilon(u, \Omega \cap S(x_i, t)) dt \\
 &\geq \frac{C(r_i - \varepsilon)}{\varepsilon} \geq \frac{Cr_i}{\varepsilon},
 \end{aligned}$$

proving the claim in this case also.

Step 2. For simplicity, we write B_i for $B(x_i, r_i)$. We claim that $\forall i \neq j$, either $B_j \subset B_i$ (in this case, we drop B_j) or $x_j \notin B_i$.

Indeed, assume that $x_j \in B_i$. By definition of the balls B_i , $\partial B_i \cap \bigcup_\ell S_\ell = \emptyset$, thus

$$S_j = (S_j \cap \overline{C B_i}) \cup (S_j \cap B_i).$$

Since $x_j \in B_i$, $S_j \cap B_i \neq \emptyset$, using the connectedness of S_j , $S_j \subset B_i$, and we can drop B_j .

The claim is proved.

Step 3. Dropping the unnecessary balls, we reduce to balls B_i such that

$$\bigcup_i S_i \subset \bigcup_k B_k \quad \text{and} \quad \forall i \neq j, x_j \notin B_i.$$

It follows from the Besicovitch covering lemma (see for instance [17], p. 44), any $x \in \bigcup B_i$ belongs to at most N of the balls, where N is an absolute constant.

Step 4. Naming C_i the connected components of $\bigcup_i B_i$, this implies that

$$\begin{aligned}
 \forall i \quad F_\varepsilon(u, C_i) &\geq \frac{1}{N} \sum_{j/B_j \subset C_i} F_\varepsilon(u, B_j) \geq C \sum_{j/B_j \subset C_i} \frac{r_j}{\varepsilon} \\
 &\geq C \frac{\text{diam } C_i}{\varepsilon},
 \end{aligned} \tag{3.10}$$

where we have used (3.9).

Step 5. Each C_i can be included in a ball B'_i of radius

$$r'_i \leq \text{diam } C_i. \tag{3.11}$$

These balls B'_i can be included in bigger balls B''_i the following way: if B'_i and B'_j intersect, we merge them into a ball of radius no higher than $r'_i + r'_j$, etc., until all intersecting balls are merged. Hence, we are left

with a family B_i'' of disjoint balls of radii r_i'' . This family satisfies

$$r_i'' \leq \sum_{j/B_j' \subset B_i''} r_j'. \tag{3.12}$$

As the C_i are disjoint, we have

$$F_\varepsilon(u, B_i'') \geq \sum_{j/C_j \subset B_i''} F_\varepsilon(u, C_j) \geq \sum_{j/C_j \subset B_i''} C \frac{\text{diam } C_j}{\varepsilon},$$

with (3.10). Using (3.11) and (3.12), we are led to

$$F_\varepsilon(u, B_i'') \geq \sum_{j/C_j \subset B_i''} C \frac{r_j'}{\varepsilon} \geq C \frac{r_i''}{\varepsilon}.$$

The family (B_i'') satisfies the desired conditions, hence the proof is complete.

3.3. Proof of Lemma 3.2

Under the assumptions of Lemma 3.3,

$$F_\varepsilon(u, S_t) \geq f_\varepsilon\left(\frac{t}{|d|}\right), \tag{3.13}$$

where

$$f_\varepsilon(s) = \frac{ab}{(a^q + b^q)^{1/q}}, \quad a(s) = \pi/s, \quad b(s) = \frac{C_2}{\varepsilon},$$

and $0 \leq q \leq 1$ is a constant.

Indeed, using Lemma 3.3, we can bound the right-hand side of (3.7) from below by

$$(m^{C'}) \frac{\pi|d|}{t} + \frac{C_2(1-m)^{C'}}{\varepsilon},$$

where $C' = \max(2, C_2) \geq 2$. This uses the fact that $0 < m < 1$. Then we minimize with respect to $m \in [0, 1]$. This yields (3.9) with $q = 1/(C' - 1)$.

We then define

$$\lambda_\varepsilon(s) = \min\left(f_\varepsilon(s), \frac{C_1}{\varepsilon}\right), \tag{3.14}$$

where C_1 is defined in Lemma 3.1 Also, we let

$$\Lambda_\varepsilon(t) = \int_0^t \lambda_\varepsilon(s) ds. \tag{3.15}$$

We prove the properties on Λ_ε . (1) is true because Λ_ε is the primitive of λ_ε which is easily seen to be decreasing. From this, we deduce that

$$\sup_{s \in \mathbb{R}_+} \frac{\Lambda_\varepsilon(s)}{s} = \lambda_\varepsilon(0) \leq \frac{C_1}{\varepsilon},$$

so that (2) is true. There remains to prove assertion (3). Recall that

$$\Lambda_\varepsilon(s) = \int_0^s \frac{a}{((a/b)^q + 1)^{1/q}} dt, \quad \text{where } a = \frac{\pi}{t}, \quad b = \frac{C_2}{\varepsilon}.$$

Denoting by C, C' generic positive constants, if C is large enough $1 \geq (1 + x^q)^{-1/q} \geq 1 - Cx$, whenever $x < 1/C$. On the other hand, it is easy to check that if $t > C'\varepsilon$, with C' large enough, then $a/b < 1/C$. Thus we may write

$$\int_{C'\varepsilon}^s a \left(1 - C \frac{a}{b}\right) dt \leq \int_{C'\varepsilon}^s \frac{a}{((a/b)^q + 1)^{1/q}} dt \leq \int_{C'\varepsilon}^s a dt.$$

It is easy to check that

$$\int_{C'\varepsilon}^s \frac{a^2}{b} dt < C,$$

and that

$$\left| \int_{C'\varepsilon}^s a dt - \pi \log \frac{s}{\varepsilon} \right| < C, \tag{3.16}$$

if $s < t_0$. Finally, combining (3.16) with the fact that

$$\int_0^{C'\varepsilon} \lambda_\varepsilon(t) dt < C,$$

the assertion is proved.

From (3.13), (3.14), and (3.15), we deduce that $\forall r > s > \varepsilon$,

$$\begin{aligned} F_\varepsilon(u, B_r) - F_\varepsilon(u, B_s) &\geq \int_s^r F_\varepsilon(u, S_t) dt \geq \int_s^r \lambda_\varepsilon\left(\frac{t}{|d|}\right) dt \\ &\geq \Lambda_\varepsilon\left(\frac{r}{|d|}\right) - \Lambda_\varepsilon\left(\frac{s}{|d|}\right). \end{aligned}$$

The lemma is proved.

3.4. Proof of Proposition 3.1

First of all, letting $\mathcal{B}(s_0)$ be the family of balls given by Lemma 3.3, we check that properties (2), (3) are verified for $s = s_0$. Indeed, from Lemma 3.1,

$$F_\varepsilon(u, B_i \cap \Omega) \geq \frac{C_1 r_i}{\varepsilon}$$

while

$$\frac{\Lambda_\varepsilon(s)}{s} \leq \frac{C_1}{\varepsilon}.$$

Thus, the second property is true. Property (3) results directly from the definition of s_0 . Notice that $\{|u| < 3/4\}$ is contained in the initial family $\mathcal{B}(s_0)$.

Now we let I be the largest interval containing s_0 such that $\forall s \in I$ there exists a finite family of disjoint balls $\mathcal{B}(s)$ verifying properties (1)–(3) above. We already know I is not empty. We now prove that I is open.

Suppose $[s_0, s_1] \subset I$. We wish to define a family $\mathcal{B}(t)$ for $t \in [s_1, s_1 + \alpha]$, for some $\alpha > 0$. Three cases occur.

Case 1. For the family $\mathcal{B}(s_1)$, all the inequalities in (3) are *strict*, i.e., $r_i(s_1) > s_1 |d_i(s_1)|$, $\forall i$ such that $\overline{B_i} \subset \Omega$. In this case we let $B_i(t) = B_i(s_1)$, i.e., we do not change the balls. This defines a family $\mathcal{B}(t)$ of disjoint balls that verifies (1) trivially, (2) also since—by Lemma 3.2— $\Lambda_\varepsilon(t)/t \leq \Lambda_\varepsilon(s_1)/s_1$ for $t \geq s_1$. Finally (3) is verified at least when $t \in [s_1, s_1 + \alpha]$, $\alpha > 0$ not too large.

Case 2. There is equality in (3) for, say, balls $B_1(s_1), \dots, B_\ell(s_1)$, but *all* the balls have disjoint closures. In this case the family $\mathcal{B}(t)$ for $t > s_1$ is defined by:

- The ball $B_i(t)$, $1 \leq i \leq \ell$ has the same center as $B_i(s_1)$ and its radius $r_i(t)$ is such that $r_i(t) = t |d_i(t)| = t |d_i(s_1)|$, as long as $\overline{B_i(t)} \subset \Omega$,

which is true when $t \in [s_1, s_1 + \alpha]$, $\alpha > 0$ not too large. (The degree remains constant as $|u| > 3/4$ on $\mathcal{B}(t) \setminus \mathcal{B}(s_0)$, $\forall t > s_0$.) In particular the balls are *growing*.

- The other balls remain unchanged.

The balls defined this way remain disjoint for $t > s_1$ not too large. Property (1) obviously is still true, (3) also. Property (2) remains true in an obvious way for the static balls, as $\Lambda_\varepsilon(s)/s$ is decreasing. For the increasing ones, we just use Lemma 3.2, which yields

$$\begin{aligned} F_\varepsilon(u, B_i(t)) &\geq F_\varepsilon(u, B_i(s_1)) + F_\varepsilon(u, B_i(t) \setminus \overline{B_i(s_1)}) \\ &\geq r_i(s_1) \frac{\Lambda_\varepsilon(s_1)}{s_1} + |d_i(t)| \left(\Lambda_\varepsilon \left(\frac{r_i(t)}{|d_i(t)|} \right) - \Lambda_\varepsilon \left(\frac{r_i(s_1)}{|d_i(s_1)|} \right) \right). \end{aligned}$$

But, we know that for the growing balls, we have $r_i(s_1) = s_1 |d_i(s_1)|$, and $r_i(t) = t |d_i(t)|$. Hence, with the constancy of the degree,

$$\begin{aligned} F_\varepsilon(u, B_i(t)) &\geq r_i(s_1) \frac{\Lambda_\varepsilon(s_1)}{s_1} + \frac{r_i(t)}{t} \Lambda_\varepsilon(t) - \frac{r_i(s_1)}{s_1} \Lambda_\varepsilon(s_1) \\ &\geq r_i(t) \frac{\Lambda_\varepsilon(t)}{t}. \end{aligned}$$

Case 3. There is equality in (3) for some of the balls in $\mathcal{B}(s_1)$ but, say, $\overline{B_1(s_1)} \cap \overline{B_2(s_1)} \neq \emptyset$. Then we modify the family $\mathcal{B}(s_1)$ as follows: group $B_1(s_1)$ and $B_2(s_1)$ into a single larger ball B of radius $r_1(s_1) + r_2(s_1)$. If B intersects, say, $\overline{B_3}$, enlarge it so that $B_1 \cup B_2 \cup B_3 \subset B$ and the radius of B is $r = r_1 + r_2 + r_3$, etc.

Thus, we have a new family of balls $\mathcal{B}'(s_1)$ whose closures are disjoint, such that the union of the balls in this family contains the balls of $\mathcal{B}(s_1)$, that verifies properties (2) and (3). This last statement is obviously true for the balls that have not changed in the process. We verify it for a ball $B \in \mathcal{B}'(s_1)$ that results from grouping $B_1, \dots, B_\ell \in \mathcal{B}(s_1)$.

- Property (2) is verified since

$$F_\varepsilon(u, B) \geq \sum_{i=1}^{\ell} F_\varepsilon(u, B_i) \geq \left(\sum_{i=1}^{\ell} r_i \right) \frac{\Lambda_\varepsilon(s_1)}{s_1} = r \frac{\Lambda_\varepsilon(s_1)}{s_1}.$$

- If $\bar{B} \subset \Omega$ and $\deg(u, \partial B) = d \neq 0$ then $d = \sum_i d_i$, where $d_i = \deg(u, \partial B_i)$. Thus

$$\frac{r}{|d|} \geq \frac{r}{\sum_{\{i|d_i \neq 0\}} |d_i|} \geq s_1,$$

since for every i such that $d_i \neq 0$, $r_i \geq s_1 |d_i|$. Thus property (3) is verified. If $d = 0$, then the property is empty.

Now, to define $\mathcal{B}(t)$ for $t > s_1$ we start from $\mathcal{B}'(s_1)$, and use case 1 or case 2.

It remains to prove that the interval I is closed. Indeed, suppose $[s_0, s_1) \subset I$ and that $\mathcal{B}(s)$ is defined on this interval by the above procedure. We wish to define $\mathcal{B}(s_1)$. First note that since the number of balls in $\mathcal{B}(s)$ is nonincreasing, and since there is initially a finite number of them, the number of balls is constant on some interval $A = [s_1 - \alpha, s_1)$. On this interval, the balls $B_1(s), \dots, B_\ell(s)$ are well defined, have their centers fixed, their radius increases continuously with s , and their degree $d_i(s) = \deg(u, \partial B_i(s))$ is constant for $s \in A$. Then we let $B_i(s_1) = \bigcup_{s \in A} B_i(s)$. This defines a family $\mathcal{B}(s_1)$ of disjoint balls that are easily seen to verify properties (1), (2). For (3), note that $\deg(u, \partial B_i(s_1))$ is either equal to $d_i(s)$, $s \in A$ if $\overline{B_i(s_1)} \subset \Omega$ or to 0. In any case, (3) will be verified.

The proposition is proved. This completes all the proofs.

REFERENCES

- [1] A. ABRIKOSOV, On the magnetic properties of superconductors of the second type, *Soviet Phys. JETP* 5 (1957) 1174–1182.
- [2] L. ALMEIDA and F. BETHUEL, Topological methods for the Ginzburg–Landau equations, *J. Math. Pures Appl.* 77 (1998) 1–49.
- [3] F. BETHUEL, H. BREZIS and F. HÉLEIN, *Ginzburg–Landau Vortices*, Birkhäuser, 1994.
- [4] F. BETHUEL and T. RIVIÈRE, Vorticit  dans les mod les de Ginzburg–Landau pour la supraconductivit , *S minaire E. D. P de l’ cole Polytechnique XVI* (1994).
- [5] F. BETHUEL and T. RIVIÈRE, Vortices for a variational problem related to superconductivity, *Annales IHP, Analyse non Lin aire* 12 (1995) 243–303.
- [6] P.G. DEGENNES, *Superconductivity of Metal and Alloys*, Benjamin, New York and Amsterdam, 1966.
- [7] R. JERRARD, Lower bounds for generalized Ginzburg–Landau Functionals, *SIAM J. Math. Anal.* 30 (4) (1999) 721–746.
- [8] J. RUBINSTEIN, Six lectures on superconductivity, in: *Proc. CRM School on “Boundaries, Interfaces, and Transitions”*.

- [9] D. SAINT-JAMES, G. SARMA and E.J. THOMAS, *Type-II Superconductivity*, Pergamon Press, 1969.
- [10] E. SANDIER, Lower bounds for the energy of unit vector fields and applications, *J. Functional Analysis* 152 (2) (1998) 379–403.
- [11] E. SANDIER and S. SERFATY, On the energy of type-II superconductors in the mixed phase, Preprint.
- [12] S. SERFATY, Solutions stables de l'équation de Ginzburg–Landau en présence de champ magnétique, *C. R. A. S. I* 326 (8) (1998) 955.
- [13] S. SERFATY, Local minimizers for the Ginzburg–Landau energy near critical magnetic field, Part I, *Comm. Contemporary Mathematics*, to appear.
- [14] S. SERFATY, Local minimizers for the Ginzburg–Landau energy near critical magnetic field, Part II, *Comm. Contemporary Mathematics*, to appear.
- [15] S. SERFATY, Stable configurations in superconductivity: Uniqueness, multiplicity and vortex-nucleation, *Arch. for Rat. Mech. Anal.*, to appear.
- [16] S. SERFATY, Sur l'équation de Ginzburg–Landau avec champ magnétique, in: *Proc. "Journées Équations aux dérivées partielles, Saint-Jean-de-Monts"*, 1998.
- [17] E.M. STEIN, *Harmonic Analysis*, Princeton Math. Series, No 43, Princeton University Press, 1969.
- [18] D. TILLEY and J. TILLEY, *Superfluidity and Superconductivity*, 2nd ed., Adam Hilger, Bristol, 1986.