




ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Ann. I. H. Poincaré – AN 21 (2004) 187–207



www.elsevier.com/locate/anihpc

On planar selfdual Electroweak vortices

Sur les vortex Électrofaibles autoduaux dans le plan

Dongho Chae^a, Gabriella Tarantello^{b,*}

^a Department of Mathematics, Seoul National University, Seoul 151-742, South Korea

^b Dipartimento di Matematica, Università degli Studi di Roma "Tor Vergata", Via della Ricerca Scientifica, 00133 Roma, Italy

Received 29 May 2002; accepted 30 January 2003

Abstract

By perturbation techniques, we obtain a new class of selfdual Electroweak vortices in \mathbb{R}^2 , whose asymptotic behavior we control in an "optimal" way to yield a sort of "quantization" property for the corresponding flux.

Our class of vortex-solutions complements that constructed by Spruck and Yang [Comm. Math. Phys. 144 (1992) 215–234]. © 2003 Elsevier SAS. All rights reserved.

Résumé

En utilisant des techniques de perturbation, nous obtenons une nouvelle classe de vortex Électrofaibles autoduaux dans \mathbb{R}^2 , dont nous pouvons contrôler de façon « optimale » le comportement asymptotique en donnant une propriété de « quantisation » pour le flot correspondant.

Notre classe de solutions complète celles construites par Spruck et Yang [Comm. Math. Phys. 144 (1992) 215–234]. © 2003 Elsevier SAS. All rights reserved.

MSC: 35J45; 35J60; 37K40; 70S15

Introduction

We shall be concerned with planar vortex-type (self-dual) solutions for the celebrated $SU(2) \times U(1)$ -Electroweak theory of Glashow, Salam and Weinberg [10]. As observed by Ambjorn and Olesen [2–4], if the physical parameters satisfy a suitable critical condition, it is possible to determine Bogomol'nyi type equations (also known as self-dual equations) to be satisfied by Electroweak-vortices when expressed in terms of the unitary gauge variables. The selfdual equations include a gauge invariant version of the Cauchy–Riemann equation, which makes it reasonable to assume that the W -boson field admits a finite number N of zeroes (counted with

* Corresponding author.

E-mail addresses: dhchae@math.snu.ac.kr (D. Chae), tarantel@mat.uniroma2.it (G. Tarantello).

multiplicity). Such an integer N is called the vortex number, and the corresponding vanishing points of W are called the vortex points.

Moreover, by means of an approach introduced by Taubes [15,16] in the context of Ginzburg–Landau vortices, it is possible to reduce the analysis of self-dual Electroweak vortices to the study of the following elliptic system:

$$(\mathcal{P}) \quad \begin{cases} -\Delta u_1 = 4g^2 e^{u_1} + g^2 e^{u_2} - 4\pi \sum_{k=1}^m n_k \delta(z - z_k), \\ \Delta u_2 = \frac{g^2}{2 \cos^2 \theta} (e^{u_2} - \phi_0^2) + 2g^2 e^{u_1} \end{cases}$$

where ϕ_0 is a given positive parameter, g is the $SU(2)$ -coupling constant, and $\theta \in (0, \pi/2)$ is the so called “Weinberg-angle”, related with the $U(1)$ -coupling constant g_* via the relation,

$$\cos \theta = \frac{g}{(g^2 + g_*^2)^{1/2}}.$$

We refer to the recent monograph of Y. Yang [17] and [5] for details, and, in particular, on how to recover the full vortex configuration out of solutions of (\mathcal{P}) . We only mention that, by virtue of the vortex ansatz and accordingly to the unitary gauge variables, the vortex solution is specified by the (complex valued) massive field W , the (scalar) field φ and the real valued 2-vector fields $P = (P_\mu)_{\mu=1,2}$ and $Z = (Z_\mu)_{\mu=1,2}$. So that, u_1 and u_2 determine the magnitude of W and φ respectively, as follows:

$$\varphi^2 = e^{u_2}, \quad |W|^2 = e^{u_1}. \quad (0.1)$$

Thus, z_k corresponds to a vortex point, the integer $n_k \in \mathbb{N}$ to its multiplicity $k = 1, \dots, m$, and $N = \sum_{k=1}^m n_k$ is the vortex number.

Furthermore, while the 2-vector fields P and Z are defined only up to gauge transformations, their gauge invariant curls,

$$P_{12} = \partial_1 P_2 - \partial_2 P_1 \quad \text{and} \quad Z_{12} = \partial_1 Z_2 - \partial_2 Z_1$$

are determined by the relations:

$$\begin{aligned} P_{12} &= \frac{g}{2 \sin \theta} \phi_0^2 + 2g \sin \theta |W|^2, \\ Z_{12} &= \frac{g}{2 \cos \theta} (\varphi^2 - \phi_0^2) + 2g \cos \theta |W|^2 \end{aligned} \quad (0.2)$$

(see [17]). Following the numerical evidence provided by Ambjorn and Olesen, the first rigorous results concerning Electroweak-vortices have been obtained by Spruck and Yang in [13,14]. In [13], the authors aim to obtain configurations of the type described in physics literature as the “Abrikosov’s mixed states” (see [1]), so they consider the selfdual equations subject to ’t Hooft boundary conditions [9]. In turn, this amounts to solve (\mathcal{P}) under periodic boundary conditions, and Spruck and Yang in [13] obtain necessary and sufficient conditions on the given physical parameters, in order to ensure the presence of periodic N -vortices in the theory. Their results are sharp for $N = 1, 2$ and recently have been improved by Bartolucci and Tarantello in [5] to a wider range of parameters.

In this paper, we shall be concerned with planar vortex-type configurations, and consider (\mathcal{P}) over \mathbb{R}^2 subject to appropriate decay assumptions at infinity. This situation has been considered in [14], where it has been observed that necessarily, the corresponding vortex-solutions must carry infinite energy. As a consequence, solutions of (\mathcal{P}) over \mathbb{R}^2 appear in abundance, and are distinguished according to their rate of decay at infinity.

In view of (0.1), also the flux of the fields P and Z is infinite, while this may not be necessarily the case for the field: $(\sin \theta)P + (\cos \theta)Z$, which bares informations about the gauge potential in the original variables. We shall be concerned with such “finite-flux” selfdual solutions, by solving (\mathcal{P}) under the condition:

$$e^{u_1}, e^{u_2} \in L^1(\mathbb{R}^2). \quad (0.3)$$

In fact, by setting:

$$\sigma_1 = \frac{2g^2}{\pi} \int_{\mathbb{R}^2} e^{u_1}, \quad \sigma_2 = \frac{g^2}{2\pi} \int_{\mathbb{R}^2} e^{u_2} \tag{0.4}$$

in view of (0.1), (0.2) we see that the flux Φ of the field $(\sin\theta)P + (\cos\theta)Z$ is given by,

$$\Phi = \int_{\mathbb{R}^2} (\sin\theta P_{12} + \cos\theta Z_{12}) = \frac{\pi}{g}(\sigma_1 + \sigma_2). \tag{0.5}$$

As a first fact, we show that necessarily

$$\sigma_1 + \sigma_2 > 2(N + 1) \tag{0.6}$$

as σ_1 and σ_2 determine the rate of decay at infinity of u_1 and u_2 (see Theorem 1.1).

The existence results for (P)–(0.3) derived by Spruck and Yang in [14] imply that the lower bound (0.6) is “sharp”, as they construct solutions satisfying:

$$\sigma_2 > 2N; \quad \sigma_1 + \sigma_2 \text{ any assigned value in } (2(N + 1), +\infty).$$

Notice that, in contrast to the periodic case, this class of solutions lacks any sort of “quantization” property for the corresponding flux (0.5). A first goal of this paper is to show that, actually, this is no longer the case, if we consider solutions of (P)–(0.3) with $\sigma_2 \leq 2N$.

In this case, and when all vortex points coincide, we show that the value σ_2 also controls from above the value σ_1 . This implies that, in certain regime, (e.g., σ_2 small) the lower bound $2(N + 1)$ in (0.6) is no longer sharp, as $\sigma_1 + \sigma_2$ is now forced to remain close to the value $4(N + 1)$, and any other value is ruled out. See Lemma 1.6.

Thus, for σ_2 small, a sort of “quantization” property is restored as σ_1 “approximately” equals to $4(N + 1)$.

Our second goal, is to show that, in fact, such situation does occur. So, in Section 2, we shall focus our attention in constructing several families of solutions of (P)–(0.3) so that

$$\sigma_2 = o(1) \quad \text{and} \quad \sigma_1 = 4(N + 1) + o(1). \tag{0.7}$$

To this end, we provide a priori estimates (see Theorem 1.1), so that the L^1 -norm of e^{u_2} uniformly estimates its L^∞ -norm. Thus, requiring a small σ_2 , implies that we must have e^{u_2} also small in L^∞ -norm. Therefore, the first equation in (P) may be viewed as a “perturbation” of the following singular Liouville equation:

$$\begin{cases} -\Delta u = 4g^2 e^u - 4\pi \sum_{k=1}^m n_k \delta(z - z_k) & \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} e^u < +\infty, \end{cases}$$

whose solutions may be classified according to Liouville formula (e.g., see [12]). When all vortex points coincide, we are able to adapt a perturbation approach introduced by Chae and Imanuvilov in [7] to construct “non-topological” Chern–Simons vortices, and obtain solutions of (P), “bifurcating” out of solutions of the singular Liouville equation mentioned above. In this way, we are able to provide rather accurate pointwise estimates on our solutions to ensure (0.7), together with a sharp control on their decay rate at infinity. We refer to Theorem 2.1 and 2.2 for the precise statements. Clearly, our result furnishes a new class of solutions for (P)–(0.3) which complements those constructed in [14].

1. Asymptotic behavior

Consider the problem,

$$(P) \begin{cases} -\Delta u_1 = 4g^2 e^{u_1} + g^2 e^{u_2} - 4\pi \sum_{k=1}^m n_k \delta(z - z_k), \\ \Delta u_2 = \frac{g^2}{2 \cos^2 \theta} (e^{u_2} - \phi_0^2) + 2g^2 e^{u_1} \quad \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} e^{u_1} < +\infty, \quad \int_{\mathbb{R}^2} e^{u_2} < +\infty, \end{cases}$$

where $\{z_1, \dots, z_m\} \subset \mathbb{R}^2$ are given points, $\{n_1, \dots, n_m\} \subset \mathbb{N}$ are given integers, and $\delta(z - p)$ denotes the Dirac measure with pole at $p \in \mathbb{R}^2$. Let

$$N = \sum_{k=1}^m n_k, \quad c_0 = \frac{g^2 \phi_0^2}{8 \cos^2 \theta}$$

with g, ϕ_0 given positive constants and $\theta \in (0, \pi/2)$. Set

$$\sigma_1 = \frac{2g^2}{\pi} \int_{\mathbb{R}^2} e^{u_1}, \quad \sigma_2 = \frac{g^2}{2\pi} \int_{\mathbb{R}^2} e^{u_2}. \tag{1.1}$$

We devote this section to obtain the asymptotic behavior, as $|x| \rightarrow +\infty$, of a solution pair (u_1, u_2) for (P) in terms of σ_1 and σ_2 , and to establish some interesting relation for such values. We have:

Theorem 1.1. *Let (u_1, u_2) be a solution pair for (P) , then*

(i) $u_1^+ \in L^\infty(\mathbb{R}^2)$,

$$u_1(x) \geq (2N - (\sigma_1 + \sigma_2)) \ln |x| - C, \quad \text{as } |x| \rightarrow +\infty \tag{1.2}$$

for suitable $C > 0$,

$$\frac{u_1(x)}{\ln |x|} \rightarrow 2N - (\sigma_1 + \sigma_2), \quad \text{as } |x| \rightarrow +\infty, \tag{1.3}$$

and so necessarily

$$\sigma_1 + \sigma_2 > 2(N + 1). \tag{1.4}$$

(ii) $u_2^+ \in L^\infty(\mathbb{R}^2)$ and for suitable $C_0 > 0$ (depending on c_0 only)

$$\|e^{u_2}\|_{L^\infty(\mathbb{R}^2)} \leq C_0 \|e^{u_2}\|_{L^1(\mathbb{R}^2)}. \tag{1.5}$$

In addition, if

$$\frac{u_2(x) + c_0|x - x_0|^2}{1 + |x|^{1+\alpha/2}} \in L^2(\mathbb{R}^2) \tag{1.6}$$

for some $x_0 \in \mathbb{R}^2$ and $\alpha \in (0, 1)$, then

$$u_1(x) = (2N - (\sigma_1 + \sigma_2)) \ln |x| + O(1), \tag{1.7}$$

$$u_2(x) = -c_0|x - x_0|^2 + \frac{1}{2} \left(\sigma_1 + \frac{\sigma_2}{\cos^2 \theta} \right) \ln |x| + O(1) \tag{1.8}$$

as $|x| \rightarrow +\infty$.

In order to derive Theorem 1.1 we are going to recall some basically well known facts about the function:

$$w(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} (\ln|x - y| - \ln(1 + |y|)) f(y) dy \tag{1.9}$$

with $f \in L^1(\mathbb{R}^2)$.

Lemma 1.1. *Let*

$$f((\ln|f|)^+ + 1) \in L^1(\mathbb{R}^2) \quad \text{and} \quad \beta = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(y) dy \tag{1.10}$$

(i) *if* $f \geq 0$, *then*

$$w(x) \leq \beta \ln(|x| + 1) + C \tag{1.11}$$

for suitable $C > 0$;

(ii) $\frac{w(x)}{\ln|x|} \rightarrow \beta, \quad \text{as } |x| \rightarrow +\infty;$ (1.12)

(iii) *if*

$$\ln(1 + |x|)f \in L^1(\mathbb{R}^2), \tag{1.13}$$

then

$$w(x) = \beta \ln|x| + O(1), \quad \text{as } |x| \rightarrow +\infty. \tag{1.14}$$

We notice that properties (1.11), (1.12) and (1.14) remain valid for any other solution v of the equation:

$$\Delta v = f \tag{1.15}$$

provided it admits a “slow” growth at infinity as expressed by one of the following conditions:

$$\frac{v^+}{\ln|x| + 1} \in L^\infty(\mathbb{R}^2), \tag{1.16}$$

or

$$\frac{v}{1 + |x|^{1+\alpha/2}} \in L^2(\mathbb{R}^2), \quad \text{for some } \alpha \in (0, 1). \tag{1.17}$$

Corollary 1.1. *Assume (1.10), and let* $v \in L^1_{\text{loc}}(\mathbb{R}^2)$ *be a solution of (1.15) which satisfies (1.16) or (1.17). Then, properties (i), (ii) and (iii) of Lemma 1.2 hold for* v .

Proof of Corollary 1.1. Notice that the function $u = v - w$ is harmonic over \mathbb{R}^2 and, in view of Lemma 1.2, satisfies (1.16) or (1.17). In either case u must be constant. Indeed, this is a well known fact in case (1.16) holds (e.g., see Lemma 4.6.1 in [17]), while it is a consequence of Proposition 1.1 in [7] in case u satisfies (1.17). \square

Proof of Lemma 1.1. Let us start by observing that

$$\ln|x - y| - \ln(1 + |y|) \leq \ln(|x| + 1), \quad \forall x, y \in \mathbb{R}^2.$$

Consequently, for $f \geq 0$, property (i) can be easily derived. To obtain (ii), let $|x| > 1$ and consider,

$$\sigma(x) = \frac{1}{\ln|x|} \int_{\mathbb{R}^2} (\ln|x - y| - \ln(1 + |y|) - \ln|x|) f(y) dy. \tag{1.18}$$

Thus, we must show that $\sigma(x) \rightarrow 0$ as $|x| \rightarrow +\infty$. For this purpose, write

$$\sigma(x) = \sigma_1(x) + \sigma_2(x) + \sigma_3(x),$$

where σ_1, σ_2 and σ_3 are defined by taking the integral in (1.18) over the regions $D_1 = \{y \in \mathbb{R}^2: |x - y| < 1\}$, $D_2 = \{y \in \mathbb{R}^2: |x - y| > 1 \text{ and } |y| < \rho\}$ and $D_3 = \{y \in \mathbb{R}^2: |x - y| > 1 \text{ and } |y| > \rho\}$ respectively, with $\rho > 1$ a fixed constant. We estimate,

$$\begin{aligned} |\sigma_1(x)| &\leq \frac{1}{\ln|x|} \int_{\{|x-y|<1\}} |\ln|x-y| - \ln(1+|y|) - \ln|x|| |f(y)| \, dy \\ &\leq \frac{1}{\ln|x|} \int_{\{|x-y|<1\}} \ln\left(\frac{1}{|x-y|}\right) |f(y)| \, dy + \frac{\ln(|x+2|) + \ln|x|}{\ln|x|} \int_{\{|x-y|<1\}} |f(y)| \, dy \\ &\leq \frac{1}{\ln|x|} \left(\int_{\{|x-y|<1\}} \frac{dy}{|x-y|} + \int_{\{|x-y|<1\}} |f(y)| (\ln|f(y)|)^+ \, dy \right) \\ &\quad + 2\left(1 + \frac{1}{\ln|x|}\right) \int_{\{|y|>|x-1\}} |f(y)| \, dy \\ &\leq \frac{1}{\ln|x|} (\pi + \|f(\ln|f|)^+\|_{L^1(\mathbb{R}^2)}) + 2\left(1 + \frac{1}{\ln|x|}\right) \int_{\{|y|>|x-1\}} |f(y)| \, dy, \end{aligned}$$

where we have used the well known inequality:

$$ab \leq e^a + b(\ln b - 1) \leq e^a + b(\ln b)^+, \quad \forall a \in \mathbb{R}, b > 0$$

with $a = \ln \frac{1}{|x-y|}$ and $b = |f(y)|$. Consequently, $\sigma_1(x) \rightarrow 0$ as $|x| \rightarrow +\infty$. Furthermore, for $x \in \mathbb{R}^2, |x| > 2\rho$, we have

$$\begin{aligned} |\sigma_2(x)| &\leq \frac{1}{\ln|x|} \int_{\{|x-y|>1, |y|<\rho\}} \left(\left| \ln\left(\frac{|x-y|}{|x|}\right) \right| + \ln(\rho+1) \right) |f(y)| \, dy \\ &\leq \frac{1}{\ln|x|} \ln(2(\rho+1)) \|f\|_{L^1(\mathbb{R}^2)} \rightarrow 0, \quad \text{as } |x| \rightarrow +\infty. \end{aligned}$$

Finally,

$$\begin{aligned} |\sigma_3(x)| &\leq \frac{1}{\ln|x|} \int_{\{|x-y|>1, \rho<|y|<2|x|\}} (\ln|x-y| + \ln(1+|y|) + \ln|x|) |f(y)| \, dy \\ &\quad + \frac{1}{\ln|x|} \int_{\{|x-y|>1, |y|>2|x|\}} \left| \ln\left(\frac{|x-y|}{1+|y|}\right) \right| |f(y)| + \int_{\{|y|>\rho\}} |f(y)| \, dy \\ &\leq 4 \int_{\{|y|>\rho\}} |f(y)| \, dy + \frac{3 \ln 3}{\ln|x|} \|f\|_{L^1(\mathbb{R}^2)} \rightarrow 4 \int_{\{|y|>\rho\}} |f(y)| \, dy \end{aligned}$$

as $|x| \rightarrow +\infty$, for any fixed $\rho > 1$. Thus, letting $\rho \rightarrow +\infty$, we obtain the desired conclusion. Now, suppose that (1.13) holds, then to establish (iii) it remains to show that

$$\left| \int_{\mathbb{R}^2} (\ln|x-y| - \ln|x|) f(y) \, dy \right| = O(1) \tag{1.19}$$

as $|x| \rightarrow +\infty$. As above, we are going to estimate the integral (1.19) over various regions. Firstly, note that for $|x| > 1$ we have:

$$\begin{aligned} \left| \int_{\{|x-y|<1\}} (\ln|x-y| - \ln|x|)f(y) \, dy \right| &\leq \int_{\{|x-y|<1\}} \ln\left(\frac{1}{|x-y|}\right)|f(y)| \, dy + \int_{\{|x-y|<1\}} \ln|x||f(y)| \, dy \\ &\leq \pi + \int_{\mathbb{R}^2} |f(y)|(\ln|f(y)|)^+ \, dy + \int_{\mathbb{R}^2} \ln(1+|y|)|f(y)| \, dy := C_1. \end{aligned}$$

On the other hand,

$$\begin{aligned} \left| \int_{\{|x-y|>1\}} (\ln|x-y| - \ln|x|)f(y) \, dy \right| &\leq \int_{\{|x-y|>1, |y|<|x|/2\}} \left| \ln\left(\frac{|x-y|}{|x|}\right) \right| |f(y)| \, dy + \int_{\{|x-y|>1, |y|>|x|/2\}} (\ln|x-y| + \ln|x|)|f(y)| \, dy \\ &\leq C_2 \left(\|f\|_{L^1} + \int_{\mathbb{R}^2} \ln(1+|y|)|f(y)| \, dy \right) \end{aligned}$$

for suitable $C_2 > 0$, and (1.19) follows. \square

Proof of Theorem 1.1. We start with the following:

Claim 1.

$$\sup_{\mathbb{R}^2} u_2 \leq \frac{c_0}{2} + \ln\left(\frac{1}{\pi} \|e^{u_2}\|_{L^1}\right). \tag{1.20}$$

Note that (1.20) immediately implies (1.5). To obtain (1.20), let us use the second equation in (P) together with Green’s representation formula, to derive that,

$$\begin{aligned} u_2(x) &\leq \frac{4c_0}{2\pi} \int_{\{|x-y|<r\}} \ln\left(\frac{r}{|x-y|}\right) \, dy + \frac{1}{2\pi r} \int_{\{|x-y|=r\}} u_2(y) \, d\sigma \\ &= c_0 r^2 + \frac{1}{2\pi r} \int_{\{|x-y|=r\}} u_2(y) \, d\sigma, \end{aligned} \tag{1.21}$$

$\forall r > 0$. Multiply both sides of (1.21) by $2r$ and integrate over $[0, 1]$ to obtain:

$$u_2(x) \leq \frac{c_0}{2} + \frac{1}{\pi} \int_{B_1(x)} u_2(y) \, dy.$$

At this point, we can use Jensen’s inequality to estimate,

$$\frac{1}{\pi} \int_{B_1(x)} u_2(y) \, dy \leq \ln\left(\frac{1}{\pi} \int_{B_1(x)} e^{u_2(y)} \, dy\right) \leq \ln\left(\frac{1}{\pi} \|e^{u_2}\|_{L^1}\right),$$

and conclude (1.20).

Claim 2.

$$u_1^+ \in L^\infty(\mathbb{R}^2). \tag{1.22}$$

To establish (1.22) note that $u_1(x) \rightarrow -\infty$ as $x \rightarrow z_k$, $k = 1, \dots, m$. So, we need to show that u_1 is bounded above outside a large ball which contains all points z_k 's, $k = 1, \dots, m$. To this end note that, if $x: |x| > \max_{k=1, \dots, m} |z_k| + 2$, then

$$\int_{B_1(x)} u_1^+ \leq \int_{\mathbb{R}^2} e^{u_1} \tag{1.23}$$

and

$$-\Delta u_1 = 4g^2 e^{u_1} + g^2 e^{u_2} \quad \text{in } B_1(x)$$

with $e^{u_2} \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$. So the argument given by Brezis and Merle [6] in the proof of Theorem 2 applies word by word and yields to the estimate:

$$\max_{B_{1/4}(x)} u_1 \leq C$$

with a suitable constant $C > 0$ independent of x .

To proceed further, let

$$w_2(x) = \frac{g^2}{2\pi} \int_{\mathbb{R}^2} (\ln|x-y| - \ln(1+|y|)) e^{u_2(y)} dy. \tag{1.24}$$

Hence, by Lemma 1.1, we have

$$w_2(x) \leq \sigma_2 \ln(|x| + 1), \quad \forall x \in \mathbb{R}^2, \tag{1.25}$$

and

$$\frac{w_2(x)}{\ln|x|} \rightarrow \sigma_2, \quad \text{as } |x| \rightarrow +\infty. \tag{1.26}$$

Define,

$$u_0(x) = \sum_{k=1}^m \ln|x - z_k|^{2n_k}, \tag{1.27}$$

and note that

$$u_1 - u_0 \in L^\infty_{\text{loc}}(\mathbb{R}^2). \tag{1.28}$$

Decompose:

$$u_1(x) = u_0(x) - w_2(x) + v_1(x) \tag{1.29}$$

so that $v_1(x)$ satisfies:

$$v_1 \in L^\infty_{\text{loc}}(\mathbb{R}^2); \quad -\Delta v_1(x) = 4g^2 e^{u_1(x)} = 4g^2 \prod_{k=1}^m |x - z_k|^{2n_k} e^{-w_2(x)} e^{v_1}, \tag{1.30}$$

and

$$\frac{v_1^+(x)}{\ln(|x| + 1)} \in L^\infty(\mathbb{R}^2).$$

Therefore, we may use Corollary 1.1 to derive that

$$-v_1(x) \leq \sigma_1 (\ln(|x| + 1)) + C \tag{1.31}$$

for suitable $C > 0$, and

$$\frac{v_1(x)}{\ln|x|} \rightarrow -\sigma_1, \quad \text{as } |x| \rightarrow +\infty. \tag{1.32}$$

In virtue of (1.25), (1.26) and (1.27), we conclude:

$$u_1(x) \geq (2N - (\sigma_1 + \sigma_2)) \ln|x| - C, \quad \text{as } |x| \rightarrow +\infty$$

and

$$\frac{u_1(x)}{\ln|x|} \rightarrow 2N - (\sigma_1 + \sigma_2) \quad \text{as } |x| \rightarrow +\infty.$$

So, (1.2) and (1.3) are established. Consequently, we must have that $\sigma_1 + \sigma_2 > 2(N + 1)$ and

$$\int_{\mathbb{R}^2} \ln(1 + |y|) e^{u_1(y)} dy < +\infty. \tag{1.33}$$

Thus, we can use part (iii) of Lemma 1.1 together with Corollary 1.1 to conclude that,

$$v_1(x) = \sigma_1 \ln \frac{1}{|x|} + O(1), \quad \text{as } |x| \rightarrow +\infty. \tag{1.34}$$

Finally, suppose that (1.6) holds, then we are in position to apply Corollary 1.1 to the function $v_2(x) = u_2(x) + c_0|x - x_0|^2$ with $f(x) = \frac{g^2}{2\cos^2\theta} e^{u_2} + 2g^2 e^{u_1}$, and conclude that

$$\frac{u_2(x) + c_0|x - x_0|^2}{\ln|x|} \rightarrow \frac{1}{2} \left(\sigma_1 + \frac{\sigma_2}{\cos^2\theta} \right) \quad \text{as } |x| \rightarrow +\infty.$$

In particular, this implies that u_2 admits exponential decay at infinity. Thus, $\int_{\mathbb{R}^2} \ln(1 + |y|) e^{u_2(y)} dy < +\infty$, and taking into account (1.33), we derive:

$$w_2(x) = \sigma_2 \ln|x| + O(1), \quad \text{as } |x| \rightarrow +\infty, \tag{1.35}$$

and

$$u_2(x) = -c_0|x - x_0|^2 + \frac{1}{2} \left(\sigma_1 + \frac{\sigma_2}{\cos^2\theta} \right) \ln|x| + O(1)$$

as $|x| \rightarrow +\infty$. Moreover, from (1.29), (1.34) and (1.35) we conclude

$$u_1(x) = \sum_{k=1}^m \ln|z - z_k|^{2n_k} - (\sigma_1 + \sigma_2) \ln|x| + O(1) \quad \text{as } |x| \rightarrow \infty,$$

and Theorem 1.1 is established. \square

Remark 1.1. We point out that, as in Lemma 1.3 of [8], it is possible to sharpen the asymptotic behavior (1.7) and (1.8) and prove that

$$\begin{aligned} \frac{\partial u_1}{\partial r} &\rightarrow 2N - (\sigma_1 + \sigma_2), & \frac{\partial u_1}{\partial \theta} &\rightarrow 0, \\ r \frac{\partial}{\partial r} (u_2 + c_0|x - x_0|^2) &\rightarrow \frac{1}{2} \left(\sigma_1 + \frac{\sigma_2}{\cos^2\theta} \right), & \frac{\partial}{\partial \theta} (u_2 + c_0|x - x_0|^2) &\rightarrow 0 \end{aligned}$$

uniformly, as $r = |x| \rightarrow +\infty$, with (r, θ) polar coordinates in \mathbb{R}^2 .

In the framework of Theorem 1.1, Spruck and Yang in [14] have constructed two families of solutions satisfying the asymptotic behavior (1.7), (1.8) (with $x_0 = 0$) and corresponding values σ_1, σ_2 free to vary as follows:

1. for the first family: $\sigma_1 \in (0, 4)$ and $\sigma_1 + \sigma_2 \in (2(N + 2), +\infty)$;
2. for the second family: $\sigma_1 \in (0, 2)$ and $2\sigma_1 + \sigma_2 \in (2(N + 1), +\infty)$.

In particular, Spruck–Yang’s construction allows one to obtain a solution pair for (P) with $\sigma_1 + \sigma_2$ equals to any prescribed value in the interval $(2(N + 1), +\infty)$. So, (1.4) is sharp. On the other hand, their construction always gives, $\sigma_2 > 2N$, while σ_1 can take values as small as wanted. For instance, according to Spruck–Yang’s result, it is possible to exhibit a family of solutions for (P) such that $\sigma_1 = o(1)$, while σ_2 can take any fixed value in $(2(N + 1), +\infty)$. Clearly, those solutions deny any sort of “quantization” property to the corresponding flux (0.5).

The aim of this paper is to show that the situation is quite different in case we consider solutions of (P) that admit the asymptotic behavior (1.7), (1.8) with σ_2 small. We see that in this case σ_1 may not be free to take any value in the interval $(2(N + 1), +\infty)$, as (1.4) would imply. In fact, in case all vortex points coincide, we see that σ_1 must remain close to the value $4(N + 1)$ and no other value is allowed. Thus solution of (P) with σ_2 small are very interesting, as they seem to restore a sort of “quantization” property for the flux in (0.5). We are going to construct such class of solutions. To this purpose, we consider the following function spaces introduced by Chae and Imanuvilov in [7].

For given $\alpha > 0$, let

$$X_\alpha = \left\{ u \in L^2_{\text{loc}}(\mathbb{R}^2) : \int_{\mathbb{R}^2} (1 + |x|^{2+\alpha}) |u(x)|^2 dx < \infty \right\},$$

which defines a Hilbert space equipped with the scalar product

$$(u, v) = \int_{\mathbb{R}^2} (1 + |x|^{2+\alpha}) uv.$$

Denote with $\|\cdot\|_{X_\alpha}$ the corresponding norm on X_α . Also let

$$Y_\alpha = \left\{ u \in W^{2,2}_{\text{loc}}(\mathbb{R}^2) : \Delta u \in X_\alpha, \frac{u}{(1 + |x|)^{1+\alpha/2}} \in L^2(\mathbb{R}^2) \right\}.$$

It defines a Hilbert space with corresponding natural scalar product and norm:

$$\|u\|_{Y_\alpha}^2 = \|\Delta u\|_{X_\alpha}^2 + \left\| \frac{u(x)}{1 + |x|^{1+\alpha/2}} \right\|_{L^2(\mathbb{R}^2)}^2.$$

We refer to [7] for the relevant properties of those spaces. In particular, from [7] we recall,

Proposition 1.1. (a) $X_\alpha \hookrightarrow L^1(\mathbb{R}^2)$ continuously, $Y_\alpha \hookrightarrow C^0_{\text{loc}}(\mathbb{R}^2)$.

(b) If $\alpha \in (0, 1)$ and $u \in Y_\alpha$ is harmonic, then u is constant.

(c) There exists a constant $C_1 > 0$ such that for all $u \in Y_\alpha$

$$|u(x)| \leq C_1 \|u\|_{Y_\alpha} ((\ln |x|)^+ + 1), \quad \forall x \in \mathbb{R}^2.$$

Notice that assumption (1.6) is motivated by considering solutions of (P) in the space $Y_\alpha \times Y_\alpha$. In fact, as an easy consequence of Theorem 1.1, it follows:

Corollary 1.2. *Under the assumptions of Theorem 1.1 we have*

- (i) $U_1 := u_1 - u_0 \in Y_\alpha, U_2 := u_2 + c_0|x - x_0|^2 \in Y_\alpha$ and

$$e^{u_1} \in X_\alpha \quad \text{for every } \alpha \in (0, 2(\sigma_1 + \sigma_2 - 2(N + 1))) \tag{1.36}$$
- (ii) $\frac{1}{2}U_1 + U_2 \in Y_\alpha$ and $e^{u_2} \in X_\alpha, \forall \alpha > 0$.

To complement the situation analyzed by Spruck and Yang in [14], we consider solutions of (P) with $\sigma_2 \leq 2N$. In this case, the following interesting estimate holds, when all vortex points coincide.

Lemma 1.2. *Let (u_1, u_2) be solutions of (P) with $z_1 = z_2 = \dots = z_m$ and such that $e^{u_2} \in X_\alpha$ with $\alpha > 2$ and $\sigma_2 \leq 2N$. For any $q \in (2, \frac{\alpha+4}{3})$ there exists a constant $C_q = C(q, c_0, \alpha) > 0$ such that*

$$|\sigma_1 + 2\sigma_2 - 4(N + 1)| \leq C_q \sigma_2^{(q-2)/q} \|e^{u_2}\|_{X_\alpha}^{2/q}. \tag{1.37}$$

Remark 1.2. Clearly, estimate (1.37) bears interesting consequences in case $\|e^{u_2}\|_{X_\alpha}$ (and hence σ_2) is small, as it forces σ_1 to remain close to the value $4(N + 1)$. Such a situation can actually occur, as we are going to construct family of solutions $(u_{1,\varepsilon}, u_{2,\varepsilon})$ for (P) (when all vortex points coincide) such that, as $\varepsilon \rightarrow 0, \|e^{u_{2,\varepsilon}}\|_{X_\alpha} = o(1)$, so

$$\sigma_{2,\varepsilon} = \frac{g^2}{2\pi} \int_{\mathbb{R}^2} e^{u_{2,\varepsilon}} = o(1), \tag{1.38}$$

$$\sigma_{1,\varepsilon} = \frac{2g^2}{\pi} \int_{\mathbb{R}^2} e^{u_{1,\varepsilon}} = 4(N + 1) + o(1). \tag{1.39}$$

Proof. Without loss of generality suppose that $z_1 = \dots = z_m = 0$. Thus, by (1.30) we have:

$$v_1 \in L^\infty_{loc}(\mathbb{R}^2), \quad -\Delta v_1 = 4g^2|x|^{2N}e^{-w_2(x)}e^{v_1}$$

with w_2 as defined in (1.24). Since $e^{u_2} \in X_\alpha$, then we also have $\int_{\mathbb{R}^2} \ln(1 + |x|)e^{u_2(x)} dx < +\infty$, and in view of Lemma 1.2, we derive

$$w_2(x) = \sigma_2 \ln(|x| + 1) + O(1).$$

Consequently, setting $R(x) = 4g^2|x|^{2N}e^{-w_2(x)}$, we see that

$$R(x) = O(|x|^{2N-\sigma_2}), \tag{1.40}$$

and, by assumption, $2N - \sigma_2 \geq 0$. Furthermore,

$$\int_{\mathbb{R}^2} R(x)e^{v_1(x)} dx = 4g^2 \int_{\mathbb{R}^2} e^{u_1} = 2\pi\sigma_1 \tag{1.41}$$

and, from (1.40), we get $\int_{\mathbb{R}^2} (1 + |x|^{2N-\sigma_2})e^{v_1} < +\infty$. Thus, we are in position to apply Theorem 1 (when $2N = \sigma_2$), or Theorem 2 (when $2N > \sigma_2$) of Chen and Li [8], to conclude:

$$\begin{aligned} \pi\sigma_1(\sigma_1 - 4) &= \int_{\mathbb{R}^2} x \cdot \nabla R(x) e^{v_1(x)} dx \\ &= 4g^2 \left(2N \int_{\mathbb{R}^2} R(x) e^{v_1} - \int_{\mathbb{R}^2} x \cdot \nabla w_2(x) R(x) e^{v_1} \right) \\ &= 4\pi N\sigma_1 - 2\pi\sigma_2\sigma_1 - 4g^2 \int_{\mathbb{R}^2} (x \cdot \nabla w_2(x) - \sigma_2) R(x) e^{v_1}. \end{aligned}$$

That is,

$$|\sigma_1 + 2\sigma_2 - 4(N + 1)| \leq \frac{4g^2}{\pi\sigma_1} \int_{\mathbb{R}^2} |x \cdot \nabla w_2(x) - \sigma_2| R(x) e^{v_1(x)} dx. \tag{1.42}$$

On the other hand, by (1.24) we have:

$$\begin{aligned} |x \cdot \nabla w_2(x) - \sigma_2| &\leq \frac{g^2}{2\pi} \int_{\mathbb{R}^2} \frac{|y|}{|x - y|} e^{u_2(y)} dy \leq \frac{g^2}{2\pi} \int_{\mathbb{R}^2} \frac{(1 + |y|)^{\frac{2+\alpha}{q}} e^{u_2(y)}}{|x - y|(1 + |y|)^{\frac{2+\alpha}{q}-1}} dy \\ &\leq \frac{g^2}{2\pi} \left(\int_{\mathbb{R}^2} (1 + |y|)^{2+\alpha} e^{qu_2} dy \right)^{\frac{1}{q}} \left(\int_{\mathbb{R}^2} \frac{dy}{|x - y|^{\frac{q}{q-1}} (1 + |y|)^{\frac{2+\alpha-q}{q-1}}} \right)^{\frac{q-1}{q}} \\ &\leq \frac{g^2}{2\pi} \|e^{u_2}\|_{L^\infty(\mathbb{R}^2)}^{\frac{q-2}{q}} \|e^{u_2}\|_{X_\alpha}^{2/q} \left(\int_{\{|x-y|<1\}} \frac{dy}{|x - y|^{\frac{q}{q-1}}} + \int_{\{|x-y|\geq 1\}} \frac{dy}{(1 + |y|)^{\frac{2+\alpha-1}{q-1}}} \right)^{\frac{q-1}{q}}. \end{aligned}$$

At this point, the desired estimate (1.37) follows, by means of (1.5), and the observation that, by the choice of $q \in (2, \frac{\alpha+4}{3})$ we have $1 < \frac{q}{q-1} < 2 < \frac{2+\alpha-q}{q-1}$. Thus,

$$|x \cdot \nabla w_2(x) - \sigma_2| \leq C_{1,q} \|e^{u_1}\|_{L^1(\mathbb{R}^2)}^{\frac{q-2}{q}} \|e^{u_2}\|_{X_\alpha}^{\frac{2}{q}} \tag{1.43}$$

with a suitable constant $C_{1,q} > 0$ independent of $x \in \mathbb{R}^2$. Hence, we can use (1.43) in (1.42), and by (1.41), obtain (1.37). \square

2. Existence result

We devote this section to the construction of a four-parameter family of solutions for (P) in case all vortex points coincide, such that (1.38) and (1.39) hold. Without loss of generality we take all vortex points to coincide with the origin and obtain,

Theorem 2.1. *Let $z_1 = z_2 = \dots = z_m = 0$, there exists $\varepsilon_0 > 0$ sufficiently small such that problem (P) admits a five-parameters family of solutions $(u_{1,\alpha}^\varepsilon, u_{2,\alpha}^\varepsilon)$ with $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$, $\alpha \in \mathbb{R}^4$ and $|\alpha| < \varepsilon_0$ satisfying:*

$$(i) \quad \frac{1}{\varepsilon} \|e^{u_{2,\alpha}^\varepsilon}\|_{X_\alpha} = O(1) \quad \forall \alpha > 0, \quad \frac{2g^2}{\pi} \int_{\mathbb{R}^2} e^{u_{1,\alpha}^\varepsilon} = 4(N + 1) + \varepsilon O(1) \quad \text{as } \varepsilon \rightarrow 0,$$

(ii)

$$u_{1,\alpha}^\varepsilon(x) = -\left[2(N+2) + \varepsilon\left(\frac{1}{2c_0}\left(\frac{(N+1)!}{c_0^{N+1}} - 1\right) + \beta_{1,\alpha}^\varepsilon\right)\right] \ln|x| + O(1),$$

$$u_{2,\alpha}^\varepsilon(x) = -c_0|x|^2 + (2(N+1) + \beta_{2,\alpha}^\varepsilon) \ln|x| + \ln\varepsilon + O(1) \quad \text{as } |x| \rightarrow \infty,$$

where $\beta_{1,\alpha}^\varepsilon \rightarrow 0, \beta_{2,\alpha}^\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0, |\alpha| \rightarrow 0$, and $O(1)$ denotes a quantity bounded uniformly in (ε, α) .

In order to obtain Theorem 2.1, we shall establish a more general existence result concerning the problem:

$$(P_\varepsilon) \quad \begin{cases} -\Delta u_1 = 4g^2 e^{u_1} + g^2 e^{u_2} - 4\pi \sum_{k=1}^m n_k \delta(z - \varepsilon z_k), \\ \Delta u_2 = \frac{g^2}{2 \cos^2 \theta} (e^{u_2} - \phi_0^2) + 2g^2 e^{u_1}, \\ \int_{\mathbb{R}^2} e^{u_1} < \infty, \quad \int_{\mathbb{R}^2} e^{u_2} < \infty \end{cases}$$

with z_1, \dots, z_m arbitrarily given points in \mathbb{R}^2 and small $\varepsilon \in \mathbb{R}$. For this purpose, it is convenient to introduce complex notation and identify the pair $x = (x_1, x_2) \in \mathbb{R}^2$ with the complex number $z = x + iy \in \mathbb{C}$. Let

$$f_\varepsilon(z) = (N+1) \prod_{k=1}^m (z - \varepsilon z_k)^{n_k}, \quad \text{and} \quad F_\varepsilon(z) = \int_0^z f_\varepsilon(\xi) d\xi. \tag{2.1}$$

For any $a \in \mathbb{C}$, define,

$$\eta_{\varepsilon,a}(z) = \frac{8|f_\varepsilon(z)|^2}{(1 + |F_\varepsilon(z) + a|^2)^2}, \tag{2.2}$$

and consider the radial function $\rho = \rho(r)$ so that,

$$\rho(|z|) := \eta_{\varepsilon=0,a=0}(z) = \frac{8(N+1)^2 |z|^{2N}}{(1 + |z|^{2N+2})^2}. \tag{2.3}$$

As well known (e.g., [12]), $\forall \varepsilon \geq 0$ and $\forall a \in \mathbb{C}$ we have

$$-\Delta \ln \eta_{\varepsilon,a} = \eta_{\varepsilon,a} - 4\pi \sum_{k=1}^m n_k \delta(z - \varepsilon z_k), \quad \text{in } \mathbb{R}^2,$$

$$\int_{\mathbb{R}^2} \eta_{\varepsilon,a} = 8\pi(N+1). \tag{2.4}$$

We shall look for solutions of (P_ε) with a specific structure. Namely, we set

$$u_1(z) = \ln \eta_{\varepsilon,a}(z) + \varepsilon(w_1(z) + v_1(z)) - 2 \ln 2g,$$

$$u_2(z) = -c_0|z|^2 + w_2(z) + \ln \varepsilon - 2 \ln g + v_2(z), \tag{2.5}$$

where w_2 is the radial function given by

$$w_2(r) = \ln(1 + r^{2N+2}), \tag{2.6}$$

and so, it satisfies

$$\Delta w_2 = \frac{\rho}{2},$$

while w_1 is the radial solution in Y_α , $\alpha \in (0, \frac{1}{2})$ of the equation:

$$\Delta w + \rho w + e^{-c_0|z|^2}(1 + |z|^{2N+2}) = 0 \quad \text{in } \mathbb{R}^2, \tag{2.7}$$

as constructed in Lemma 2.1 of [7]. Consequently, relative to the unknowns (v_1, v_2) , problem (P_ε) becomes:

$$(P'_\varepsilon) \quad \begin{cases} -\Delta v_1 = e^{-c_0|z|^2+w_2+v_2} + \eta_{\varepsilon,a} \frac{(e^{\varepsilon(v_1+w_1)} - 1)}{\varepsilon} - \rho w_1 - e^{-c_0|z|^2+w_2}, \\ \Delta v_2 = \frac{\varepsilon}{2 \cos^2 \theta} e^{-c_0|z|^2+w_2+v_2} + \frac{1}{2} \eta_{\varepsilon,a} e^{\varepsilon(v_1+w_1)} - \frac{\rho}{2}, \end{cases}$$

as it follows by direct computations. Notice that, by continuity problem (P'_ε) may be considered also at $\varepsilon = 0$, and for $a = 0$, $(P'_{\varepsilon=0})$ admits the (trivial) solution $v_1 = v_2 = 0$. For ε small we aim to construct solutions for (P'_ε) which “bifurcate” from such trivial solution. To this purpose, we start by collecting some useful properties of the function w_1 , which will be established in Appendix A.

Lemma 2.1. *Problem (2.7) admits a radial solution w_1 such that*

- (i) $w_1 \in Y_\alpha, \quad \forall \alpha \in (0, \frac{1}{2});$
- (ii)

$$w_1(r) = \frac{1}{2c_0} \left(1 - \frac{(N+1)!}{c_0^{N+1}} \right) \ln r + O(1), \quad \text{as } r \rightarrow +\infty;$$

$$\frac{r dw_1}{dr} \rightarrow \frac{1}{2c_0} \left(1 - \frac{(N+1)!}{c_0^{N+1}} \right), \quad \text{as } r \rightarrow +\infty;$$
- (iii)
$$\int_0^{+\infty} \left(-2\rho(r)w_1(r) \frac{r^{2(N+1)}}{(1+r^{2(N+1)})^2} + \frac{e^{-c_0 r^2} r^{2(N+1)}}{1+r^{2(N+1)}} \right) r \, dr = \frac{1}{2c_0}.$$

From now on, the function w_1 in (2.5) is chosen according to Lemma 2.1.

For fixed $\alpha > 0$ define the operator

$$P : Y_\alpha^2 \times \mathbb{C} \times \mathbb{R} \rightarrow X_\alpha^2$$

by setting $P = (P_1, P_2)$ with

$$P_1(v_1, v_2, a, \varepsilon) = \Delta v_1 + e^{-c_0|z|^2+w_2+v_2} + \eta_{\varepsilon,a} \frac{(e^{\varepsilon(v_1+w_1)} - 1)}{\varepsilon} - \rho w_1 - e^{-c_0|z|^2+w_2},$$

$$P_2(v_1, v_2, a, \varepsilon) = \Delta v_2 - \frac{\varepsilon}{2 \cos^2 \theta} e^{-c_0|z|^2+w_2+v_2} - \frac{1}{2} \eta_{\varepsilon,a} e^{\varepsilon(v_1+w_1)} - \frac{\rho}{2},$$

and extended by continuity at $\varepsilon = 0$. Thus, $P(0, 0, 0, 0) = (0, 0)$, and finding a solution for problem (P'_ε) is now reduced to finding a small $\varepsilon_0 > 0$ and an implicit function

$$\varepsilon \mapsto (v_{1,\varepsilon}, v_{2,\varepsilon}, a_\varepsilon) : (-\varepsilon_0, \varepsilon_0) \rightarrow Y_\alpha^2 \times \mathbb{C}$$

satisfying $P(v_{1,\varepsilon}, v_{2,\varepsilon}, a_\varepsilon, \varepsilon) = 0, \forall \varepsilon \in (-\varepsilon_0, \varepsilon_0)$. To this end, let $a = a_1 + ia_2$ and observe that,

$$\left. \frac{\partial \eta_{\varepsilon,a}(z)}{\partial a_1} \right|_{a=0, \varepsilon=0} = -4\rho\varphi_+, \quad \left. \frac{\partial \eta_{\varepsilon,a}(z)}{\partial a_2} \right|_{a=0, \varepsilon=0} = -4\rho\varphi_-, \tag{2.8}$$

where, in polar coordinates,

$$\varphi_+(r, \theta) = \frac{r^{N+1} \cos(N+1)\theta}{1+r^{2N+2}}, \quad \varphi_-(r, \theta) = \frac{r^{N+1} \sin(N+1)\theta}{1+r^{2N+2}}.$$

Therefore, the linearized operator $P'_{(v_1, v_2, a)} = (P'_{1, (v_1, v_2, a)}, P'_{2, (v_1, v_2, a)})$ of P at $(0, 0, 0, 0)$ may be easily computed as given by,

$$\begin{aligned} P'_{1, (v_1, v_2, a)}(0, 0, 0, 0)[\psi_1, \psi_2, b] &= \Delta\psi_1 + \rho\psi_1 + e^{-c_0|z|^2+w_2}\psi_2 - 4\rho w_1\varphi_+b_1 - 4\rho w_1\varphi_-b_2, \\ P'_{2, (v_1, v_2, a)}(0, 0, 0, 0)[\psi_1, \psi_2, b] &= \Delta\psi_2 + 2\rho\varphi_+b_1 + 2\rho\varphi_-b_2. \end{aligned}$$

Set $P'_{(v_1, v_2, a)}(0, 0, 0, 0) = A$, we have:

Proposition 2.1. *Let $\alpha \in (0, \frac{1}{2})$, then the operator $A : Y_\alpha^2 \times \mathbb{C} \rightarrow X_\alpha^2$ is onto, and*

$$\text{Ker } A = \text{Span}\{(\varphi_+, 0), (\varphi_-, 0), (\varphi_0, 0), (\omega_1, 1)\} \times \{\mathbf{0}\},$$

where φ_0 is the radial function:

$$\varphi_0(r) = \frac{1-r^{2(N+1)}}{1+r^{2(N+1)}}, \quad r = |z|.$$

Proof. In order to prove Proposition 2.1, we recall the following result established in [7] for the operator

$$L = \Delta + \rho : Y_\alpha \rightarrow X_\alpha.$$

Namely, if $\alpha \in (0, \frac{1}{2})$, then

$$\text{Ker } L = \text{Span}\{\varphi_+, \varphi_-, \varphi_0\}, \tag{2.9}$$

(see Lemma 2.4 in [7]) and

$$\text{Im } L = \left\{ f \in X_\alpha : \int_{\mathbb{R}^2} f\varphi_\pm = 0 \right\} \tag{2.10}$$

(see Proposition 2.2 in [7]). Now, let $(f_1, f_2) \in X_\alpha^2$, we seek $(\psi_1, \psi_2, b) \in Y_\alpha^2 \times \mathbb{C}$, $b = b_1 + ib_2$ such that

$$\Delta\psi_1 + \rho\psi_1 + e^{-c_0|z|^2+w_2}\psi_2 - 4\rho w_1\varphi_+b_1 - 4\rho w_1\varphi_-b_2 = f_1, \tag{2.11}$$

and

$$\Delta\psi_2 + 2\rho\varphi_+b_1 + 2\rho\varphi_-b_2 = f_2. \tag{2.12}$$

We start by considering (2.12). Decompose

$$\psi_2 = \phi_2 + 2d_1\varphi_+ + 2d_2\varphi_-, \tag{2.13}$$

where the constants d_1, d_2 will be specified later, and $\phi_2 \in Y_\alpha$:

$$\int_{\mathbb{R}^2} \Delta\phi_2\varphi_\pm = 0. \tag{2.14}$$

Therefore, recalling that $\Delta\varphi_\pm + \rho\varphi_\pm = 0$, (2.12) holds if and only if

$$\Delta\phi_2 = 2(d_1 - b_1)\rho\varphi_+ + 2(d_2 - b_2)\rho\varphi_- + f_2. \tag{2.15}$$

In order to meet the orthogonality condition (2.14), necessarily:

$$\int_{\mathbb{R}^2} f_2 \varphi_+ + 2(d_1 - b_1) \int_{\mathbb{R}^2} \rho \varphi_+^2 = 0; \quad \int_{\mathbb{R}^2} f_2 \varphi_- + 2(d_2 - b_2) \int_{\mathbb{R}^2} \rho \varphi_-^2 = 0. \quad (2.16)$$

By elementary calculations, we see that

$$\int_{\mathbb{R}^2} \rho \varphi_{\pm}^2 = \frac{2}{3} \pi (N + 1).$$

So (2.16) requires the choice:

$$d_1 = b_1 - \frac{3}{4\pi(N+1)} \int_{\mathbb{R}^2} f_1 \varphi_+, \quad d_2 = b_2 - \frac{3}{4\pi(N+1)} \int_{\mathbb{R}^2} f_2 \varphi_-. \quad (2.17)$$

Hence, inserting (2.17) into (2.15), and letting

$$f = f_2 - \left(\frac{3}{2\pi(N+1)} \int_{\mathbb{R}^2} f_2 \varphi_+ \right) \rho \varphi_+ - \left(\frac{3}{2\pi(N+1)} \int_{\mathbb{R}^2} f_2 \varphi_- \right) \rho \varphi_- \in X_{\alpha},$$

it suffices to take

$$\phi_2(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \ln|x-y| f(y) dy, \quad (2.18)$$

in order to obtain a solution for (2.14)–(2.15) in Y_{α} .

Thus ψ_2 in (2.13) is determined by (2.17) and (2.18). Now, insert such ψ_2 into (2.11) to obtain:

$$\Delta \psi_1 + \rho \psi_1 = g_1 - 2(e^{-c_0|z|^2+w_2} - 2\rho w_1)(b_1 \varphi_+ + b_2 \varphi_-) \quad (2.19)$$

with

$$g_1 = f_1 - e^{-c_0|z|^2+w_2} \left[\phi_2 - \frac{3}{2\pi(N+1)} \left(\varphi_+ \int_{\mathbb{R}^2} f_2 \varphi_+ + \varphi_- \int_{\mathbb{R}^2} f_2 \varphi_- \right) \right],$$

and ϕ_2 given in (2.18). Next, we are going to choose b_1 and b_2 in order to insure that the right hand side of (2.19) satisfies to the orthogonality conditions required by (2.10). Namely, we impose:

$$\int_{\mathbb{R}^2} g_1 \varphi_+ - 2b_1 \int_{\mathbb{R}^2} (e^{-c_0|z|^2+w_2} - 2\rho w_1) \varphi_+^2 = 0, \quad (2.20)$$

$$\int_{\mathbb{R}^2} g_1 \varphi_- - 2b_2 \int_{\mathbb{R}^2} (e^{-c_0|z|^2+w_2} - 2\rho w_1) \varphi_-^2 = 0. \quad (2.21)$$

Recalling (2.6), we can use part (iii) of Lemma 2.1 to obtain:

$$\int_{\mathbb{R}^2} (e^{-c_0|z|^2+w_2} - 2\rho w_1) \varphi_{\pm}^2 dx = \pi \int_0^{+\infty} (e^{-c_0 r^2} (1 + r^{2(N+1)}) - 2\rho(r) w_1(r)) \frac{r^{2(N+1)}}{(1 + r^{2(N+1)})^2} r dr = \frac{\pi}{2c_0}.$$

Consequently, we may choose:

$$b_1 = \frac{c_0}{\pi} \int_{\mathbb{R}^2} g_1 \varphi_+, \quad b_2 = \frac{c_0}{\pi} \int_{\mathbb{R}^2} g_1 \varphi_-,$$

and obtain that the right-hand side of (2.19) belongs to the image of the operator $L = \Delta + \rho$ when defined over Y_α (see (2.10)). Thus we derive ψ_1 and the desired conclusion follows. Next, to determine $\text{Ker } A$, take $f_1 = f_2 = 0$ in the above computations. We find immediately, that $d_1 = b_1 = 0$, $d_2 = b_2 = 0$, and $\Delta\phi_2 = 0$. Hence, by Proposition 1.1(b), ϕ_2 must be a constant. If $\phi_2 = 0$, then $g_1 = 0$ and we must take $\psi_1 \in \text{ker } L$. If $\phi_2 \neq 0$, say $\phi_2 = 1$, then $g_1 = -e^{-c_0|z|^2 + \omega_2}$ and ψ_1 must satisfy: $\Delta\psi_1 + \rho\psi_1 + e^{-c_0|z|^2 + \omega_2} = 0$. Thus, $\psi_1 \in \omega_1 + \text{ker } L$ and we conclude that $\text{ker } A = W \times \{0\}$ with $W \subset Y_\alpha^2$ given by

$$W = \text{Span}\{\text{ker } L \times \{0\}, (\omega_1, 1)\} \tag{2.22}$$

as claimed. \square

Denote by $V_\alpha \subset Y_\alpha^2$ the space orthogonal to W defined in (2.22) with respect to the scalar product in Y_α^2 . Hence, we may write:

$$Y_\alpha^2 = V_\alpha \oplus W.$$

Furthermore, for given $r > 0$ denote by:

$$Q_r = \{(\varepsilon, \alpha) \in \mathbb{R}^5: \varepsilon \in (-r, r) \text{ and } \alpha = (s, \alpha_0, \alpha_1, \alpha_2) \in \mathbb{R}^4, |\alpha| < r\}.$$

By a direct application of the Implicit Function Theorem (see Theorem 2.7.5 in [11]) we may conclude:

Theorem 2.2. *Let $\alpha \in (0, 1/2)$ there exists $\varepsilon_0 > 0$ sufficiently small and continuous functions*

$$a(\varepsilon, \alpha): Q_{\varepsilon_0} \rightarrow \mathbb{C}, \quad v(\varepsilon, \alpha) = (v_1(\varepsilon, \alpha), v_2(\varepsilon, \alpha)): Q_{\varepsilon_0} \rightarrow V_\alpha,$$

all vanishing at $\varepsilon = 0$ and $\alpha = 0$, such that $\forall \varepsilon \in (-\varepsilon_0, \varepsilon_0)$ problem (P_ε) admits a four-parameter family of solutions $(u_{1,\alpha}^\varepsilon, u_{2,\alpha}^\varepsilon)$, $\alpha = (s, \alpha_0, \alpha_1, \alpha_2) \in \mathbb{R}^4$, $|\alpha| < \varepsilon_0$, which decompose as follows:

$$\begin{aligned} u_{1,\alpha}^\varepsilon &= \ln \eta_{\varepsilon,a(\varepsilon,\alpha)} + \varepsilon((1+s)w_1 + \alpha_0\varphi_0 + \alpha_1\varphi_+ + \alpha_2\varphi_- + v_1(\varepsilon, \alpha)) - 2 \ln 2g, \\ u_{2,\alpha}^\varepsilon &= -c_0|z|^2 + w_2(z) + \ln \varepsilon - 2 \ln g + s + v_2(\varepsilon, \alpha). \end{aligned}$$

At this point, if we take $z_1 = z_2 = \dots = z_m = 0$, then problem (P) and (P_ε) are one and the same, and in view of (2.2), (2.4), (2.6), Lemma 2.1(ii), and Proposition 1.1(c), we easily derive Theorem 2.1.

Final remark. It would be interesting to know whether or not a result similar to Theorem 2.1 remains valid even when the vortex points do not coincide.

Acknowledgements

The authors wish to thank M. Nolasco for useful comments.

This research is supported partially by the grant no.2000-2-10200-002-5 from the basic research program of the KOSEF Korea, and by M.U.R.S.T. project: Variational Methods and Non Linear Differential Equation, Italy.

Appendix A

In order to establish Lemma 2.1, let us consider the operator

$$L_1 = \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} + \rho \tag{A.1}$$

given by the restriction of the operator $L = \Delta + \rho$ over the radial functions. Thus, we wish to solve

$$L_1 w = f \tag{A.2}$$

with $f(r) = -e^{-c_0 r^2 + w_2(r)} = -e^{-c_0 r^2} (1 + r^{2(N+1)})$. Chae and Imanuvilov in [7] have obtained an integral representation for solutions of (A.2) in case

$$f \in C^1(\mathbb{R}^+) \cap X_\alpha \quad \text{and} \quad \alpha \in \left(0, \frac{1}{2}\right). \tag{A.3}$$

Lemma A.1 (see Lemma 2.1 in [7]). *Assume (A.3), then (A.2) admits a solution $w \in Y_\alpha$ given by the formula*

$$w(r) = \varphi_0(r) \left\{ \int_0^r \frac{\phi_f(s) - \phi_f(1)}{(1-s)^2} ds + \frac{\phi_f(1)r}{1-r} \right\} \tag{A.4}$$

with

$$\phi_f(r) = (\varphi_0(r))^{-2} \frac{(1-r)^2}{r} \int_0^r \varphi_0(t) t f(t) dt, \quad \varphi_0(r) = \frac{1 - r^{2(N+1)}}{1 + r^{2(N+1)}},$$

where $\phi_f(1)$ and $w(r)$ are extended by continuity at $r = 1$.

In order to obtain the solution w_1 as claimed in Lemma 2.1, we shall use such a representation formula with $f(r) = -e^{-c_0 r^2} (1 + r^{2(N+1)})$. To this purpose, for given $n \in \mathbb{N}$, let

$$I_n(r) = \int_0^r (1 - t^{2n}) t e^{-c_0 t^2} dt, \tag{A.5}$$

and notice that

$$\phi_f(r) = -(\varphi_0(r))^{-2} \frac{(1-r)^2}{r} I_{N+1}(r). \tag{A.6}$$

Lemma A.2. *The following identity holds:*

$$I_n(r) = \frac{1}{2c_0} \left[1 - \frac{n!}{c_0^n} - e^{-c_0 r^2} \left(1 - \sum_{k=0}^n \frac{r^{2(n-k)}}{c_0^k} \frac{n!}{(n-k)!} \right) \right]. \tag{A.7}$$

Proof. We shall proceed by induction. For $n = 1$,

$$\begin{aligned} I_{n=1}(r) &= \int_0^r (1 - t^2) t e^{-c_0 t^2} dt \\ &= -\frac{1}{2c_0} e^{-c_0 t^2} \Big|_{t=0}^{t=r} + \frac{1}{2c_0} \int_0^r \frac{d}{dt} e^{-c_0 t^2} t^2 dt \\ &= \frac{1}{2c_0} \left(1 - e^{-c_0 r^2} + r^2 e^{-c_0 r^2} - 2 \int_0^r t e^{-c_0 t^2} dt \right) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2c_0} \left[1 - e^{-c_0 r^2} + r^2 e^{-c_0 r^2} + \frac{1}{c_0} (e^{-c_0 r^2} - 1) \right] \\ &= \frac{1}{2c_0} \left[1 - \frac{1}{c_0} - e^{-c_0 r^2} \left(1 - r^2 - \frac{1}{c_0} \right) \right], \end{aligned}$$

and (A.7) is established for $n = 1$. Now assume that (A.7) holds for $n \geq 1$, and we proceed to prove it for $n + 1$. To this end, notice that

$$\begin{aligned} I_{n+1}(r) &= \int_0^r (1 - t^{2(n+1)}) t e^{-c_0 t^2} dt \\ &= \frac{1}{2c_0} \left(1 - e^{-c_0 r^2} + r^{2(n+1)} e^{-c_0 r^2} - 2(n+1) \int_0^r t^{2n} t e^{-c_0 t^2} dt \right) \\ &= \frac{1}{2c_0} \left(1 - e^{-c_0 r^2} + r^{2(n+1)} e^{-c_0 r^2} + 2(n+1) I_n(r) - 2(n+1) \int_0^r t e^{-c_0 t^2} dt \right) \\ &= \frac{1}{2c_0} \left[1 - e^{-c_0 r^2} + r^{2(n+1)} e^{-c_0 r^2} + 2(n+1) I_n(r) + \frac{n+1}{c_0} (e^{-c_0 r^2} - 1) \right]. \end{aligned}$$

Thus, if we apply the inductive assumption (A.7), substituting above, we find

$$\begin{aligned} I_{n+1}(r) &= \frac{1}{2c_0} \left[1 - e^{-c_0 r^2} + r^{2(n+1)} e^{-c_0 r^2} + \frac{2(n+1)}{2c_0} \left(1 - \frac{n!}{c_0^n} \right. \right. \\ &\quad \left. \left. - e^{-c_0 r^2} \left(1 - \sum_{k=0}^n \frac{r^{2(n-k)}}{c_0^k} \frac{n!}{(n-k)!} \right) \right) + \frac{n+1}{c_0} e^{-c_0 r^2} - \frac{n+1}{c_0} \right] \\ &= \frac{1}{2c_0} \left[1 - \frac{(n+1)!}{c_0^{n+1}} - e^{-c_0 r^2} \left(1 - r^{2(n+1)} - \sum_{k=0}^n \frac{r^{2(n-k)}}{c_0^{k+1}} \frac{(n+1)!}{(n-k)!} \right) \right] \\ &= \frac{1}{2c_0} \left[1 - \frac{(n+1)!}{c_0^{n+1}} - e^{-c_0 r^2} \left(1 - r^{2(n+1)} - \sum_{k=1}^{n+1} \frac{r^{2[(n+1)-k]}}{c_0^k} \frac{(n+1)!}{(n+1-k)!} \right) \right] \\ &= \frac{1}{2c_0} \left[1 - \frac{(n+1)!}{c_0^{n+1}} - e^{-c_0 r^2} \left(1 - \sum_{k=0}^{n+1} \frac{r^{2[(n+1)-k]}}{c_0^k} \frac{(n+1)!}{(n+1-k)!} \right) \right] \end{aligned}$$

and the desired identity is established. \square

An immediate consequence of Lemma 3.2 gives

$$I_{N+1}(r) \rightarrow \frac{1}{2c_0} \left(1 - \frac{(N+1)!}{c_0^{N+1}} \right) := \gamma_0 \quad \text{as } r \rightarrow +\infty. \tag{A.8}$$

Furthermore, for $r > 2$, inserting (A.7) into (A.4), we find

$$w_1(r) = -\varphi_0(r) \int_2^r \left(\frac{1 + t^{2(N+1)}}{1 - t^{2(N+1)}} \right)^2 \frac{I_{N+1}(t)}{t} dt + O(1)$$

$$\begin{aligned} &= -\varphi_0(r) \int_0^r \left(\frac{1+t^{2(N+1)}}{1-t^{2(N+1)}} \right)^2 \frac{\gamma_0}{t} dt + O(1) \\ &= -\gamma_0 \varphi_0(r) \ln r + O(1). \end{aligned}$$

Consequently,

$$w_1(r) = \gamma_0 \ln r + O(1), \quad \text{as } r \rightarrow +\infty. \tag{A.9}$$

On the other hand, for $r \rightarrow +\infty$,

$$\begin{aligned} \frac{r dw_1}{dr}(r) &= -r\varphi_0'(r)(\varphi_0(r))^{-1}w_1(r) - (\varphi_0(r))^{-1}I_{N+1}(r) + O\left(\frac{1}{r}\right) \\ &= -(\varphi_0(r))^{-1} \left[I_{N+1}(r) + \frac{4(N+1)r^{2(N+1)}}{(1+r^{2(N+1)})^2}w_1(r) + O\left(\frac{1}{r}\right) \right]. \end{aligned}$$

So, taking into account (A.9), we immediately conclude that

$$\frac{r dw_1}{dr}(r) \rightarrow \gamma_0 \quad \text{as } r \rightarrow +\infty \tag{A.10}$$

and, part (ii) of Lemma 2.1 follows by (3.9) and (3.10). In order to establish (iii), notice that, by direct computation, we find

$$\frac{1}{2}L_1\left(\frac{1}{(1+r^{2(N+1)})^2}\right) = \frac{8(N+1)^2r^{4N+2}}{(1+r^{2(N+1)})^4} = \frac{\rho(r)r^{2(N+1)}}{(1+r^{2(N+1)})^2},$$

while, by definition, $L_1w_1 = -e^{-c_0r^2}(1+r^{2(N+1)})$. Therefore, in view of the asymptotic behaviors (A.9) and (A.10) of w_1 as $r \rightarrow +\infty$, we can use integration by parts, to obtain:

$$\begin{aligned} &\int_0^{+\infty} \rho(r) \frac{r^{2(N+1)}}{(1+r^{2(N+1)})^2} w_1(r) r dr \\ &= \frac{1}{2} \int_0^{+\infty} L_1\left(\frac{1}{(1+r^{2(N+1)})^2}\right) w_1(r) r dr \\ &= \frac{1}{2} \int_0^{+\infty} \frac{d}{dr} \left(r \frac{d}{dr} \frac{1}{(1+r^{2(N+1)})^2} \right) w_1(r) dr + \frac{1}{2} \int_0^{+\infty} \rho(r) w_1(r) \frac{1}{(1+r^{2(N+1)})^2} r dr \\ &= \frac{1}{2} \int_0^{+\infty} \frac{d}{dr} \left(r \frac{d}{dr} w_1(r) \right) \frac{1}{(1+r^{2(N+1)})^2} dr + \frac{1}{2} \int_0^{+\infty} \rho(r) w_1(r) \frac{1}{(1+r^{2(N+1)})^2} r dr \\ &= \frac{1}{2} \int_0^{+\infty} L_1 w_1 \frac{1}{(1+r^{2(N+1)})^2} r dr = -\frac{1}{2} \int_0^{+\infty} \frac{e^{-c_0r^2}}{1+r^{2(N+1)}} r dr. \end{aligned}$$

Consequently,

$$\int_0^{+\infty} \left(\frac{e^{-c_0r^2} r^{2(N+1)}}{1+r^{2(N+1)}} - \frac{2\rho(r)w_1(r)r^{2(N+1)}}{(1+r^{2(N+1)})^2} \right) r dr = \int_0^{+\infty} e^{-c_0r^2} r dr = \frac{1}{2c_0}. \quad \square$$

References

- [1] A.A. Abrikosov, On the magnetic properties of superconductors of second group, *Sov. Phys. JETP* 5 (1957) 1174–1182.
- [2] J. Ambjorn, P. Olesen, A magnetic condensate solution of the classical electroweak theory, *Phys. Lett. B* 218 (1989) 67–71.
- [3] J. Ambjorn, P. Olesen, On electroweak magnetis, *Nucl. Phys. B* 315 (1989) 606–614.
- [4] J. Ambjorn, P. Olesen, A condensate solution of the electroweak theory which interpolates between the broken and symmetry phase, *Nucl. Phys. B* 330 (1990) 193–204.
- [5] D. Bartolucci, G. Tarantello, The Liouville equations with singular data and their applications to electroweak vortices, *Comm. Math. Phys.* 229 (2002) 3–47.
- [6] H. Brezis, F. Merle, Uniform estimates and blow-up behaviour for solutions of $-\Delta u = V(x)e^u$ in two dimensions, *Comm. Partial Differential Equations* 16, (8,9), 1223–1253.
- [7] D. Chae, O.Yu. Imanuvilov, The existence of non-topological multivortex solutions in the relativistic self-dual Chern–Simons theory, *Comm. Math. Phys.* 215 (2000) 119–142.
- [8] W. Chen, C. Li, Qualitative properties of solutions to some nonlinear elliptic equations in \mathbb{R}^2 , *Duke Math. J.* 71 (2) (1993) 427–439.
- [9] G. 't Hooft, A property of electric and magnetic flux in nonabelian gauge theories, *Nucl. Phys. B* 153 (1979) 141–160.
- [10] C.H. Lai (Ed.), *Selected Papers on Gauge Theory of Weak and Electromagnetic Interactions*, World Scientific, Singapore.
- [11] L. Nirenberg, *Topics in Nonlinear Analysis*, in: *Courant Lecture Notes in Math.*, American Mathematical Society, 2001.
- [12] J. Prajapat, G. Tarantello, On a class of elliptic problems in \mathbb{R}^2 : symmetry and uniqueness results, *Proc. Royal Soc. Edinburgh* 131 (4) (2001) 967–985.
- [13] J. Spruck, Y. Yang, On multivortices in the electroweak theory I: existence of periodic solutions, *Comm. Math. Phys.* 144 (1992) 1–16.
- [14] J. Spruck, Y. Yang, On multivortices in the electroweak theory II: existence of Bogomol'nyi solutions in \mathbb{R}^2 , *Comm. Math. Phys.* 144 (1992) 215–234.
- [15] C.H. Taubes, Arbitrary N -vortex solutions to the first order Ginzburg–Landau equation, *Comm. Math. Phys.* 72 (1980) 277–292.
- [16] C.H. Taubes, On the equivalence of first order and second order equations for gauge theories, *Comm. Math. Phys.* 75 (1980) 207–227.
- [17] Y. Yang, *Solitons in Field Theory and Nonlinear Analysis*, in: *Springer Monographs in Math.*, Springer-Verlag, New York, 2001.