G. RANGAN

***p*-adic almost periodicity and representations**

Annales mathématiques Blaise Pascal, tome 2, n° 1 (1995), p. 237-243

http://www.numdam.org/item?id=AMBP_1995__2_1_237_0

© Annales mathématiques Blaise Pascal, 1995, tous droits réservés.

L'accès aux archives de la revue « Annales mathématiques Blaise Pascal » (<http://math.univ-bpclermont.fr/ambp/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

P-ADIC ALMOST PERIODICITY

AND REPRESENTATIONS

G. Rangan

Abstract– In the first international conference on p -adic functional analysis, the question whether it is possible to get the structure of the Banach Algebra $A_c(G)$ of p -adic valued continuous almost periodic functions on a totally disconnected topological IB-group G through the structure of its non-archimedean Bohr compactification $\hat{G}$ was raised. We affirmatively answer this question here. This structure of $A_c(G)$ helps one to study the p -adic regular representation of G using the known theory of representations for compact groups.

1991 Mathematics subject classification: 46S10

1 Introduction

Let G be a group and K a complete ultra-metric valued field. When G carries a topology under which G is a topological group, we have studied in earlier papers Rangan [5], [6], [7] and [8] continuous almost periodic functions on G with values in K . In Rangan [8] we conjectured that a structure theory for the Banach algebra $A = A_c(G)$ of continuous almost periodic functions on G can be obtained using the known structure theory of the group algebra of a compact group by going to the Bohr compactification $\hat{G}$ of G . In this paper we give an affirmative answer to the conjecture. The observation that G is an IB-group if and only if the Bohr compactification $\hat{G}$ is an IB-group or equivalently a p -free group, where p is the characteristic of the residue class field of K , which is implicitly contained in the results proved in Rangan [7], helps us to establish the conjecture.

When G is an arbitrary group and K is a locally compact field we consider the subgroup topology on G defined by the normal subgroups of finite index in G under which

G becomes a O -dimensional group. The space of continuous almost periodic functions on G described above coincides with the space of almost periodic functions $AP(G \rightarrow K)$ defined by Schikof [10] using compactoid. This enables us to prove that there exists an invariant mean on $AP(G \rightarrow K)$ or equivalently the pair (G, K) is a.p.i.m. in the sense of Diarra [2](p.23, N.B.(i)) if and only if G is a IB-group or equivalently a p -free group (see Rangan [7]). Thus in the case when the base field is locally-compact, the problem of characterising (G, K) pairs which are a.p.i.m posed by Diarra is solved. The problem still remains open for non-locally compact fields. This also gives rise to the structure theory for $AP(G \rightarrow K)$ which is got by going to its Bohr compactification.

The structure theory so arrived at for the algebra of almost periodic functions gives rise to a study of representations of G taking the base space for representation to be the space of almost periodic functions on G . This may give rise to an alternative approach to representation theory developed by Diarra [1] using Hopf algebras. We intend discussing the details in another paper. Using the structure theory of $AP(G \rightarrow K)$, we prove that the regular representation decomposes as a direct sum of finite-dimensional representations.

2 Notations and Definitions

G is a group and K is a complete ultra metric rank one valued field, p denotes the characteristic of the residue class field. For $f : G \rightarrow K$, $x, s \in G$ we put $f_s(x) := f(s^{-1}x)$, $f^s(x) := f(xs)$, $f^v(x) := f(x^{-1})$, $f_G = \{f_s : s \in G\}$ and $f^G := \{f^s : s \in G\}$. A function f defined on G is called almost periodic if f_G is pre-compact or equivalently if for every $\epsilon > 0$ there exists a covering of G by a finite collection of subsets $A_1, A_2, \dots, A_n$ such that for $x, y \in A_i$ for $i = 1, 2, \dots, n$ $|f(cxd) - f(cyd)| < \epsilon$ for all $c, d \in G$ (See Maak [4]). Interestingly it turns out that for a given $\epsilon > 0$ and an almost periodic function f on G , the covering consisting of minimum number of subsets $A_1, A_2, \dots, A_n$ such that for $x, y \in A_i$, $|f(cxd) - f(cyd)| < \epsilon$ for $i = 1, 2, \dots, n$ is the covering by cosets of a suitable normal subgroup $H(f, \epsilon)$ called the ϵ -kernel of finite index n in G . If f is a continuous almost periodic function on a topological group G , $H(f, \epsilon)$ is also an open and closed subgroup of finite index in G . A (topological) group is called an IB-group (Index Bounded group) if $\inf |n| > 0$, as n varies over all the indices of (closed) subgroups of finite index of G . We take $c = \inf |n|$. G is p -free if only if $c = 1$ or equivalently $|n| = 1$ for each index n . There exists a Mean M with $\|M\| = 1$ (sup norm) on $A_c(G)$ if and only if G is a p -free group.

Schikhof [10] calls a function $f : G \rightarrow K$ almost periodic if f_G is a compactoid in $B(G, K)$, the space of bounded functions on G with the supremum norm. The set of all almost periodic functions from G to K is denoted by $AP(G \rightarrow K)$. The almost periodic functions which are analogous of the classical case discussed earlier are called strictly almost periodic and the space of such functions is denoted by $SAP(G \rightarrow K)$. When G is a topological group the space of continuous strictly almost periodic functions is the space $A_c(G)$ of

the earlier papers of the author. In general $SAP(G \rightarrow K) \subset AP(G \rightarrow K)$; however when the base field is locally compact $SAP(G \rightarrow K) = AP(G \rightarrow K)$. Diarra [1] has shown that χ_N the characteristic function of a normal subgroup N belongs to $AP(G \rightarrow K)$ if and only if N is of finite index in G .

3 Existence of Mean

Theorem 3.1 *If G is a topological 0-dimensional group then G is an IB-group if and only if its Bohr compactification $\hat{G}$ is an IB-group or equivalently a p -free group.*

Proof: Let G be an IB-group. Then Theorem 3.3. [5] implies that there exists a Mean M on $A_c(G)$. Again by Theorem 3.8. [7] M defines an invariant integral for continuous functions on $\hat{G}$ and so $\hat{G}$ is a p -free group or equivalently an IB-group.

Conversely if $\hat{G}$ is an IB-group or equivalently a p -free group, the integral on $\hat{G}$ induces an invariant mean on $A_c(G)$. and so G is a p -free group or an IB-group with $c = 1$. ■

Remark 1: When G is compact the collection of open and closed subgroups coincides with the collection of closed subgroups of finite index in G and so the p -free condition in the usual sense coincides with the IB-condition on G .

Remark 2: When the base field K is locally-compact Diarra has given (corollary 2, p.13, [1]) several equivalent criteria for the existence of mean on $AP(G \rightarrow K)$ in terms of almost periodic representations, existence of Haar measure on the Bohr compactification etc. The above theorem which gives a criterion for the existence of mean in $AP(G \rightarrow K)$ enables one to conclude that Diarra's equivalent formulations holds when and only when the group is p -free.

If G is an arbitrary group. Let τ_B be the subgroup topology on G for which the collection of all normal subgroups of finite index is a fundamental system of neighbourhoods at the identity of G . With this topology, G is a topological group.

Proposition 3.2 *When K is locally compact and G is an arbitrary group, $AP(G \rightarrow K) = SAP(G \rightarrow K) = A_c(G)$, where $A_c(G)$ is the space of all continuous (in the subgroup topology defined above) of almost periodic functions in the sense of Maak.*

Proof: When K is locally compact every closed bounded subset of K is compact and so $SAP(G \rightarrow K) = AP(G \rightarrow K)$ (See Schikhof [10], p.3); clearly $A_c(G) \subset AP(G \rightarrow K)$. If $f \in AP(G \rightarrow K)$, $f \in SAP(G \rightarrow K)$. Hence for $\epsilon > 0$, there exists a normal subgroup of finite index $H = H(f, \epsilon)$ such that

- (i) $G = \cup_{i=1}^n Hx_i, x_i \in G$
- (ii) for $x, y \in Hx_i, i = 1, 2, \dots, n$
- $$|f(cxd) - f(cyd)| < \epsilon \text{ for all } c, d \in G.$$

In particular for $x, y \in H, |f(x) - f(y)| < \epsilon$, i.e. f is uniformly continuous with respect to the subgroup topology τ_B on G and so $f \in A_c(G)$. This proves the proposition. ■

The next theorem gives a necessary and sufficient condition for the existence of Mean on $AP(G \rightarrow K)$ in tune with the earlier conditions for the existence of Haar measure etc. (see van Rooij [8]) where G is an arbitrary group which solves the problem posed by Schikhof [10] in the case of the locally compact base field K . See also Diarra [1] theorem 4 and Schikhof [10], Theorem 8.2.

Theorem 3.3 *Let K be a locally compact field. An invariant Mean M on $AP(G \rightarrow K)$ exists if and only if G is p -free.*

Proof: We consider the subgroup topology τ_B on G given by the normal subgroups of finite index as a neighbourhood base at the identity. By the earlier proposition 3.2, $AP(G \rightarrow K) = SAP(G \rightarrow K) = A_c(G)$. Now the Theorem follows from Theorem 3.3 of Rangan [5]. ■

Example: Let G be any free-group. Then for every $x \in G, x$ different from the identity of G , there exists a normal subgroup of finite index $N, x \notin N$. (See Hewitt and Ross [3]). Hence the subgroup topology on G given by the family of normal subgroups of finite index as a neighbourhood base is a Hausdorff topology on G . Hence $AP(G \rightarrow K) = SAP(G \rightarrow K) = A_c(G)$. G is a maximally almost periodic group. An invariant Mean exists on $AP(G \rightarrow K)$ if and only if G is p -free.

Remark: When K is locally compact for the study of continuous almost periodic functions on a totally disconnected topological group, only the topology τ_B on G matters. For if (G, τ) be a totally-disconnected topological group. G is a totally disconnected topological group also with respect to the topology τ_B defined by closed (in τ) normal subgroups of finite index in (G, τ) . The topology τ_B is weaker than τ . By Theorem 4.1 Rangan [6], and proposition 3.2 above it follows that $A_c(G, \tau) = A_c(G, \tau_B) = AP(G \rightarrow K)$.

4 Structure of $A = A_c(G)$

Throughout this section we assume that K is locally compact and G is either a totally disconnected topological group or an arbitrary group G considered as a topological

group with respect to the subgroup topology τ_B defined by the normal subgroups of finite index in G . So $A_c(G, \tau) = A_c(G, \tau_B) = AP(G \rightarrow K)$. We assume G to be a p -free group.

Theorem 4.1 *The algebra $A = A_c(G)$ is isometrically isomorphic to the group algebra $L(\hat{G})$ of the Bohr compactification $\hat{G}$ of G .*

Proof: The map $\theta : A \rightarrow L(\hat{G})$ given by $f \rightarrow \hat{f}$ where $\hat{f}$ is the associated continuous function on the compact group $\hat{G}$ to f (see Rangan [6], Theorem 4.4). If ρ is the homomorphism which imbeds G in $\hat{G}$, for $x \in G$, $f(x) = \hat{f}(\rho(x))$. θ is one-to-one: For $\theta(f) = \theta(g) \Rightarrow \hat{f} = \hat{g} \Rightarrow f(x) = g(x)$ for all $x \in G \Rightarrow f = g$. θ is onto: if $h \in L(\hat{G})$, h is a continuous function on $\hat{G}$. Define $f(x) = h(\rho(x))$ for $x \in G$ then $\hat{f} = h$. θ is an algebra homomorphism: For

$$\begin{aligned} f * g(x) &= M_y(f(y)g(y^{-1}x)) \\ &= \int_G f(y)g(y^{-1}x)dy = \hat{f} * \hat{g}(x) \end{aligned}$$

where the integral is the Haar integral and it exists since $\hat{G}$ is p -free, G being so.

θ is an isometry: When G is p -free $|n| = 1$ for every normal subgroup of finite index and so $c = 1$. Hence for $f \in A$,

$$\|f\| = \sup_{x \in G} |f(x)| = \sup_{x \in G} |\hat{f}(\rho(x))| = \sup_{x \in \hat{G}} |\hat{f}(x)|$$

since $\rho(G)$ is dense in $\hat{G}$. ■

Proposition 4.2 *A is the closure of the K -linear span of the idempotents of A .*

Proof: Since $A = A_c(G) = A_c(G, \tau_B) = AP(G \rightarrow K) = SAP(G \rightarrow K)$ the proposition follows from Lemma 4.4, Schikhof [10], which is now easily seen to be a restatement of the approximation Theorem 7.4 of Rangan [5]. ■

Theorem 4.3 *For a p -free group G , $A = \oplus A_e$ where $A_e = e * A$ is a finite-dimensional two sided ideal of A and for every $f \in A$,*

$$f = \sum_{e \in E} e * f \text{ and } \|f\| = \sup_{e \in E} \|e * f\|$$

and every non-zero minimal two sided ideal in A is an A_e for a suitable $e \in E$. If I is a closed two sided ideal in A then

$$I = cl \sum_{e \in I} A_e$$

where E is the set of all minimal non-zero central idempotents of A .

Proof: Follows from 8.14 Theorem van Rooij [9] since by the earlier theorem A and $L(\hat{G})$ are isometrically isomorphic. ■

It is not difficult to prove, using the existence of the approximate identity (U_H) , (H varying over the collection Γ'_G of normal subgroups of finite index in G) that the closed ideals in A are same as closed invariant subspaces. For $f \in A$, defining $(L_a f)(x) = f(a^{-1}x)$ for $x \in G$, we get the (left) regular representation $a \rightarrow L_a$ on G . A_e being invariant subspaces in view of Theorem 4.3, L_a decomposes as a direct sum of finite-dimensional representations. Thus we get the following result.

Theorem 4.4 *The regular Representation decomposes as a direct sum of finite-dimensional representations.*

References

- [1] Diarra, B. : Ultrametric almost periodic Linear representations (Preprint).
- [2] Diarra, B. : On reducibility of ultrametric almost periodic linear representation, (Preprint)
- [3] Hewitt, E. and K.A. Ross. : *Abstract Harmonic Analysis* Vol.1 (Springer-Verlag, Berlin, 1963)
- [4] Maak, W. : *Fast periodische Funktionen*, Berlin - Göttingen - Heidelberg, Springer 1950.
- [5] Rangan, G. : Non-archimedean valued almost periodic functions, *Indag.Math.*, **31** (1969), 345-353.
- [6] Rangan, G. : Non-archimedean Bohr Compactification of a topological group, *Indag.Math.*, **31** (1969), 354-360.
- [7] Rangan, G. : On the existence of non-archimedean valued invariant Mean *Publ. Math.* (Debrecen) **29** (1982), 57-63.
- [8] Rangan, G. and M.S.Saleemullah. : Banach algebra of p-adic valued almost periodic functions: *p-adic Functional Analysis*, (Eds. J.M. Bayod, N.De Grade-De Kimpe and J. Martinez-Maurica), Marcel Dekker, New York 1991, 141 - 150.

- [9] Rooij van, A.C.M. : *Non-archimedean Functional Analysis*, Marcel-Dekker-New York, 1978.
- [10] Schikhof, W.H. : An approach to p-adic almost periodicity by Means of Compactoids, *Report 8809*, Department of Mathematics, Catholic University, Nijmegen, 1988.

RANGAN, G.
University Building
Chepauk, Triplicane P.O.
MADRAS, 600005
INDIA, INDE