

ANNALES SCIENTIFIQUES DE L'É.N.S.

NANHUA XI

The based ring of the lowest two-sided cell of an affine Weyl group. II

Annales scientifiques de l'É.N.S. 4^e série, tome 27, n° 1 (1994), p. 47-61

http://www.numdam.org/item?id=ASENS_1994_4_27_1_47_0

© Gauthier-Villars (Éditions scientifiques et médicales Elsevier), 1994, tous droits réservés.

L'accès aux archives de la revue « Annales scientifiques de l'É.N.S. » (<http://www.elsevier.com/locate/ansens>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

THE BASED RING OF THE LOWEST TWO-SIDED CELL OF AN AFFINE WEYL GROUP, II

BY NANHUA XI ⁽¹⁾

ABSTRACT. — We show that the lowest based ring of an affine Weyl group W is very interesting to understand some simple representations of the corresponding Hecke algebra H_{q_0} ($q_0 \in \mathbb{C}^*$) even when q_0 is a root of 1.

Let H_{q_0} be the Hecke algebra (over \mathbb{C}) attached by Iwahori and Matsumoto [IM] to an affine Weyl group W and to a parameter $q_0^2 \in \mathbb{C}^*$.

When q_0 is not a root of 1 or $q_0^2 = 1$, the simple H_{q_0} -modules have been classified (see [KL2]). However we know little about the simple H_{q_0} -modules when q_0 is a root of 1. In this paper we give some discussion to the representations of H_{q_0} with q_0 a root of 1. Namely, let J be the asymptotic Hecke algebra defined in [L3, III]. There exists a natural injection $\phi_{q_0}: H_{q_0} \rightarrow J$. Let $K(J)$ [resp. $K(H_{q_0})$] be the Grothendieck group of J -modules (resp. H_{q_0} -modules) of finite dimension over \mathbb{C} , then ϕ_{q_0} induces a surjective homomorphism $(\phi_{q_0})_*: K(J) \rightarrow K(H_{q_0})$, when q_0 is not a root of 1 or $q_0^2 = 1$, $(\phi_{q_0})_*$ is an isomorphism (*loc. cit.*). For each two-sided cell c of W , we can define the direct summand $K(J_c)$ [resp. $K(H_{q_0,c})$] of $K(J)$ [resp. $K(H_{q_0})$]. Thus $(\phi_{q_0})_*$ induces a homomorphism $(\phi_{q_0})_{*,c}: K(J_c) \rightarrow K(H_{q_0,c})$. The map $(\phi_{q_0})_{*,c}$ remains surjective and is an isomorphism if q_0 is not a root of 1 or $q_0^2 = 1$. In this paper we mainly discuss the map $(\phi_{q_0})_{*,c_0}$, where c_0 is the lowest two-sided cell of W .

1. Introduction

1.1. Let G be a simply connected, almost simple complex algebraic group and T a maximal torus. Let $P \subseteq X = \text{Hom}(T, \mathbb{C}^*)$ be the root lattice. The Weyl group $W_0 = N_G(T)/T$ of G acts on X in a natural way and this action is stable on P . Thus we can form the affine Weyl group $W_a = W_0 \ltimes P$, which is a normal subgroup of the extended affine Weyl group $W = W_0 \ltimes X$. There exists a finite abelian subgroup Ω of W such that $W = \Omega \ltimes W_a$. Let $S = \{r_0, r_1, \dots, r_n\}$ be the set of simple reflections of W_a with $r_0 \notin W_0$. Then we have a standard length function l on W_a which can be extended

⁽¹⁾ Supported in part by an N.S.F. Grant (DMS 9100383).

to W by defining $l(\omega w) = l(w)$ for any $\omega \in \Omega$, $w \in W_a$. We keep the same notation for the extension of l .

1.2. For any $u = \omega_1 u_1$, $w = \omega_2 w_1$, $\omega_1, \omega_2 \in \Omega$, $u_1, w_1 \in W_a$, we define $P_{u, w}$ to be P_{u_1, w_1} , as in [KL 1] if $\omega_1 = \omega_2$ and $P_{u, w}$ to be zero if $\omega_1 \neq \omega_2$. We say that $u \underset{LR}{\leq} w$ or $u \underset{L}{\leq} w$ if $u_1 \underset{LR}{\leq} w_1$, or $u_1 \underset{L}{\leq} w_1$ in the sense of [KL 1], we say that $u \underset{R}{\leq} w$ if $u^{-1} \underset{L}{\leq} w^{-1}$. These relations generate equivalence relations $\underset{LR}{\sim}$, $\underset{L}{\sim}$, $\underset{R}{\sim}$ in W , respectively, and the corresponding equivalence classes are called two-sided cells, left cells, right cells of W , respectively. The relation $\underset{LR}{\leq}$ (resp. $\underset{L}{\leq}$, $\underset{R}{\leq}$) in W then induces a partial order $\underset{LR}{\leq}$ (resp. $\underset{L}{\leq}$, $\underset{R}{\leq}$) in the set of two-sided (resp. left, right) cells of W . We extend the Bruhat order \leq in W_a to W by defining $u \leq w$ if and only if $\omega_1 = \omega_2$ and $u_1 \leq w_1$.

Let q be an indeterminate and let $A = \mathbb{C}[q, q^{-1}]$. Let H be the Hecke algebra of W over A , that is a free A -module with basis $T_w (w \in W)$ and multiplication defined by

$$(T_r - q^2)(T_r + 1) = 0 \quad \text{if } r \in S \quad \text{and} \quad T_w T_{w'} = T_{ww'} \quad \text{if } l(ww') = l(w) + l(w').$$

For each $w \in W$, let

$$C_w = q^{-l(w)} \sum_{u \leq w} P_{u, w}(q^2) T_u \in H.$$

And we write

$$C_w C_u = \sum_z h_{w, u, z} C_z \in H, \quad h_{w, u, z} \in A.$$

For each $z \in W$, there is a well defined integer $a(z) \geq 0$ such that

$$\begin{aligned} q^{a(z)} h_{w, u, z} &\in \mathbb{C}[q] \quad \text{for all } w, u \in W \\ q^{a(z)-1} h_{w, u, z} &\notin \mathbb{C}[q] \quad \text{for some } w, u \in W \end{aligned}$$

(see [L 3, I, 7.3]). We have $a(z) \leq l(w_0)$, where w_0 is the longest element of W_0 . It is known that

$$c_0 = \{ w \in W \mid a(w) = l(w_0) \}$$

is a two-sided cell of W (see [S, I]) which is the lowest one for the partial order $\underset{LR}{\leq}$.

1.3. Let $\gamma_{w, u, z}$ be the constant term of $q^{a(z)} h_{w, u, z} \in \mathbb{C}[q]$. We have $\gamma_{w, u, z} \in \mathbb{N}$. Moreover (see [L 3, II])

$$(a) \quad \gamma_{w, u, z} \neq 0 \quad \Rightarrow \quad \underset{L}{w} \sim \underset{L}{u}^{-1}, \quad \underset{L}{u} \sim \underset{L}{z}, \quad \underset{R}{w} \sim \underset{R}{z}.$$

Let J be the \mathbb{C} -vector space with basis $(t_w)_{w \in W}$. This is an associative \mathbb{C} -algebra with multiplication

$$t_w t_u = \sum_z \gamma_{w, u, z} t_z.$$

It has a unit element $1 = \sum_{d \in \mathcal{D}} t_d$, where $\mathcal{D} = \{d \in W_a \mid a(d) = l(d) - 2 \deg P_{e,d}\}$ (e is the unit of W) (see [L 3, II]).

For each two-sided cell c of W , let J_c be the subspace of J spanned by t_w , $w \in c$, then $J = \bigoplus_c J_c$, where the sum is over the set of all two-sided cells of W . By (a) we see that J_c is a two-sided ideal of J and in fact is an associative \mathbb{C} -algebra with unit $\sum_{d \in \mathcal{D} \cap c} t_d$.

1.4. For each $q_0 \in \mathbb{C}^*$, we denote $H_{q_0} = H \otimes_A \mathbb{C}$, where \mathbb{C} is an A -algebra with q_0 acting as scalar multiplication by q_0 . We shall denote $T_w \otimes 1$, $C_w \otimes 1$ in H_{q_0} again by T_w , C_w . We also use the notation $h_{w,u,z}$ for the specialization at $q_0 \in \mathbb{C}^*$ of $h_{w,u,z}$.

The A -linear map $\phi: H \rightarrow J \otimes_{\mathbb{C}} A$ defined by

$$\phi(C_w) = \sum_{\substack{d \in \mathcal{D} \\ z \in W \\ a(z) = a(d)}} h_{w,d,z} t_z$$

is a homomorphism of A -algebra with 1 (see [L 3, II]). Let $\phi_{q_0}: H_{q_0} \rightarrow J$ be the induced homomorphism for any $q_0 \in \mathbb{C}^*$.

Any (left) J -module E gives rise, via $\phi_{q_0}: H_{q_0} \rightarrow J$, to a (left) H_{q_0} -module E_{q_0} . We denote by $K(J)$ [resp. $K(H_{q_0})$] the Grothendieck group of (left) J -modules (resp. H_{q_0} -modules) of finite dimension over \mathbb{C} . The correspondence $E \rightarrow E_{q_0}$ defines a homomorphism $(\phi_{q_0})_*: K(J) \rightarrow K(H_{q_0})$.

We similarly define $K(J_c)$ for any two-sided cell c of W . Then we have $K(J) = \bigoplus_c K(J_c)$, where the sum is over the set of all two-sided cells of W . Now we define $K(H_{q_0})_c$. For any simple H_{q_0} -module M , we attach to M a two-sided cell c_M of W by the following two conditions:

$$\begin{aligned} C_w M &\neq 0 \text{ for some } w \in c_M \\ C_w M &= 0 \text{ for any } w \text{ in a two-sided cell } c \text{ with } c \underset{\text{LR}}{\leq} c_M, c \neq c_M. \end{aligned}$$

Then c_M is well defined since there are only a finite number of two-sided cells in W . Let $K(H_{q_0})_c$ be the subgroup of $K(H_{q_0})$ spanned by simple H_{q_0} -modules M with $c_M = c$. Obviously we have $K(H_{q_0}) = \bigoplus_c K(H_{q_0})_c$. Thus for a two-sided cell c of W ,

$(\phi_{q_0})_*$ induces a homomorphism

$$(\phi_{q_0})_{*,c}: K(J_c) \rightarrow K(H_{q_0})_c.$$

The following result is due to Lusztig (see [L 3, III, 1.9 and 3.4]).

PROPOSITION 1.5. — *The map $(\phi_{q_0})_{*,c}$ is surjective for any $q_0 \in \mathbb{C}^*$, moreover, $(\phi_{q_0})_{*,c}$ is an isomorphism when q_0 is not a root of 1 or $q_0^2 = 1$.*

Now we state a conjecture.

CONJECTURE 1.6. — The map $(\phi_{q_0})_{*,c}$ is injective if $(\phi_{q_0})_{*,c'}$ is injective for some two-sided cell c' of W with $c' \leq c$.
LR

By proposition 1.6 one knows that $(\phi_{q_0})_{*,c}$ is injective is equivalent to that $(\phi_{q_0})_{*,c}$ is bijective.

We mainly discuss $(\phi_{q_0})_{*,c_0}$, where c_0 is the lowest two-sided cell of W . We prove that if $\sum_{w \in W_0} q_0^{2l(w)} \neq 0$, then $(\phi_{q_0})_{*,c_0}$ is injective (see Theorem 3.4) and show that $(\phi_{q_0})_{*,c_0}$ is likely not injective if $\sum_{w \in W_0} q_0^{2l(w)} = 0$ (see Theorem 3.6).

1.7. Let H'_{q_0} be the subalgebra of H_{q_0} spanned by T_w , $w \in W_0$. And let J' be the subspace of J spanned by t_w , $w \in W_0$. J' is a \mathbb{C} -algebra with unit $\sum_{d \in \mathcal{D} \cap W_0} t_d$. Let

$\phi'_{q_0}: H'_{q_0} \rightarrow J'$ be defined by

$$\phi'_{q_0}(C_w) = \sum_{\substack{d \in \mathcal{D} \cap W_0 \\ z \in W_0 \\ a(d) = a(z)}} h_{w,d,z}(q_0) t_z, \quad w \in W_0,$$

then ϕ'_{q_0} is a \mathbb{C} -algebra homomorphism preserving 1.

As in 1.4 we define $K(H'_{q_0})$, $K(J')$, $K(H'_{q_0/c'})$, $K(J'_c')$, $(\phi'_{q_0})_{*,c}$, $(\phi'_{q_0})_{*,c'}$, etc., where c' is a two-sided cell of W_0 . We also have

PROPOSITION 1.8. — $(\phi'_{q_0})_{*,c'}$ is surjective for any $q_0 \in \mathbb{C}^*$. Moreover $(\phi_{q_0})_{*,c'}$ is an isomorphism when q_0 is not a root of 1 or $q_0^2 = 1$.

CONJECTURE 1.9. — $(\phi_{q_0})_{*,c'}$ is injective if $(\phi'_{q_0})_{*,c'}$ is injective for some two-sided cell c'' of W_0 with $c'' \leq c'$.
LR

When c' is the lowest two-sided cell of W_0 , it is easy to see that $(\phi'_{q_0})_{*,c'}$ is injective if and only if $\sum_{w \in W_0} q_0^{2l(w)} \neq 0$.

2. The two-side cell c_0 and the ring J_{c_0}

In this section we recall and prove some results on c_0 and J_{c_0} .

2.1. We denote by w_0 the longest element in W_0 . Let

$$\mathfrak{S} = \{w \in W \mid l(w w_0) = l(w) + l(w_0) \quad \text{and} \quad w w_0 r \notin c_0 \quad \text{for any } r \in S \cap W_0\}.$$

Then $\mathcal{D}_0 = \mathcal{D} \cap c_0 = \{w w_0 w^{-1} \mid w \in \mathfrak{S}\}$ and $|\mathfrak{S}| = |W_0|$ (see [S, II]).

Let $X^+ = \{w \in W \mid l(rx) > l(x) \text{ for any } r \in S'\}$, where $S' = S \cap W_0$. Let $x_i \in X^+$ ($i \in \{1, 2, \dots, n\} = I_0$) be the i -th basic dominant weight, then x_i has the properties: $l(x_i r_i) < l(x_i)$, $x_i r_j = r_j x_i$, $l(x_i r_j) = l(x_i) + 1$ if $i \neq j \in I_0$. We have

$$c_0 = \{w' w_0 x w^{-1} \mid w, w' \in \mathfrak{S}, x \in X^+\} \quad (\text{see [S, II]}).$$

Moreover $l(w' w_0 x w^{-1}) = l(w') + l(w_0) + l(x) + l(w^{-1})$.

LEMMA 2.2. — *Let $u \in c_0$, then $C_u = h C_{w_0} h'$ for some $h, h' \in H_{q_0}$, i. e., the two-sided ideal $\bigoplus_{u \in c_0} \mathbb{C} C_u$ of H_{q_0} is generated by the element C_{w_0} .*

Proof. — Write $u = w' w_0 w$ for some $w', w \in W$ such that $l(u) = l(w') + l(w_0) + l(w)$. We use induction on $l(u)$ to prove that C_u is in the two-sided ideal N of H_{q_0} generated by C_{w_0} .

When $l(u) = l(w_0)$, then $C_u = C_\omega C_{w_0} C_{\omega'}$ for some $\omega, \omega' \in \Omega$. Now assume that $l(w') > 0$. Let $s \in S$ be such that $sw' \leq w'$, then

$$C_s \cdot C_{su} = C_u + \sum_{\substack{z \in c_0 \\ l(z) < l(u)}} a_z C_z, \quad a_z \in \mathbb{N} \quad (\text{see [KL 1]}).$$

By induction hypothesis we know that $C_u \in N$. Similarly we can prove that $C_u \in N$ if $l(w) > 0$. The lemma is proved.

COROLLARY 2.3. — *For a simple H_{q_0} -module M , we have $c_M = c_0$ if and only if $C_{w_0} M \neq 0$.*

For $w \in W$, set $L(w) = \{r \in S \mid rw \leq w\}$ and $R(w) = \{r \in S \mid wr \leq w\}$.

LEMMA 2.4. — (i) *Let w' be the longest element in the Weyl group generated by $L(w)$ (or $R(w)$), then $w = w' w''$ (or $w = w'' w'$) for some $w'' \in W$ and $l(w) = l(w') + l(w'')$.*

(ii) *Let w' be the longest element in the Weyl group W' generated by $S - L(w)$ [resp. $S - R(w)$], then $l(w' w) = l(w') + l(w)$ [resp. $l(w w') = l(w) + l(w')$].*

Proof. — (i) follows from $T_w C_w = q^{l(w')} C_w$ or $C_w T_w = q^{l(w')} C_w$.

(ii) follows from the fact that w is the shortest element in $W' w$ or $w W'$.

Let Γ_0 be the left cell in c_0 containing w_0 , then

$$\begin{aligned} \Gamma_0 &= \{w w_0 x \mid x \in X^+, w \in \mathfrak{S}\} \\ &= \{w \in W \mid R(w) = S'\} \end{aligned}$$

LEMMA 2.5. — *Any element $u \in \Gamma_0$ has the form $w x w_j$, where $w \in W_0$, $x = \prod_{i=1}^n x_i^{a_i} \in X^+$. w_j is the longest element in $W_j = \langle r_j \mid a_j = 0, j \in I_0 \rangle$, moreover $l(u) = l(w) + l(x) + l(w_j)$.*

Proof. — Choose $x = \prod_{i=1}^n x_i^{a_i} \in X^+$ such that $u \in \Gamma_0 \cap W_0 x W_0$.

Then the shortest element in $W_0 x W_0$ is $xw_j w_0$ and the shortest element in $\Gamma_0 \cap W_0 x W_0$ is xw_j by lemma 2.4 (i), where w_j is the longest element in $W_J = \langle r_j \mid a_j = 0, j \in I_0 \rangle$. The lemma is proved.

LEMMA 2.6. – (i) Let $J \subseteq K \subseteq I_0$, then in H_{q_0} we have $C_{w_j} C_{w_K} = C_{w_K} C_{w_j} = \eta_J C_{w_K}$, where w_j, w_K are the longest element in $W_J = \langle r_j \mid j \in J \rangle, W_K = \langle r_k \mid k \in K \rangle$, respectively, $\eta_J = q_0^{-l(w_j)} \sum_{w \in W_J} q_0^{2l(w)}$.

(ii) $C_{ww_j} = h C_{w_j}, C_{w_j w'} = C_{w_j} h'$ for some $h, h' \in H_{q_0}$ if $l(ww_j) = l(w) + l(w_j), l(w_j w') = l(w_j) + l(w')$.

Proof. – First we prove (ii). We use induction on $l(w)$. Assume that $l(w) > 0$. Choose $r \in S$ such that $rw \leq w$, then

$$C_r C_{rw_j} = C_{ww_j} + \sum_{\substack{z \in W \\ l(z) < l(ww_j)}} a_z C_z, \quad a_z \in \mathbb{N} \quad (\text{see [KL 1]}).$$

Moreover $a_z \neq 0$ implies that $z \leq_L rw_j$. So $R(z) \supseteq \{r_j \mid j \in J\}$ (see [KL 1]).

By Lemma 2.4 we see that $z = z' w_j$ for some $z' \in W$ and $l(z) = l(z') + l(w_j)$. By induction hypothesis we know that $C_{ww_j} = h C_{w_j}$ for some $h \in H_{q_0}$. Similarly we have $C_{w_j w'} = C_{w_j} h'$ for some $h' \in H_{q_0}$.

(i) follows from $C_J C_J = \eta_J C_J$ and (ii).

COROLLARY 2.7. – Let x, w_j be as in 2.5, then in H_{q_0} we have

$$C_{w_0} C_{xw_j} = \eta_J \sum_{\substack{y \in X^+ \\ w_0 y \leq w_0 x}} a_{x,y} C_{w_0 y} \in \mathbb{C}, \quad a_{x,y} \in \mathbb{C} \quad \text{and} \quad a_{x,x} = 1.$$

Proof. – By 2.1 and 2.6(ii) we see that $C_{xw_j} = C_{w_j x} = C_{w_j} h$, where

$$h = \sum_{\substack{w \in W \\ l(w_j w) = l(w_j) + l(w) \\ w_j w \leq w_j x}} a_w T_w, \quad a_w \in \mathbb{C}, \quad a_x = q_0^{-l(x)}.$$

By (2.6(i)) we know that

$$(a) \quad C_{w_0} \cdot C_{xw_j} = C_{w_0} \cdot C_{w_j} h = \eta_J C_{w_0} h.$$

Note that $h_{w_0, xw_j, z} \neq 0$ implies that $z \sim_L xw_j, z \sim_R w_0$ (see [L 3, I]), we have $z \in \Gamma_0 \cap \Gamma_0^{-1} = \{w_0 y \mid y \in X^+\}$. So by (a) we get

$$C_{w_0} C_{xw_j} = \eta_J \sum_{y \in X^+} a_{x,y} C_{w_0 y}, \quad a_{x,y} \in \mathbb{C}.$$

Since $a_x = q_0^{-l(x)}$ and $l(w) < l(x)$ if $a_w \neq 0$, $w \neq x$. We have $a_{x,x} = 1$ and $a_{x,y} = 0$ if $l(y) > l(x)$ or $l(y) = l(x)$ but $x \neq y$. Let $w \in W$ be such that $a_w \neq 0$. Consider the expression

$$C_{w_0} \cdot T_w = \sum_{z^{-1} \in \Gamma_0} b_z C_z, \quad b_z \in \mathbb{C}.$$

Since $w_j w \leq w_j x$, we have $b_z \neq 0$ implies that $z \leq w_0 x$. Thus by (a) we know that $a_{x,y} \neq 0$ implies that $w_0 y \leq w_0 x$. The Corollary is proved.

2.8. For any $x \in X$, we choose $x', x'' \in X^+$ such that $x = x' x''^{-1}$ and then define $\tilde{T}_x = q_0^{-l(x')} T_{x'} (q_0^{-l(x'')} T_{x''})^{-1}$. \tilde{T}_x is independent of the choices x' and x'' . We denote the conjugacy class of $x \in X$ in W by O_x and let $z_x = \sum_{x' \in O_x} \tilde{T}_{x'}$. z_x is in the center of

H_{q_0} . For $x \in X^+$, denote $d(x', x)$ the dimension of the x' -weight space $V(x)_{x'}$ of $V(x)$, where $V(x)$ is the irreducible representation of G with highest weight x . We set $S_x = \sum_{x' \in X^+} d(x', x) z_{x'}$, $x \in X^+$.

LEMMA 2.9. (see [X]). — In H_{q_0} we have $C_{w' w_0 w^{-1}} S_x = S_x C_{w' w_0 w^{-1}} = C_{w' w_0 x w^{-1}}$ for any $w', w \in \mathfrak{S}$, $x \in X^+$.

LEMMA 2.10. — Let $u \in \Gamma_0$, then

$$C_u = \sum_{\substack{y \in X^+ \\ I \subseteq I_0}} h_{I,y} C_{x_I w_I} S_y,$$

where $h_{I,y} \in H'_{q_0} = \bigoplus_{w \in W_0} \mathbb{C} T_w = \bigoplus_{w \in W_0} \mathbb{C} C_w$, $x_I = \prod_{i \in I} x_i$, $I' = I_0 - I$.

Proof. — By 2.5 we see that $u = w x w_j$, where $w \in W_0$, $x = \prod_{i=1}^n x_i^{a_i}$, $J = \{j \in I_0 \mid a_j = 0\}$.

We use induction on $l(u)$, when $w = e$, by 2.9 we see that $C_u = C_{x_J w_J} S_y$, where $J' = I_0 - J$, $y = \prod_{j \in J'} x_j^{a_j - 1}$, i.e. the lemma is true. Now assume that $l(w) > 0$ and choose $r \in S'$ such

that $rw \leq w$, then

$$C_r \cdot C_{r w x w_j} = C_{w x w_j} + \sum_{\substack{z \in \Gamma_0 \\ l(z) < l(w x w_j)}} a_z C_z, \quad a_z \in \mathbb{N}.$$

By induction hypothesis we know that there exists $h_{I,y} \in H'_{q_0}$ such that $C_u = \sum_{\substack{y \in X^+ \\ I \subseteq I_0}} h_{I,y} C_{x_I w_I} S_y$. The lemma is proved.

2.11. Let R_G be the ring of the rational representations ring of G tensor with \mathbb{C} . Then R_G is a \mathbb{C} -algebra with a \mathbb{C} -basis $V(x)$, $x \in X^+$. Let $M_{\mathfrak{S} \times \mathfrak{S}}(R_G)$ be the $\mathfrak{S} \times \mathfrak{S}$ matrix

ring over R_G . Then we have

THEOREM 2.12 (see [X]). — *There is a natural isomorphism $J_{c_0} \xrightarrow{\sim} M_{\mathfrak{S} \times \mathfrak{S}}(R_G)$ such that $t_{w'w_0xw}^{-1} \rightarrow (m_{w_1, w_2}) \in M_{\mathfrak{S} \times \mathfrak{S}}(R_G)$, $w', w^{-1}, w_1, w_2 \in \mathfrak{S}$,*

$$m_{w_1, w_2} = \begin{cases} V(x) & \text{if } w_1 = w', \quad w_2 = w \\ 0 & \text{otherwise.} \end{cases}$$

Hereafter we identify J_{c_0} with $M_{\mathfrak{S} \times \mathfrak{S}}(R_G)$.

3. The homomorphism $(\phi_{q_0})_{*, c_0}$

3.1. For any semisimple conjugacy class s in G , we have a simple representation ψ_s of $J_{c_0} \simeq M_{\mathfrak{S} \times \mathfrak{S}}(R_G)$:

$$\begin{aligned} \psi_s: M_{\mathfrak{S} \times \mathfrak{S}}(R_G) &\rightarrow M_{\mathfrak{S} \times \mathfrak{S}}(\mathbb{C}) \\ (m_{w, w'}) &\rightarrow (tr(s, m_{w, w'})), \quad w, w' \in \mathfrak{S}. \end{aligned}$$

Any simple representation of J_{c_0} is isomorphic to some ψ_s (see [X]). Let E_s be the simple J_{c_0} -module providing the representation ψ_s . E_s gives rise, via

$$\phi_{q_0, c_0}: H_{q_0} \rightarrow J \rightarrow J_{c_0},$$

to an H_{q_0} -module E_{s, q_0} . Note that $\phi_{q_0, c_0}(S_x) = \sum_{w \in \mathfrak{S}} t_{ww_0xw}^{-1}$ for any $x \in X^+$, we see that S_x acts on E_{s, q_0} by scalar $tr(s, V(x))$.

PROPOSITION 3.2. — *The set $\Lambda = \{(\phi_{q_0})_{*, c_0}(E_s) \mid s \text{ semisimple conjugacy class of } G\} - \{0\}$ is a base of $K(H_{q_0})_{c_0}$.*

Proof. — It is easy to see that $(\phi_{q_0})_{*, c_0}(E_s) = \sum_M a_M M$, where the sum is over the set of composition factors M of E_{s, q_0} with $c_M = c_0$ and a_M is the multiplicity of M in E_{s, q_0} .

Now let $F_i = (\phi_{q_0})_{*, c_0}(E_{s_i}) \in \Lambda$, $1 \leq i \leq k$, and suppose that $\sum_{i=1}^k m_i F_i = 0$, $m_i \in \mathbb{Z}$. Let $F_i = \sum_{M_{ij}} a_{M_{ij}} M_{ij}$, M_{ij} simple H_{q_0} -module with $c_{M_{ij}} = c_0$. Since S_x acts on E_{s_i, q_0} by scalar $tr(s_i, V(x))$. S_x acts on M_{ij} by scalar $tr(s_i, V(x))$ if $a_{M_{ij}} \neq 0$. $F_i \in \Lambda$ implies that $a_{M_{ij}} \neq 0$ for some M_{ij} . Therefore $m_i = 0$. By 1.6 we know that $(\phi_{q_0})_{*, c_0}$ is surjective, hence Λ is a base of $K(H_{q_0})_{c_0}$. The proposition is proved.

COROLLARY 3.3. — *E_{s, q_0} has at most one composition factor to which the attached two-sided cell is c_0 . Moreover the multiplicity a_M is 1 if E_{s, q_0} has such a composition factor M .*

THEOREM 3.4. — *If $\sum_{w \in W_0} q_0^{2l(w)} = q_0^{l(w)} \eta_{1_0} \neq 0$, then $(\phi_{q_0})_{*, c_0}$ is injective, so $(\phi_{q_0})_{*, c_0}$ is an isomorphism by 1.6.*

Proof. — We have

$$\phi_{q_0, c_0}(C_{w_0}) = \sum_{\substack{w \in \mathfrak{S} \\ x \in X^+}} h_{w_0, ww_0w^{-1}, w_0xw^{-1}} t_{w_0xw^{-1}} \in J_{c_0}.$$

We identify J_{c_0} with $M_{\mathfrak{S} \times \mathfrak{S}}(\mathbb{R}_G)$, then $\phi_{q_0, c_0}(C_{w_0}) = (m_{w', w}) \in M_{\mathfrak{S} \times \mathfrak{S}}(\mathbb{R}_G)$ and

$$m_{w', w} = \begin{cases} \sum_{x \in X^+} h_{w_0, ww_0w^{-1}, w_0xw^{-1}} V(x), & \text{if } w' = e \\ 0 & \text{if } w' \neq e \end{cases}.$$

Note that $C_{w_0} C_{w_0} = \eta_{1_0} C_{w_0}$, we see that $m_{e, e} = \eta_{1_0} \neq 0$, where e is the unit in W . So $C_{w_0} E_{s, q_0} \neq 0$ for any semisimple conjugacy class s of G since $\psi_s \phi_{q_0, c_0}(C_{w_0}) \neq 0$. Now let $0 = F_0 \subseteq F_1 \subseteq \dots \subseteq F_k = E_{s, q_0}$ be a composition series of E_{s, q_0} and let i be the integer such that $C_{w_0} F_i \neq 0$ and $C_{w_0} F_{i-1} = 0$. Then $C_{w_0} M \neq 0$ where $M = F_i/F_{i-1}$, otherwise, $C_{w_0} F_i \subseteq F_{i-1}$, choose $v \in F_i$ such that $C_{w_0} v \neq 0$, we have $C_{w_0}^2 v = \eta_{1_0} C_{w_0} v \neq 0$. A contradiction, so $C_{w_0} M \neq 0$, i. e., $c_M = c_0$. That is to say $(\phi_{q_0})_{*, c_0}(E_s) \neq 0$. The theorem follows from proposition 3.2.

3.5. In the subsequent part of this section we assume that $\eta_{1_0} = 0$, i. e., $\sum_{w \in W_0} q_0^{2l(w)} = 0$.

Let $\Delta_{q_0} = \{I \subseteq I_0 \mid \eta_I \neq 0 \text{ but } \eta_{I' \cup \{i\}} = 0 \text{ for any } i \in I\}$. Here we use the convention that I' always denotes the complement of I in I_0 i. e., $I' = I_0 - I$.

THEOREM 3.6. — *Let s be a semisimple conjugacy class of G , then $(\phi_{q_0})_{*, c_0}(E_s) = 0$ if and only if $\alpha_I = 0$ for all $I \in \Delta_{q_0}$, where*

$$\alpha_I = \sum_{x \in X^+} h_{w_0, x_1 w_I, w_0 x} \text{tr}(s, V(x)) \text{ for any } I \subseteq I_0.$$

We need two lemmas.

LEMMA 3.7. — *The following conditions are equivalent.*

- (i) $C_{w_0} E_{s, q_0} = 0$.
- (ii) $\psi_s \phi_{q_0, c_0}(C_{w_0}) = 0$.
- (iii) $\sum_{x \in X^+} h_{w_0, ww_0w^{-1}, w_0xw^{-1}} \text{tr}(s, V(x)) = 0$ for all $w \in \mathfrak{S}$.
- (iv) $\sum_{x \in X^+} h_{w_0, ww_0, w_0x} \text{tr}(s, V(x)) = 0$ for all $w \in \mathfrak{S}$.
- (v) $\alpha_I = \sum_{x \in X^+} h_{w_0, x_1 w_I, w_0 x} \text{tr}(s, V(x)) = 0$ for all $I \subseteq I_0$.
- (vi) $\alpha_I = \sum_{x \in X^+} h_{w_0, x_1 w_I, w_0 x} \text{tr}(s, V(x)) = 0$ for all $I \in \Delta_{q_0}$.

Proof. — (i) and (ii) are obviously equivalent.

Note that $h_{w_0, ww_0w^{-1}, z} \neq 0$ implies that $z = w_0xw^{-1}$ for some $x \in X^+$ and that $\phi_{q_0, c_0}(C_{w_0}) = (m_{w', w})$,

$$m_{w', w} = \begin{cases} \sum_{x \in X^+} h_{w_0, ww_0w^{-1}, w_0xw^{-1}} V(x), & \text{if } w' = e \\ 0, & \text{otherwise} \end{cases}$$

we see that (ii) \Leftrightarrow (iii).

By theorem 2.9 in [X] we have $h_{w_0, ww_0, w_0x} = h_{w_0, ww_0w^{-1}, w_0xw^{-1}}$. So we have (iii) \Leftrightarrow (iv).

By Lemma 2.4 (i) we see that $x_1 w_{\Gamma} = w w_0$ for some $w \in W$. Using the method in [S] one knows that $w \in \mathfrak{S}$. Thus we have (iv) \Rightarrow (v). Now we show that (v) \Rightarrow (iv). Let $w \in \mathfrak{S}$, then $ww_0 \in \Gamma_0$, hence by 2.10

$$C_{ww_0} = \sum_{\substack{y \in X^+ \\ I \subseteq I_0}} h_{1, y} C_{x_1 w_{\Gamma}} S_y, \quad h_{1, y} \in H'_{q_0}.$$

Since $C_{w_0} h_{1, y} = a_{1, y} C_{w_0}$ for some $a_{1, y} \in \mathbb{C}$, we have

$$\sum_{x \in X^+} h_{w_0, ww_0, w_0x} \text{tr}(s, V(x)) = \sum_{\substack{y \in X^+ \\ I \subseteq I_0}} a_{1, y} \alpha_1 \text{tr}(s, V(y)) = 0.$$

Finally we prove that (v) and (vi) are equivalent.

One direction is obvious. Now assume that (vi) holds. Let $J \subseteq I_0$. We use induction on $l(x_j)$ to prove that $\alpha_j = 0$. When $\eta_{J'} = 0$ or $J \in \Delta_{q_0}$ we have $\alpha_j = 0$ by 2.7 or by (vi). Suppose $\eta_{J'} \neq 0$ and $J \notin \Delta_{q_0}$. Choose $j \in J$ such that $\eta_{J' \cup \{j\}} \neq 0$. Let $K = J - \{j\}$, then $K' = J' \cup \{j\}$. We have

$$\begin{aligned} C_{w_0} C_{x_j w_{J'}} &= \frac{1}{\eta_{K'}} C_{w_0} C_{w_{K'}} C_{x_j w_{J'}} \quad (\text{by 2.6}) \\ &= \frac{\eta_{J'}}{\eta_{K'}} C_{w_0} (C_{w_{K'} x_K x_j} + \sum_{\substack{I \subseteq I_0 \\ y \in X^+}} h_{1, y} C_{x_1 w_{\Gamma}} S_y), \quad h_{1, y} \in H'_{q_0} \quad (\text{by 2.6, 2.10}). \end{aligned}$$

Let $C_{w_0} h_{1, y} = a_{1, y} C_{w_0}$, $a_{1, y} \in \mathbb{C}$. By 2.7 we see that $a_{1, y} \eta_{J'} \neq 0$ implies that $l(x_1 y) < l(x_j)$. Obviously $l(x_K) < l(x_j)$. Using induction hypothesis we get

$$\alpha_j = \frac{\eta_{J'}}{\eta_{K'}} (\alpha_K \text{tr}(s, V(x_j)) + \sum_{\substack{I \subseteq I_0 \\ y \in X^+}} a_{1, y} \alpha_1 \text{tr}(s, V(y))) = 0.$$

The lemma is proved.

LEMMA 3.8. — $(\phi_{q_0})_{*, c_0}(E_s) = 0$ if and only if $C_{w_0} E_{s, q_0} = 0$.

Proof. — The “if” part is obvious. The “only if” part need to do a little more.

Assume that $C_{w_0}E_{s, q_0} \neq 0$. By 3.7 we see that $\alpha_I \neq 0$ for some $I \subseteq I_0$. As in [LX] we define an automorphism $\alpha: W \rightarrow W$ by

$$\alpha(wx) = w_0 wx^{-1} w_0, \quad w \in W_0, \quad x \in X.$$

One verifies that α leaves stable W_0, X, S, S' . In particular, α induces a bijection $\alpha: I_0 \rightarrow I_0$ and an automorphism $\sigma: H_{q_0} \rightarrow H_{q_0}$ by defining $C_u \rightarrow C_{\alpha(u)}$, $u \in W$. Let $J = \alpha(I)$, we have $\alpha(x_I) = x_J$, $\alpha(w_I) = w_J$. Consider

$$\psi_s \phi_{q_0, c_0}(C_{x_J^{-1} w_J}) = (n_{w', w}) \in M_{\mathfrak{S} \times \mathfrak{S}}(\mathbb{C}).$$

By 2.4 and 2.12, we know that $n_{w', w} = 0$ if $w' \neq e$ and

$$n_{e, w} = \sum_{x \in X^+} h_{x_J^{-1} w_J, w w_0 w^{-1}, w_0 x w^{-1}} tr(s, V(x)).$$

In particular,

$$n_{e, e} = \sum_{x \in X^+} h_{x_J^{-1} w_J, w_0, w_0 x} tr(s, V(x)).$$

We claim that $n_{e, e} = \alpha_I$. In fact, let ι be the antiautomorphism of H_{q_0} defined by $C_u \rightarrow C_{u^{-1}}$, $u \in W$. Apply ι to the equality

$$C_{w_0} C_{x_I w_I} = \sum_{x \in X^+} h_{w_0, x_I w_I, w_0 x} C_{w_0 x}.$$

We get

$$C_{x_I^{-1} w_I} C_{w_0} = \sum_{x \in X^+} h_{w_0, x_I w_I, w_0 x} C_{x^{-1} w_0}.$$

Apply σ to the above identity we obtain

$$C_{x_J^{-1} w_J} C_{w_0} = \sum_{x \in X^+} h_{w_0, x_I w_I, w_0 x} C_{w_0 x}.$$

Therefore $h_{x_J^{-1} w_J, w_0, w_0 x} = h_{w_0, w_I w_I, w_0 x}$ and $n_{e, e} = \alpha_I \neq 0$. By this and $n_{w', w} = 0$ if $w' \neq e$ we see that α_I is an eigenvalue of $\psi_s \phi_{q_0, c_0}(C_{x_J^{-1} w_J})$. Let $0 \neq v \in E_{s, q_0}$ be such that $C_{x_J^{-1} w_J} v = \alpha_I v$. Let F be the H_{q_0} -submodule of E_{s, q_0} generated by v . Then F has a maximal H_{q_0} -submodule F_0 which doesn't contain v . F/F_0 is an irreducible H_{q_0} -module. Moreover $C_{x_J^{-1} w_J}(F/F_0) \neq 0$ since $v \notin F_0$. We have proved that $(\phi_{q_0})_{*, c_0}(E_s) \neq 0$.

Theorem 3.6 follows from 3.7 and 3.8.

3.9. There are two special cases. One is that $\eta_{I_0} = 0$ but $\eta_I \neq 0$ for any proper subset I of I_0 . In this case we have $\Delta_{q_0} = \{\{i\} \mid i \in I_0\}$. Let $i' = I_0 - \{i\}$. By 2.7 we have $h_{w_0, x_I w_I, w_0 x} = \eta_{i'} a_{i, x}$ for some $a_{i, x} \in \mathbb{C}$. Moreover, $a_{i, x} \neq 0$ implies that $w_0 x \leq w_0 x_i$ and $a_{i, x_i} = 1$. By this we see that the equation system

$$\alpha_{\{i\}} = \eta_{i'} \sum_{\substack{x \in X^+ \\ w_0 x \leq w_0 x_i}} a_{i, x} tr(s, V(x)) = 0, \quad i \in I_0$$

uniquely determines $tr(s, V(x_i))$, $i \in I_0$. In other words, there exists a unique semisimple conjugacy class s of G such that $\alpha_{\{i\}} = 0$ for all $i \in I_0$. By 3.6 we have got the following.

PROPOSITION. — *There exists a unique semisimple conjugacy class s of G such that $(\phi_{q_0})_{*, c_0}(E_s) = 0$ when $\eta_{I_0} = 0$ but $\eta_I \neq 0$ for any proper subset I of I_0 .*

When W is of type \tilde{A}_n . We can determine the semisimple conjugacy class s in the proposition explicitly. We have $a_{i,x} = 0$ if $x \neq x_i$ since x_i is a minimal dominant weight for any $i \in I_0$. So $\alpha_{\{i\}} = \eta_i \cdot tr(s, V(x_i))$. Let T be the diagonal subgroup of $G = SL_{n+1}(\mathbb{C})$. We may require that $x_i \in \text{Hom}(T, \mathbb{C}^*)$ is defined by $x_i(t) = t_1 t_2 \dots t_i$ where $t = \text{diag}(t_1, t_2, \dots, t_{n+1}) \in T$. Thus, we have

$$tr(s, V(x_i)) = \sum_{\substack{j_a \in I_0 \cup \{n+1\} \\ j_a \neq j_b \text{ if } a \neq b}} t_{j_1} t_{j_2} \dots t_{j_i}$$

where $t = \text{diag}(t_1, t_2, \dots, t_{n+1}) \in s \cap T$, s a semisimple conjugacy class of G . Hence, $tr(s, V(x_i)) = 0$, $1 \leq i \leq n$ is equivalent to that t_i ($1 \leq i \leq n+1$) is the solution of the equation $\lambda^{n+1} + (-1)^{n+1} = 0$. So if $\eta_{I_0} = 0$ but $\eta_I \neq 0$ for any proper subset I of I_0 , $(\phi_{q_0})_{*, c_0}(E_s) = 0$ if and only if the eigenpolynomial of s is $\lambda^{n+1} + (-1)^{n+1}$.

Another special case is that $q_0 + q_0^{-1} = 0$. In this case $\Delta_{q_0} = \{I_0\}$. So $(\phi_{q_0})_{*, c_0}(E_s) = 0$ if and only if $\alpha_{I_0} = 0$. If we identify the set $\{\text{semisimple conjugacy classes of } G\}$ with \mathbb{C}^n through the bijection

$$s \rightarrow (tr(s, V(x_1)), tr(s, V(x_2)), \dots, tr(s, V(x_n))),$$

then $\alpha_{I_0} = 0$ defines a hypersurface in \mathbb{C}^n . That is to say, the set $\{\text{semisimple conjugacy class } s \text{ of } G \mid (\phi_{q_0})_{*, c_0}(E_s) = 0\}$ is a variety of dimension $n-1$.

When W_0 is of rank 2, if $\eta_{I_0} = 0$, then either $\eta_I \neq 0$ for any proper subset $I \subseteq I_0$ or $q_0 + q_0^{-1} = 0$. The above discussion shows that $(\phi_{q_0})_{*, c_0}$ is an isomorphism if and only if $\eta_{I_0} \neq 0$.

3.10. In general it is difficult to compute $C_{w_0} C_{x_1 w_1}$ in H . Now we compute it for the simplest case: x_1 is the highest short root.

When $x_1 \in X^+$ is the highest short root, $x_1 w_1 = r_0 w_0$, and $w_0 x \leq w_0 x_1$, $x \in X^+$ implies that $x = e$ or x_1 . So by 2.7, in H we have

$$C_{w_0} C_{r_0 w_0} = C_{w_0} C_{x_1 w_1} = \sigma_{I'}(C_{w_0 x_1} + a C_{w_0}),$$

where $\sigma_{I'} \in A = \mathbb{C}[q, q^{-1}]$ is determined by $C_{w_1} C_{w_1} = \sigma_{I'} C_{w_1}$, $a \in A$. We need to determine the coefficient a . Comparing the coefficient of T_e in both sides we get

$$q^{-l(w_0)-1} \sigma_{I_0} = q^{-l(w_0 w_1)} \sigma_{I'} P_{e, w_0 x_1}(q^2) + a q^{-l(w_0)} \sigma_{I'}.$$

i. e.

$$(a) \quad \sigma_{I_0} = q^{1-l(x_1)} \sigma_{I'} P_{w_0, w_0 w_1}(q^2) + a q \sigma_{I'}.$$

Using the formula 8.10 in [L2] we get the following

PROPOSITION 3.11. — *If x_1 is the highest short weight, then*

$$P_{w_0, w_0 x_1} = \begin{cases} \sum_{i=1}^n q^{e_i-1} & \text{for type } \tilde{A}_n, \tilde{D}_n, \tilde{E}_n. \\ 1 & \text{for type } \tilde{C}_n, \tilde{G}_2. \\ \frac{q^{2(n-1)} - 1}{q^2 - 1} & \text{for type } \tilde{B}_n. \\ q^4 + 1 & \text{for type } \tilde{F}_4. \end{cases}$$

where e_1, \dots, e_n are the exponents of W_0 .

By the proposition and 3.10(a) we obtain the following

PROPOSITION 3.12. — *In H we have*

$$C_{w_0} C_{r_0 w_0} = C_{w_0} C_{x_1 w_1} = \sigma_{1'} C_{w_0 x_1} + \frac{\sigma_{1_0}}{[e_n + 1]} [e_n] C_{w_0},$$

where e_n is the largest exponent of W_0 and $[i] = (q^i - q^{-i}) / (q - q^{-1})$ for any $i \in \mathbb{N}$.

3.13. When W is of type \tilde{A}_n , the highest short weight is $x_1 x_n$.

$$\eta_{1_0} = [2]_{q_0} [3]_{q_0} \dots [n+1]_{q_0},$$

where $[i]_{q_0}$ is the specialization at $q_0 \in \mathbb{C}^*$ of $[i]$. By 3.12, in H_{q_0} we have

$$C_{w_0} C_{r_0 w_0} = [2]_{q_0} [3]_{q_0} \dots [n-1]_{q_0} (C_{w_0 x_1 x_n} + [n]_{q_0}^2 C_{w_0}).$$

Now suppose $[n]_{q_0} = 0$ but $[i]_{q_0} \neq 0$ for $i, 1 \leq i \leq n-1$, then $\Delta_{q_0} = \{\{1, n\}, \{2\}, \{3\}, \dots, \{n-1\}\}$. By 3.9 and 3.12 we see that $\alpha_1 = 0, I \in \Delta_{q_0}$ is equivalent to $\text{tr}(s, V(x_1 x_n)) = 0, \text{tr}(s, V(x_i)) = 0, 2 \leq i \leq n-1$. Note that $\text{tr}(s, V(x_1 x_n)) = \text{tr}(s, V(x_1)) \text{tr}(s, V(x_n)) - 1$, by 3.9, we know that $\alpha_1 = 0, I \in \Delta_{q_0}$ if and only if the eigenpolynomial of s has the form $\lambda^{n+1} - a\lambda^n + (-1)^n a^{-1} \lambda + (-1)^{n+1}, a \in \mathbb{C}^*$. In other words, if $[n]_{q_0} = 0, [i]_{q_0} \neq 0, 1 \leq i \leq n-1$, then $(\phi_{q_0})_{*, c_0}(E_s) = 0$ if and only if the eigenpolynomial of s has the form $\lambda^{n+1} - a\lambda^n + (-1)^n a^{-1} \lambda + (-1)^{n+1}, a \in \mathbb{C}^*$.

4. Examples

4.1. Type \tilde{A}_1 . In this case $G = \text{SL}_2(\mathbb{C}), S = \{r_0, r_1\}, x_1 = r_0 \omega, \Omega = \{e, \omega\}, \eta_{1_0} = q_0 + q_0^{-1}, c_0 = \{w \in W \mid l(w) > 0\}$. Another two-sided cell c of W is Ω .

J_c has two irreducible modules F_0, F_1 . Both have dimension 1 and t_ω acts on F_i by scalar $(-1)^i, i=0, 1$. Via, $\phi_{q_0, c}: H_{q_0} \rightarrow J_c, F_i$ becomes H_{q_0} -module F_{i, q_0} . T_ω acts on F_{i, q_0} by scalar $(-1)^i$ and T_{r_i} acts on F_{i, q_0} by scalar -1 . $(\phi_{q_0})_{*, c}$ is an isomorphism for any $q_0 \in \mathbb{C}^*$.

For c_0 , we have $J_{c_0} = M_{2 \times 2}(\mathbb{R}_G)$ and

$$\begin{aligned}\phi_{q_0, c_0}(C_{r_1}) &= \begin{pmatrix} \eta_{1_0} & V(x_1) \\ 0 & 0 \end{pmatrix} \\ \phi_{q_0, c_0}(C_{r_0}) &= \begin{pmatrix} 0 & 0 \\ V(x_1) & \eta_{1_0} \end{pmatrix} \\ \phi_{q_0, c_0}(C_\omega) &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.\end{aligned}$$

Suppose that $\eta_{1_0} \neq 0$. Let s be the semisimple conjugacy class of G containing $\begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \in G$, then E_{s, q_0} is irreducible if and only if $\eta_{1_0} \neq \pm(t+t^{-1})$. When $\eta_{1_0} = \pm(t+t^{-1})$, $E_{s, q_0}/F_{i, q_0} \simeq M_{s, q_0}$, where $i=0$ if $\eta_{1_0} = -(t+t^{-1})$ and $i=1$ if $\eta_{1_0} = t+t^{-1}$. T_ω acts on M_{s, q_0} by scalar $(-1)^{i-1}$ and T_{r_i} acts on M_{s, q_0} by scalar q_0^2 . $(\phi_{q_0})_{*, c}(E_s) = E_{s, q_0}$ if $\eta_{1_0} \neq \pm(t+t^{-1})$, $(\phi_{q_0})_{*, c_0}(E_s) = M_{s, q_0}$ if $\eta_{1_0} = \pm(t+t^{-1})$. In particular, when $\eta_{1_0} \neq 0$, $(\phi_{q_0})_*$ is an isomorphism.

When $\eta_{1_0} = 0$, one verifies that E_{s, q_0} is irreducible if $t+t^{-1} \neq 0$ and $E_{s, q_0} = F_{0, q_0} \oplus F_{1, q_0}$ if $t+t^{-1} = 0$. In particular $\text{rank ker } (\phi_{q_0})_* = 1$.

4.2. Type \tilde{A}_2 . In this case we have $G = \text{SL}_3(\mathbb{C})$, $S = \{r_0, r_1, r_2\}$, $\Omega = \{1, \omega, \omega^2\}$ and $\omega r_0 = r_1 \omega$, $\omega r_1 = r_2 \omega$, $\omega r_2 = r_0 \omega$, $x_1 = r_0 r_2 \omega$, $x_2 = r_0 r_1 \omega^2$. W has three two-sided cells: $c = \Omega$, c_0 , $c' = W - c \cup c_0$. c' is the two-sided cell of W containing r_0, r_1, r_2 .

It is obviously $(\phi_{q_0})_{*, c}$ is an isomorphism.

Now consider $J_{c'}$. Any element in c' has one of the following forms: $\omega^i r_1 x_1^a \omega^j$, $\omega^{i+1} x_1^a \omega^j$, $\omega^{i+2} r_2 x_2^a \omega^{j+1}$, $\omega^{i+1} x_2^a \omega^{j+1}$, $i, j = 0, 1, 2$. We define a \mathbb{C} -linear map $\theta: J_{c'} \rightarrow M_{3 \times 3}(A)$, $A = \mathbb{C}[q, q^{-1}]$, by $\theta(t_w) = (\mathcal{M}_{ab}) \in M_{3 \times 3}(A)$, $w \in c'$. Assume that w is of one of the above forms, then $m_{ab} = 0$ except $(a, b) = (i+1, j+1)$ and

$$m_{i+1, j+1} = \begin{cases} q^{2a} & \text{if } w = \omega^i r_1 x_1^a \omega^j \\ q^{2a-1} & \text{if } w = \omega^{i+1} x_1^a \omega^j \\ q^{-2a} & \text{if } w = \omega^{i+2} r_2 x_2^a \omega^{j+1} \\ q^{-2a+1} & \text{if } w = \omega^{i+1} x_2^a \omega^{j+1}. \end{cases}$$

By [L 1, 3.8] we know that θ is a \mathbb{C} -algebra isomorphism. We have

$$\begin{aligned}\theta \phi_{q_0, c'}(C_{r_1}) &= \begin{pmatrix} [2]_{q_0} & q^{-1} & q \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ \theta \phi_{q_0, c'}(C_\omega) &= \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.\end{aligned}$$

Specialize q to $a \in \mathbb{C}^*$, we get a simple representation ψ_a of $J_{c'} = M_{3 \times 3}(A)$ and any simple representation of $J_{c'}$ is isomorphic to some ψ_a , $a \in \mathbb{C}^*$. Let E_a be a simple $J_{c'}$ -module providing ψ_a .

