

BULLETIN DE LA S. M. F.

GOPAL PRASAD

Non-vanishing on the first cohomology

Bulletin de la S. M. F., tome 105 (1977), p. 415-418

http://www.numdam.org/item?id=BSMF_1977__105__415_0

© Bulletin de la S. M. F., 1977, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (<http://smf.emath.fr/Publications/Bulletin/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

NON-VANISHING
OF THE FIRST COHOMOLOGY

BY

GOPAL PRASAD
[Tata Institute, Bombay]

RÉSUMÉ. — On démontre que, pour les réseaux Γ du type fini dans les groupes semi simples sur les corps locaux de caractéristique positive, $H^1(\Gamma, \text{Ad})$ ne s'annule pas; ceci est bien différent de ce que passe dans le cas de caractéristique zéro.

ABSTRACT. — It is shown here that, for any finitely generated lattice Γ in certain semi simple groups over local fields of positive characteristics, $H^1(\Gamma, \text{Ad})$ is non-vanishing; this is in sharp contrast with the situation in characteristic zero.

Let K be a local field (i. e. a non-discrete locally compact field), and let \mathbf{G} be a connected semi simple algebraic group defined over K . Let $G = \mathbf{G}(K)$, and let $r = K\text{-rank } \mathbf{G}$. The topology on K induces a locally compact Hausdorff topology on G ; in the sequel, we assume G endowed with this topology. G is then a K -analytic group. Let Γ be a lattice in G i.e., a discrete subgroup of G such that G/Γ carries a finite G -invariant Borel measure. We assume that Γ is *irreducible*, i.e. no subgroup of Γ of finite index is a direct product of two infinite normal subgroups.

In case $K = \mathbf{R}$ and G is not locally isomorphic to either $SL(2, \mathbf{R})$ or $SL(2, \mathbf{C})$, it is known that $H^1(\Gamma, \text{Ad}) = 0$; where, as usual, Ad denotes the adjoint representation of G on its Lie algebra (see WEIL [9], [10] for uniform lattices; for non-uniform lattices in groups of \mathbf{R} -rank > 1 , this vanishing theorem follows from the results of RAGHUNATHAN [8], combined with the results of MARGULIS [4] on arithmeticity; for non-uniform lattices in groups of \mathbf{R} -rank 1, it is contained in GARLAND-RAGHUNATHAN [2]).

It is also known, in view of a recent result of MARGULIS ([5], theorem 8), that in case K is non-archimedean but of characteristic zero, $H^1(\Gamma, \text{Ad}) = 0$ when $r > 1$.

The object of this note is to show that *when K is of positive characteristic, then it is not in general true that $H^1(\Gamma, \text{Ad}) = 0$.*

We shall in fact prove the following theorem.

THEOREM. — *Let F be a finite field, and let K be the local field $F((t))$. Let \mathbf{G} be a connected semi simple algebraic group, with trivial center, defined over F . Let $G = \mathbf{G}(K)$, let Γ be a finitely generated lattice in G . Then $H^1(\Gamma, \text{Ad}) \neq 0$.*

Remark. — If \mathbf{G} has no K -rank 1 factors, then according to a well-known theorem of D.A. KAZHDAN (see [1]), every lattice in G is finitely generated.

For the proof of the theorem, we need to recall a result of WEIL [10].

We introduce some notation and a definition.

Let Λ be a finitely generated abstract group. We shall let $\mathcal{A}(\Lambda, G)$ denote the space of all homomorphisms of Λ in G with the topology of pointwise convergence. There is a natural action of G on $\mathcal{A}(\Lambda, G)$ induced by the inner automorphism.

Now assume that Λ is a finitely generated subgroup of G , and let $\iota : \Lambda \rightarrow G$ be the natural inclusion. Then Λ is said to be *locally* (or *infinitesimally*) *rigid* if the orbit of ι under G is open in $\mathcal{A}(\Lambda, G)$. According to a result of WEIL [10], vanishing of $H^1(\Lambda, \text{Ad})$ implies local rigidity of Λ .

Proof of the theorem. — In view of the above result of WEIL, to prove that $H^1(\Gamma, \text{Ad}) \neq 0$, it suffices to show that Γ is not infinitesimally rigid.

For $i > 1$, $t \mapsto t + t^i$ extends uniquely to give a continuous automorphism a_i of $F((t))/F$. It is evident that, for any fixed $x \in F((t))$, the sequence $\{a_i(x)\}$ converge to x .

Now since \mathbf{G} is defined over F , a_i induces a continuous automorphism α_i of G . Therefore, for all i , $\alpha_i \cdot \iota$ is an embedding of Γ in G ; where $\iota : \Gamma \rightarrow G$ is the natural inclusion of Γ in G . It is also obvious that the sequence $\{\alpha_i \cdot \iota\}$ converges to ι in $\mathcal{A}(\Gamma, G)$. We shall show that none of the $\alpha_i \cdot \iota$ lie in the G -orbit of ι . This will prove that Γ is not locally rigid and hence $H^1(\Gamma, \text{Ad}) \neq 0$.

If possible, assume that, for some i , $\alpha_i \cdot \iota = \text{Int } g_i \cdot \iota$. Then $(\text{Int } g_i^{-1} \cdot \alpha_i) \cdot \iota = \iota$, and the main theorem of PRASAD [6] implies that $\text{Int } g_i^{-1} \cdot \alpha_i$ is the identity automorphism of G . Hence, $\alpha_i = \text{Int } g_i$.

We now fix a 1-dimensional torus $\mathbf{T} (\subset \mathbf{G})$ which is defined and split over the finite field F (existence of such a torus follows from Lang's theorem [3]). Let $T = \mathbf{T}(K)$. Then since \mathbf{T} is defined over F , $\alpha_i(T) = T$. Moreover,

for any rational character χ on \mathbf{T} and all $t \in T$,

$$\chi(\alpha_i(t)) = a_i(\chi(t)).$$

Since $\alpha_i = \text{Int } g_i$ and $\alpha_i(T) = T$, it follows that g_i normalizes T and hence also \mathbf{T} . Therefore, for any rational character χ on \mathbf{T} :

$$\chi(\alpha_i(t)) = \chi(g_i t g_i^{-1}) = \chi^d(t),$$

where $d = +1$ or -1 . Hence,

$$(\star) \quad a_i(\chi(t)) = \chi^d(t), \quad \text{where } d = +1 \quad \text{or} \quad -1.$$

Now take χ to be one of the generators of the group of rational characters on \mathbf{T} . Then it follows from (\star) that, for all $k \in K$, either

$$a_i(k) = k \quad \text{or} \quad a_i(k) = k^{-1}.$$

But it is obvious from the definition of a_i , that this is not the case. Hence, none of the $\alpha_i \cdot \mathfrak{v}$ lie in the G -orbit of \mathfrak{v} . This proves that $H^1(\Gamma, \text{Ad}) \neq 0$.

Remark. — As the above proof shows, Γ is not locally rigid. However, in case K -rank $\mathbf{G} > 1$ and Γ is an irreducible uniform lattice, it is *strongly rigid* (see PRASAD [7], § 8).

REFERENCES

- [1] DELAROCHE (C.) and KIRILLOV (A.). — Sur les relations entre l'espace dual d'un groupe et la structure de ses sous-groupes fermés, *Séminaire Bourbaki*, 20^e année, 1967/1968, n^o 343, 22 p.
- [2] GARLAND (H.) and RAGHUNATHAN (M. S.). — Fundamental domains for lattices in \mathbf{R} -rank 1 semi simple groups, *Annals of Math*, Series 2, t. 92, 1970, p. 279-326.
- [3] LANG (S.). — Algebraic groups over finite fields, *Amer. J. of Math.*, t. 78, 1956, p. 555-563.
- [4] MARGULIS (G. A.). — Arithmetic properties of discrete subgroups [in Russian], *Uspekhi Mat. Nauk*, t. 29, 1974, p. 49-98.
- [5] MARGULIS (G. A.). — Discrete groups of motions of manifolds of non-positive curvature [in Russian], "Proceedings of the International Congress of Mathematicians [1974, Vancouver]", Vol. 2, p. 21-34. — Vancouver, Canadian mathematical Congress, 1975.
- [6] PRASAD (G.). — Triviality of certain automorphisms of semi-simple groups over local fields, *Math. Annalen*, t. 218, 1975, p. 219-227.
- [7] PRASAD (G.). — Lattices in semi simple groups over local fields, *Advances in Math.* (to appear).
- [8] RAGHUNATHAN (M. S.). — Cohomology of arithmetic subgroups of algebraic groups I and II, *Annals of Math.*, t. 86, 1967, p. 409-424, and t. 87, 1968, p. 279-304.

- [9] WEIL (A.). — Discrete subgroups of Lie groups, I and II, *Annals of Math.*, t. 72, 1960, p. 369-384, and t. 75, 1962, p. 578-602.
- [10] WEIL (A.). — Remarks on the cohomology of groups, *Annals of Math.*, t. 80, 1964, p. 149-157.

(Texte reçu le 18 février 1977)

Gopal PRASAD,
School of Mathematics,
Tata Institute of Fundamental Research
Homi Bhabha Road,
Bombay 400 005,
Inde.
