

# BULLETIN DE LA S. M. F.

CHRISTIAN BERG

JESPER LAUB

## **The resolvent for a convolution kernel satisfying the domination principle**

*Bulletin de la S. M. F.*, tome 107 (1979), p. 373-384

[http://www.numdam.org/item?id=BSMF\\_1979\\_\\_107\\_\\_373\\_0](http://www.numdam.org/item?id=BSMF_1979__107__373_0)

© Bulletin de la S. M. F., 1979, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (<http://smf.emath.fr/Publications/Bulletin/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

**THE RESOLVENT FOR A CONVOLUTION KERNEL  
SATISFYING THE DOMINATION PRINCIPLE**

BY

CHRISTIAN BERG and JESPER LAUB (\*)  
[Københavns Universitet]

---

**ABSTRACT.** — Let  $N$  be a convolution kernel on a locally compact abelian group. It is shown that if  $N$  satisfies the domination principle and is non-singular, then there exists a splitting  $N = N_0 + N'$  of  $N$  in which  $N_0$  is a resolvent kernel and  $N'$  is  $N$ -invariant. Furthermore, the singular part  $N'$  of  $N$  is either  $N_0$ -invariant or a  $N_0$ -potential of a  $N$ -invariant measure. These results simplify Theorems of M. Irô.

**RÉSUMÉ.** — Soit  $N$  un noyau de convolution dans un groupe abélien localement compact. Pour  $N$  satisfaisant au principe de domination et étant non singulier, on démontre qu'il existe une partition  $N = N_0 + N'$  de  $N$ , où  $N_0$  est un noyau à résolvante et  $N'$  est  $N$ -invariante. De plus, la partie singulière  $N'$  de  $N$  est ou bien  $N_0$ -invariante ou bien un  $N_0$ -potentiel d'une mesure  $N$ -invariante. Ces résultats simplifient des théorèmes de M. Irô.

**Introduction**

Let  $G$  be a locally compact abelian group and  $N$  a convolution kernel on  $G$  satisfying the domination principle. In [2], Irô introduced a family  $(N_p)_{p>0}$  of convolution kernels, which in later papers ([3], [5]) turned out to be the resolvent family for the regular part  $N_0$  of  $N$ . Some of the proofs in these papers are complicated, so it is of interest to give a simple and unified treatment of the resolvent and the regular part of  $N$  based entirely on the Riesz decomposition theorem, and this is the aim of the present paper.

A complete proof of the Riesz decomposition theorem was given in [7], which will be a prerequisite for the present paper. Less general versions of the Riesz decomposition Theorem appeared in [3] and [4], and the treatment in [3] assumes knowledge of the resolvent and the regular part.

---

(\*) Texte reçu le 16 octobre 1978.

Christian BERG and Jesper LAUB, Københavns Universitets Matematiske Institut, Universitetsparken 5, DK-2100 København Ø (Danemark).

The idea behind our treatment is as follows:

For each  $p > 0$ , we have  $pN + \varepsilon_0 < N$ , so let  $N = (pN + \varepsilon_0) \star N_p + \eta_p$  be the Riesz decomposition of  $N$  with respect to  $pN + \varepsilon_0$  as sum of a  $(pN + \varepsilon_0)$ -potential, generated by a measure  $N_p$ , and a  $(pN + \varepsilon_0)$ -invariant measure  $\eta_p$ . The measures  $N_p$  and  $\eta_p$  are uniquely determined, and this leads to the resolvent equation for  $(N_p)_{p>0}$ . For  $p$  tending to zero,  $N_p$  increases to the regular part of  $N$ .

### Preliminaries

In the following,  $G$  denotes an arbitrary locally compact abelian group, and  $N$  a convolution kernel on  $G$  satisfying the domination principle.

A positive measure  $\xi$  on  $G$  is called  $N$ -excessive, if  $N$  satisfies the relative domination principle with respect to  $\xi$  ( $N < \xi$ ) (cf. [4], [7]). The set of  $N$ -excessive measures is a vaguely closed convex cone  $E(N)$ , which is infimum-stable, and every  $\xi \in E(N)$  is the vague limit of an increasing net of  $N$ -potentials. For an open subset  $\Omega \subseteq G$  and a measure  $\xi \in E(N)$ , the reduced measure  ${}_N R_\xi^\Omega$  of  $\xi$  over  $\Omega$  (with respect to  $N$ ) is defined (cf. [7]) as

$${}_N R_\xi^\Omega = \inf \{ \tau \in E(N); \tau \geq \xi \text{ in } \Omega \}.$$

We write  $R_\xi^\Omega$  instead of  ${}_N R_\xi^\Omega$  when  $N$  is clear from the context.

Let  $\mathcal{V}$  denote the set of compact neighbourhoods of 0 in  $G$ . A measure  $\xi \in E(N)$  is called  $N$ -invariant if  $R_\xi^{V'} = \xi$  for all  $V \in \mathcal{V}$ . The set of  $N$ -invariant measures is a convex cone  $I(N)$ , closed under increasing limits.

*Definition.* — The singular part  $N'$  of  $N$  is the limit  $N' = \lim_{V \uparrow G} R_N^{V'}$  of the decreasing net  $(R_N^{V'})_{V \in \mathcal{V}}$ , when  $V \in \mathcal{V}$  increases to  $G$ .

The regular part  $N_0$  of  $N$  is  $N_0 = N - N'$ . Note that  $N_0 \geq 0$ .

The convolution kernel  $N$  is called *singular* (resp. *non-singular*) if  $N_0 = 0$  (resp.  $N_0 \neq 0$ ).

The following Riesz decomposition theorem will be essential later (cf. [7]).

**PROPOSITION 1.** — *Suppose  $N$  is non-singular. Every  $\xi \in E(N)$  has a decomposition*

$$\xi = N \star \mu + \eta, \quad \text{where } \eta \in I(N).$$

*The invariant part  $\eta$  is uniquely determined, and the measure  $\mu$  is uniquely determined if (and only if)  $N$  satisfies the principle of unicity of mass.*

We shall use some alternative characterizations of  $N$ -excessive and  $N$ -invariant measures.

LEMMA 2 (ITÔ [4], LAUB [7]). — Suppose  $N$  is non-singular. A positive measure  $\xi$  is  $N$ -invariant if (and only if) there exists a net  $(\lambda_\alpha)_{\alpha \in A}$  of positive measures with compact support such that

$$N \star \lambda_\alpha \uparrow \xi, \quad \lambda_\alpha \rightarrow 0.$$

In Corollary 6 below the conclusion of Lemma 2 is shown to be valid also for singular kernels.

The following result is well-known and not difficult to establish.

LEMMA 3. — Let  $N = 1/a \sum_{n=0}^{\infty} \sigma^n$  be an elementary kernel ( $a > 0$ ). Then

- (i)  $\xi \in E(N) \Leftrightarrow \sigma \star \xi \leq \xi,$
- (ii)  $\eta \in I(N) \Leftrightarrow \sigma \star \eta = \eta.$

For  $c > 0$ , the convolution kernel  $N + c \varepsilon_0$  satisfies the domination principle and the principle of unicity of mass, where  $\varepsilon_0$  denotes the Dirac measure at 0. Moreover, it is easily seen that if  $N < \xi$  then  $N + c \varepsilon_0 < \xi$ , i. e.  $E(N) \subseteq E(N + c \varepsilon_0)$ . For  $V \in \mathcal{V}$ , we consequently have

$${}_N R_N^{cV} \geq_{(N+c\varepsilon_0)} R_{N+c\varepsilon_0}^{cV},$$

so that  $N + c \varepsilon_0$  is non-singular.

The following Lemma is an extension of [4] (Corollaire 1, p. 340); the hypothesis of  $N$  being non-singular is removed.

LEMMA 4. — For  $c > 0$ , we have

$$I(N) = I(N + c \varepsilon_0).$$

*Proof.* — Suppose first that  $\eta \in I(N + c \varepsilon_0)$ . By Lemma 2, there exists a net  $(\lambda_\alpha)_{\alpha \in A}$  of positive measures such that  $(N + c \varepsilon_0) \star \lambda_\alpha \uparrow \eta$  and  $\lambda_\alpha \rightarrow 0$ . Therefore, we have  $\eta = \lim N \star \lambda_\alpha$  and then  $\eta \in E(N)$ . For  $V \in \mathcal{V}$ , we find

$$\eta =_{(N+c\varepsilon_0)} R_\eta^{cV} \leq_N R_\eta^{cV} \leq \eta.$$

hence  $\eta \in I(N)$ .

Suppose next that  $\eta \in I(N)$ . Then  $\eta \in E(N) \subseteq E(N + c \varepsilon_0)$ , and since  $N + c \varepsilon_0$  is non-singular,  $\eta$  has a Riesz decomposition

$$\eta = (N + c \varepsilon_0) \star \nu_c + \eta_c,$$

where  $\eta_c \in I(N + c \varepsilon_0) \subseteq I(N)$ . We shall prove that  $\nu_c = 0$ .

Let  $V \in \mathcal{V}$ , and choose a net of positive measures  $(\mu_\alpha)_{\alpha \in A}$  with compact support in  $\overline{\mathbb{C}}V$  such that  $N \star \mu_\alpha \uparrow R_N^{\mathbb{C}V}$ . Since  $N < \eta$  the net  $(\eta \star \mu_\alpha)_{\alpha \in A}$  is increasing and

$$\lim_A \eta \star \mu_\alpha \leq \eta.$$

Now choose  $(\lambda_\beta)_{\beta \in B}$  such that  $N \star \lambda_\beta \uparrow \eta$ , and if  $N$  is non-singular with the additional property that  $\lambda_\beta \rightarrow 0$ . We then claim that

$$\lim_B (N - R_N^{\mathbb{C}V}) \star \lambda_\beta = 0.$$

This is true, because if  $N$  is singular then  $N = R_N^{\mathbb{C}V}$ , and if  $N$  is non-singular then  $N - R_N^{\mathbb{C}V}$  has compact support and  $\lambda_\beta \rightarrow 0$ . We then have

$$\begin{aligned} \eta &= \lim_B N \star \lambda_\beta = \lim_B R_N^{\mathbb{C}V} \star \lambda_\beta \\ &= \lim_B (\lim_A N \star \mu_\alpha \star \lambda_\beta) \leq \lim_A \eta \star \mu_\alpha, \end{aligned}$$

hence  $\eta = \lim_A \eta \star \mu_\alpha$ .

Since  $\eta_c \in I(N)$ , we similarly find  $\lim_A \eta_c \star \mu_\alpha = \eta_c$ . If  $\mu_{\mathbb{C}V}$  denotes a vague accumulation point of  $(\mu_\alpha)_{\alpha \in A}$ , we may assume that  $\mu_\alpha \rightarrow \mu_{\mathbb{C}V}$ , and since  $N \star \mu_\alpha \leq N$  and  $N \star v_c$  exists, Deny's convergence Lemma ([1], Lemma 5.2) shows that

$$\lim_A \mu_\alpha \star v_c = \mu_{\mathbb{C}V} \star v_c.$$

If we convolve all terms in the Riesz decomposition of  $\eta$  with  $\mu_\alpha$  and go to the limit, we obtain

$$\eta = R_N^{\mathbb{C}V} \star v_c + c \mu_{\mathbb{C}V} \star v_c + \eta_c.$$

Finally, letting  $V$  increase to  $G$ , Deny's convergence Lemma shows that  $\mu_{\mathbb{C}V} \star v_c \rightarrow 0$  because  $\text{supp}(\mu_{\mathbb{C}V}) \subseteq \overline{\mathbb{C}V}$ , and hence

$$\eta = N' \star v_c + \eta_c,$$

which compared to the original decomposition gives  $v_c = 0$ , so we have

$$\eta = \eta_c \in I(N + c\varepsilon_0).$$

As an application of Lemma 4, we prove the following result which will not be used in the sequel, but it might be of independent interest.

**PROPOSITION 5.** — *The following conditions about  $N$  are equivalent:*

- (i)  $N$  is singular.
- (ii)  $I(N) = E(N)$ .
- (iii) *There exists a net  $(\lambda_\alpha)_{\alpha \in A}$  of positive measures with compact support such that  $N \star \lambda_\alpha \uparrow N$  and  $\lambda_\alpha \rightarrow 0$ .*

*Proof.*

(i)  $\Rightarrow$  (ii): Let  $\mu$  be a positive measure such that  $N \star \mu$  exists. By Lemma 1.8 in [7], the net  $R_N^{\mathfrak{C}V} \star \mu$  decreases to  $N \star \mu$  as  $V$  increases to  $G$ , hence  $R_N^{\mathfrak{C}V} \star \mu = N \star \mu$  for all  $V \in \mathcal{V}$  so that  $N \star \mu \in I(N)$ . Since every measure  $\xi \in E(N)$  is an increasing limit of potentials  $N \star \mu$ , we get  $E(N) \subseteq I(N)$ .

(ii)  $\Rightarrow$  (iii): By Lemma 4, we have  $N \in I(N + \varepsilon_0)$ , so by Lemma 2 there exists a net  $(\lambda_\alpha)_{\alpha \in A}$  such that  $(N + \varepsilon_0) \star \lambda_\alpha \uparrow N$  and  $\lambda_\alpha \rightarrow 0$ . Therefore,  $N \star \lambda_\alpha \rightarrow N$ , and  $(N \star \lambda_\alpha)_{\alpha \in A}$  is increasing because  $N + \varepsilon_0 < N$ .

(iii)  $\Rightarrow$  (i): Let  $V \in \mathcal{V}$ , and suppose that  $N \star \lambda_\alpha \uparrow N$  and  $\lambda_\alpha \rightarrow 0$ . Writing  $\lambda_\alpha$  as sum of its restrictions  $\lambda_\alpha|_W$  and  $\lambda_\alpha|\mathfrak{C}W$  to  $W$  and  $\mathfrak{C}W$ , where  $W \in \mathcal{V}$  is a compact neighbourhood of  $V$ , we have  $N \star (\lambda_\alpha|\mathfrak{C}W) \rightarrow N$ . By the domination principle for measures, we find  $R_N^{\mathfrak{C}V} \geq N \star (\lambda_\alpha|\mathfrak{C}W)$ , so taking limits for  $\alpha \in A$  we get  $N \leq R_N^{\mathfrak{C}V}$ , which proves (i).

**COROLLARY 6.** — *The conclusion of Lemma 2 is valid also for singular convolution kernels satisfying the domination principle.*

*Proof.* — In order to show that a  $N$ -invariant measure  $\xi$  is the limit of an increasing net  $(N \star \lambda_\alpha)$  for which  $\lambda_\alpha \rightarrow 0$  one proceeds like in (ii)  $\Rightarrow$  (iii) above. In order to prove the converse one proceeds like in (iii)  $\Rightarrow$  (i).

A family  $(N_p)_{p>0}$  of convolution kernels is called a *resolvent* if

$$N_p = N_q + (q - p) N_p \star N_q \quad \text{for } p, q > 0.$$

A convolution kernel  $N$  is called a *resolvent kernel* if there exists a resolvent  $(N_p)_{p>0}$  such that  $N = \lim_{p \rightarrow 0} N_p$ .

A resolvent kernel  $N$  satisfies the domination principle, and for every  $V \in \mathcal{V}$  there exists a balayaged measure  $\varepsilon'_{\mathfrak{C}V}$  of  $\varepsilon_0$  on  $\mathfrak{C}V$  with respect to  $N$  such that  $R_N^{\mathfrak{C}V} = N \star \varepsilon'_{\mathfrak{C}V}$  (cf. [4], § 3). Since  $\text{supp } \varepsilon'_{\mathfrak{C}V} \subseteq \overline{\mathfrak{C}V}$ , we have  $\lim_{V \uparrow G} \varepsilon'_{\mathfrak{C}V} = 0$ , and therefore  $N' = \lim_{V \uparrow G} R_N^{\mathfrak{C}V} = 0$  because of the dominated convergence property of a resolvent kernel (cf. [4] or [6]).

Suppose now that  $N$  is a non-zero resolvent kernel. KISHI showed in [6] that  $\lim_{p \rightarrow \infty} p N_p$  exists and is the normalized Haar measure  $\omega_K$  of a compact subgroup  $K$  of  $G$ . The group  $K$  is the periodicity group for  $N$ , i. e.  $K = \{x \in G; N \star \varepsilon_x = N\}$ .

If  $\mu$  is a positive measure such that  $N \star \mu = N$ , it follows that  $N_p \star \mu = N_p$  for all  $p$ , hence by the convergence Lemma of DENY that  $\mu \star \omega_K = \omega_K$ .

This shows that  $\mu$  is a probability measure supported by  $K$ . In particular, every pseudo-period of  $N$  (i. e. a point  $x \in G$  such that  $N \star \varepsilon_x$  is proportional to  $N$ ) is a period for  $N$ .

Denoting by  $\mathcal{V}_K$  the set of compact neighbourhoods of  $K$  we consequently have  $N \star \varepsilon'_V \neq N$  for any  $V \in \mathcal{V}_K$ . This implies that the series  $\sum_{n=0}^{\infty} (\varepsilon'_V)^n$  converges and the following formula holds

$$N = (N - N \star \varepsilon'_V) \star \sum_{n=0}^{\infty} (\varepsilon'_V)^n, \quad V \in \mathcal{V}_K.$$

Using this notation, the sets  $E(N)$  and  $I(N)$  can be characterized in the following way:

**PROPOSITION 7.** — *Let  $N$  be a non-zero resolvent kernel with resolvent  $(N_p)_{p>0}$ . Then*

- (i)  $\xi \in E(N) \Leftrightarrow \forall p > 0 : p N_p \star \xi \leq \xi$ .
- (ii)  $\eta \in I(N) \Leftrightarrow \forall p > 0 (\exists p > 0) : p N_p \star \eta = \eta$ .
- (iii)  $\xi \in E(N) \Leftrightarrow \forall V \in \mathcal{V}_K : \varepsilon'_V \star \xi \leq \xi$  and  $\omega_K \star \xi = \xi$ .
- (iv)  $\eta \in I(N) \Leftrightarrow \forall V \in \mathcal{V}_K : \varepsilon'_V \star \eta = \eta$  and  $\omega_K \star \eta = \eta$ .

The invariant part of  $\xi \in E(N)$  is given as  $\lim_{p \rightarrow 0} p N_p \star \xi$ .

*Proof:*

(ii): If  $N$  is a resolvent kernel then  $N + 1/p \varepsilon_0 = 1/p \sum_{n=0}^{\infty} (p N_p)^n$  is an elementary kernel for every  $p > 0$  and hence Lemma 3 and 4 show that

$$\eta \in I(N) \Leftrightarrow \eta \in I\left(N + \frac{1}{p} \varepsilon_0\right) \Leftrightarrow p N_p \star \eta = \eta.$$

“(i)  $\Rightarrow$  ”: For  $\xi \in E(N)$ , Proposition 1 shows that  $\xi = N \star \mu + \eta$ , where  $\eta \in I(N)$ , and from this Riesz decomposition we obtain

$$p N_p \star \xi = p N_p \star N \star \mu + \eta \leq \xi.$$

“(i)  $\Leftarrow$  ”: From  $p N_p \star \xi \leq \xi$  follows by Lemma 3 that  $N + (1/p) \varepsilon_0 < \xi$ . Letting  $p$  tend to infinity we find that  $N < \xi$ .

“(iii)  $\Rightarrow$  ”: The statement holds for  $N$ -potentials and hence for every  $N$ -excessive measure.

“(iii)  $\Leftarrow$  ”: Lemma 3 proves that  $\xi$  is excessive with respect to the elementary kernel  $N_V = \sum_{n=0}^{\infty} (\varepsilon'_V)^n$ , so there exists a net  $(\lambda_\alpha)_{\alpha \in A}$  of positive measures such that  $N_V \star \lambda_\alpha \uparrow \xi$ . Using  $N = (N - N \star \varepsilon'_V) \star N_V$ , we get

$$N \star \lambda_\alpha \uparrow \xi \star (N - N \star \varepsilon'_V),$$

so that  $\xi \star (N - N \star \varepsilon'_{\mathfrak{t}V}) \in E(N)$ . Defining  $a_V = (N - N \star \varepsilon'_{\mathfrak{t}V})(G)$ , we have

$$\frac{1}{a_V} (N - N \star \varepsilon'_{\mathfrak{t}V}) \rightarrow \omega_K \text{ as } V \downarrow K,$$

which implies that

$$\xi = \xi \star \omega_K \in E(N).$$

“(iv)  $\Rightarrow$  ”: By Lemma 2 there exists a net  $(\lambda_\alpha)_{\alpha \in A}$  of positive measures such that  $N \star \lambda_\alpha \uparrow \eta$ ,  $\lambda_\alpha \rightarrow 0$ . Since  $N - N \star \varepsilon'_{\mathfrak{t}V}$  has compact support, we get

$$\eta - \eta \star \varepsilon'_{\mathfrak{t}V} = \lim_A (N - N \star \varepsilon'_{\mathfrak{t}V}) \star \lambda_\alpha = 0.$$

“(iv)  $\Leftarrow$  ”: By (iii)  $\eta \in E(N)$  and hence  $\eta = N \star \mu + \zeta$ , where  $\zeta \in I(N)$ , but since  $\eta = \eta \star \varepsilon'_{\mathfrak{t}V} = (N \star \varepsilon'_{\mathfrak{t}V}) \star \mu + \zeta$ , we get  $\mu = 0$  and then  $\eta \in I(N)$ .

If  $\xi \in E(N)$  has the Riesz decomposition  $\xi = N \star \mu + \eta$ , where  $\eta \in I(N)$ , we find  $p N_p \star \xi = (N - N_p) \star \mu + \eta$ , hence

$$\eta = \lim_{p \rightarrow 0} p N_p \star \xi.$$

**Main result**

**THEOREM 8.** — *Let  $N$  be a non-singular convolution kernel satisfying the domination principle. There exist a non-zero resolvent  $(N_p)_{p>0}$  and a positive measure  $\nu$  such that*

$$(1) \quad N = N_p \star (pN + \varepsilon_0 + \nu) \quad \text{for } p > 0.$$

The resolvent kernel  $\tilde{N} = \lim_{p \rightarrow 0} N_p$  exists, and denoting by  $K$  the compact periodicity group of  $\tilde{N}$ , the measure  $\nu$  can be chosen such that  $\nu \star \varepsilon_x = \nu$  for all  $x \in K$  and  $\nu \in I(N)$ .

*Proof.* — Let  $p > 0$  be fixed. Then  $pN + \varepsilon_0$  is a non-singular convolution kernel satisfying the domination principle and  $N \in E(pN + \varepsilon_0)$ . By Proposition 1, there exist positive measures  $N_p$  and  $\eta_p$  such that

$$N = (pN + \varepsilon_0) \star N_p + \eta_p,$$

where  $\eta_p \in I(pN + \varepsilon_0) = I(N)$ . Furthermore,  $N_p$  and  $\eta_p$  are uniquely determined,  $N_p$  because  $pN + \varepsilon_0$  satisfies the principle of unicity of mass.

For  $q > p > 0$ , we have

$$\begin{aligned} N &= (qN + \varepsilon_0) \star N_q + \eta_q = (q-p)N \star N_q + (pN + \varepsilon_0) \star N_q + \eta_q, \\ &= (q-p)((pN + \varepsilon_0) \star N_p + \eta_p) \star N_q + (pN + \varepsilon_0) \star N_q + \eta_q, \\ &= (pN + \varepsilon_0) \star (N_q + (q-p)N_p \star N_q) + \eta_q + (q-p)\eta_p \star N_q. \end{aligned}$$



The measure  $\eta_q + (q-p)\eta_p \star N_q$  is  $N$ -invariant because both  $\eta_p$  and  $\eta_q$  are so (cf. [7]). By the unicity of the Riesz decomposition with respect to  $pN + \varepsilon_0$ , we conclude

$$(2) \quad \begin{aligned} N_p &= N_q + (q-p)N_p \star N_q & \text{for } 0 < p < q, \\ \eta_p &= \eta_q + (q-p)\eta_p \star N_q & \text{for } 0 < p < q. \end{aligned}$$

This shows that  $(N_p)_{p>0}$  is a resolvent family, and since  $N_p \leq N$  for all  $p$  we get that  $\tilde{N} = \lim_{p \rightarrow 0} N_p$  exists. The resolvent kernel  $\tilde{N}$  is non-zero, because  $\tilde{N} = 0$  would imply that  $N \in I(pN + \varepsilon_0) = I(N)$ , hence that  $N$  is singular. Moreover  $N \geq \eta_p \geq \eta_q$  for  $p < q$  so the limit  $\eta_0 = \lim_{p \rightarrow 0} \eta_p$  exists and belongs to  $I(N)$ . From (2), we get

$$(3) \quad \eta_0 = \eta_q + qN_q \star \eta_0 \quad \text{for } q > 0,$$

which by Proposition 7 shows that  $\eta_0$  is excessive with respect to the resolvent kernel  $\tilde{N}$ , but since from (3)  $\lim_{q \rightarrow 0} qN_q \star \eta_0 = 0$ , the  $\tilde{N}$ -invariant part of  $\eta_0$  is 0. There exists consequently a positive measure  $\nu$  such that  $\eta_0 = \tilde{N} \star \nu$ . The measure  $\nu$  need not be uniquely determined. In fact,  $\tilde{N}$  has a compact periodicity group  $K$ , and denoting the normalized Haar measure of  $K$  by  $\omega_K$ , we have as well  $\eta_0 = \tilde{N} \star (\omega_K \star \nu)$ , so by replacing  $\nu$  by  $\omega_K \star \nu$ , we may and will assume that  $\nu$  is periodic with each  $x \in K$  as period. In this case,  $\nu$  is easily seen to be uniquely determined. From (3) follows

$$\eta_q = \tilde{N} \star \nu - qN_q \star \tilde{N} \star \nu = N_q \star \nu,$$

which implies (1).

Using  $pN_p \star \tilde{N} \leq \tilde{N}$  and that  $\tilde{N} \star \nu$  exists, the convergence Lemma of DENY implies that  $\lim_{p \rightarrow \infty} p\eta_p = \omega_K \star \nu = \nu$ , so  $\nu \in E(N)$ . Since  $\eta_0 = \tilde{N} \star \nu \in I(N)$ , it is easy to see that  $\nu \in I(N)$ , (cf. [7], Corollary 2.4).

LEMMA 9. — *Let  $N_1$  and  $N_2$  be non-zero convolution kernels satisfying the domination principle and  $N_1 < N_2$ . Then*

$$I(N_1) \cap E(N_2) \subseteq I(N_2).$$

*Proof.* — The relation  $<$  being transitive (cf. [4]), it follows that  $E(N_2) \subseteq E(N_1)$ . For  $\eta \in I(N_1) \cap E(N_2)$  and  $V \in \mathcal{V}$ , we then have

$$\eta = {}_{N_1}R_\eta^{\mathbf{C}^V} \leq {}_{N_2}R_\eta^{\mathbf{C}^V} \leq \eta,$$

which proves that  $\eta \in I(N_2)$ .

**THEOREM 10.** — *Let  $N$  be a non-singular convolution kernel satisfying the domination principle, and let  $(N_p)_{p>0}$  and  $\nu$  be as in Theorem 8.*

*Then we have  $\lim_{p \rightarrow 0} N_p = N_0$  and  $N_0 \prec N$ , where  $N_0$  is the regular part of  $N$ . The Riesz decomposition of  $N$  with respect to  $N_0$  is*

$$(4) \quad N = N_0 \star (\varepsilon_0 + \nu) + N^i$$

and

$$N^i \in I(N_0) \cap I(N).$$

*The singular part  $N'$  of  $N$  is  $N$ -invariant and given as*

$$(5) \quad N' = N_0 \star \nu + N^i.$$

*Proof.* — From Theorem 8 we know that the resolvent kernel  $\tilde{N} = \lim_{p \rightarrow 0} N_p$  exists, and also that  $p N_p \star N \leq N$ , hence  $\tilde{N} \prec N$ . Furthermore,  $N$  has the  $\tilde{N}$ -invariant part  $N^i = \lim_{p \rightarrow 0} p N_p \star N$ , so by Lemma 9  $N^i$  is also  $N$ -invariant. Letting  $p \rightarrow 0$  in (1), we find

$$(6) \quad N = \tilde{N} \star (\varepsilon_0 + \nu) + N^i.$$

Since  $\eta_0 = \tilde{N} \star \nu \in I(N)$ , we have  $\tilde{N} \star \nu + N^i \in I(N)$ , so for  $V \in \mathcal{V}$ :

$$\tilde{N} \star \nu + N^i = {}_N R_{N \star \nu + N^i}^{\xi_V} \leq {}_N R_N^{\xi_V},$$

which implies

$$(7) \quad \tilde{N} \star \nu + N^i \leq N',$$

where  $N'$  is the singular part of  $N$ .

For  $V \in \mathcal{V}_K$ , let  $\varepsilon'_{\mathfrak{t}V}$  be a  $\tilde{N}$ -balayaged measure of  $\varepsilon_0$  on  $\mathfrak{t}V$ . Then

$$\xi_V = N \star \varepsilon'_{\mathfrak{t}V} + (\tilde{N} - \tilde{N} \star \varepsilon'_{\mathfrak{t}V}) \star \nu \in E(N),$$

and using (6) and  $\varepsilon'_{\mathfrak{t}V} \star N^i = N^i$ , we find

$$\xi_V = \tilde{N} \star \varepsilon'_{\mathfrak{t}V} + \tilde{N} \star \nu + N^i.$$

In  $\mathfrak{t}V$ , we have  $\tilde{N} \star \varepsilon'_{\mathfrak{t}V} = \tilde{N}$ , hence  $\xi_V = N$  in  $\mathfrak{t}V$ , so by the definition of reduced measure, we get  ${}_N R_N^{\xi_V} \leq \xi_V$ . Letting  $V$  increase to  $G$ , we find using  $\lim_{V \uparrow G} \tilde{N} \star \varepsilon'_{\mathfrak{t}V} = 0$  that

$$N' = \lim_{V \uparrow G} {}_N R_N^{\xi_V} \leq \lim_{V \uparrow G} \xi_V = \tilde{N} \star \nu + N^i,$$

which combined with (7) yields  $N' = \tilde{N} \star \nu + N^i$  and hence  $N_0 = \tilde{N}$ .

With the notation as in Theorem 8 and 10, we further have the following proposition.

PROPOSITION 11. — *If  $N' = N_0 \star v + N^i$  is the Riesz decomposition of the singular part  $N'$  of  $N$  with respect to the regular part  $N_0$  of  $N$ , then either  $v$  or  $N^i$  is zero.*

*Proof.* — Suppose that  $v \neq 0$ . Since  $v \in E(N)$  there exists a net  $(\lambda_\alpha)_{\alpha \in A}$  of positive measures such that  $N \star \lambda_\alpha \uparrow v$ , and since  $N_0 \star v$  exists, this shows that also  $N \star N_0$  exists. Finally, since  $N^i \leq N$  also  $N^i \star N_0$  exists. Using  $N^i \in I(N_0)$ , it follows that

$$N^i = p N_p \star N^i \leq p N_0 \star N^i \quad \text{for all } p > 0,$$

and hence  $N^i = 0$ .

PROPOSITION 12. — *Let  $N$  be a non-singular convolution kernel with regular part  $N_0$ . Then  $N$  and  $N_0$  have the same pseudo-periods. In particular, the group of pseudo-periods for a non-singular convolution kernel is compact.*

*Proof.* — Suppose that  $N_0 \star \varepsilon_x = c N_0$ . Since  $N_0 \prec N$ , it follows that  $N \star \varepsilon_x = c N$ . Conversely, if  $N \star \varepsilon_x = c N$ , then  $N' \star \varepsilon_x = c N'$  because  $N \prec N'$ . Using  $N = N_0 + N'$ , we get  $N_0 \star \varepsilon_x = c N_0$ .

Let  $V \in \mathcal{V}$  be fixed. For every open relatively compact set  $\omega \subseteq G$  such that  $V \subseteq \omega$ , let  $\mu_{\omega \setminus V}$  be a balayaged measure of  $\varepsilon_0$  on  $\omega \setminus V$  with respect to  $N$  such that  ${}_N R_N^{\omega \setminus V} = N \star \mu_{\omega \setminus V}$ . With this notation, we have the following result.

PROPOSITION 13.

(i) *Every accumulation point for the net  $(\mu_{\omega \setminus V})_\omega$  as  $\omega$  increases to  $G$  is a balayaged measure of  $\varepsilon_0$  on  $\complement V$  with respect to  $N_0$ .*

(ii)  ${}_N R_N^{\complement V} = {}_{N_0} R_{N_0}^{\complement V} + N'$ .

(iii) *If  $N$  satisfies the principle of unicity of mass  $\lim_{\omega \uparrow G} \mu_{\omega \setminus V}$  exists and  ${}_{N_0} R_{N_0}^{\complement V} = N_0 \star \lim_{\omega \uparrow G} \mu_{\omega \setminus V}$ .*

*Proof.* — Since  $N \star \mu_{\omega \setminus V} \leq N$ , the net  $(\mu_{\omega \setminus V})_\omega$  is vaguely bounded. Let  $\mu_{\complement V}$  be an accumulation point and assume that  $\mu_{\omega \setminus V} \rightarrow \mu_{\complement V}$  (For notational simplicity we do not write the subnet). From (1) follows

$$(8) \quad N \star \mu_{\omega \setminus V} = p N_p \star N \star \mu_{\omega \setminus V} + N_p \star \mu_{\omega \setminus V} + v \star N_p \star \mu_{\omega \setminus V}.$$

We have  $N \star \mu_{\omega \setminus V} = {}_N R_N^{\omega \setminus V} \uparrow {}_N R_N^{\complement V}$  so the first term on the right-hand side increases to  $p N_p \star {}_N R_N^{\complement V}$ . Since  $N_p \star N$  exists, Deny's convergence Lemma implies that

$$\lim_{\omega} N_p \star \mu_{\omega \setminus V} = N_p \star \mu_{\complement V}.$$

Finally, since  $v \star N_p \in I(N)$ , we have as in the proof of Lemma 4 that  $\lim_{\omega} v \star N_p \star \mu_{\omega \setminus V} = v \star N_p$  so (8) leads to

$$(9) \quad {}_N R_N^{\xi_V} = p N_p \star {}_N R_N^{\xi_V} + N_p \star (\mu_{\xi_V} + v).$$

This shows that  ${}_N R_N^{\xi_V} \in E(N_0)$ , and since  $N' \leqslant {}_N R_N^{\xi_V} \leqslant N$  the  $N_0$ -invariant part of  ${}_N R_N^{\xi_V}$  is equal to  $N^i$  which is the  $N_0$ -invariant part of  $N'$  as well as of  $N$ . Letting  $p \rightarrow 0$  in (9), we get

$${}_N R_N^{\xi_V} = N_0 \star (\mu_{\xi_V} + v) + N^i = N_0 \star \mu_{\xi_V} + N',$$

so it is clear that  $\mu_{\xi_V}$  is a balayaged measure of  $\varepsilon_0$  on  $\mathbb{C}V$  with respect to  $N_0$ .

Let  $\varepsilon'_{\xi_V}$  be a balayaged measure of  $\varepsilon_0$  on  $\mathbb{C}V$  with respect to  $N_0$  such that  ${}_{N_0} R_{N_0}^{\xi_V} = N_0 \star \varepsilon'_{\xi_V}$ . Then  $N_0 \star \varepsilon'_{\xi_V} \leqslant N_0 \star \mu_{\xi_V}$  and with the notation from the proof of Theorem 10, we have

$${}_N R_N^{\xi_V} \leqslant \xi_V = N_0 \star \varepsilon'_{\xi_V} + N',$$

hence

$$(10) \quad {}_N R_N^{\xi_V} = N_0 \star \mu_{\xi_V} + N' \geqslant N_0 \star \varepsilon'_{\xi_V} + N' \geqslant {}_N R_N^{\xi_V}.$$

We shall finally prove (iii). When  $N$  satisfies the principle of unicity of mass,  $N$  and hence also  $N_0$  have no pseudo-periods, so  $N_0$  is a Hunt kernel. Therefore  $\varepsilon'_{\xi_V}$  is uniquely determined by the formula  ${}_{N_0} R_{N_0}^{\xi_V} = N_0 \star \varepsilon'_{\xi_V}$ , and every accumulation point  $\mu_{\xi_V}$  of  $(\mu_{\omega \setminus V})_{\omega}$  is equal to  $\varepsilon'_{\xi_V}$ . Therefore  $\lim_{\omega} \mu_{\omega \setminus V} = \varepsilon'_{\xi_V}$ .

*Remarks*

1° The singular part of  $N + c \varepsilon_0$  is equal to the singular part  $N'$  of  $N$ .

In fact, for  $V \in \mathcal{V}$ , we have observed that

$${}_N R_N^{\xi_V} \geqslant {}_{N+c\varepsilon_0} R_{N+c\varepsilon_0}^{\xi_V},$$

hence  $N' \geqslant (N+c \varepsilon_0)'$ . Since  $N' \in I(N) = I(N+c \varepsilon_0)$ , we also have

$$N' = {}_{N+c\varepsilon_0} R_{N+c\varepsilon_0}^{\xi_V} \leqslant {}_{N+c\varepsilon_0} R_{N+c\varepsilon_0}^{\xi_V},$$

which shows that  $N' \leqslant (N+c \varepsilon_0)'$ .

2° Suppose that  $N' = N_0 \star v$  where  $v \neq 0$ . If  $N$  is shift-bounded (i. e. the set  $\{ N \star \varepsilon_x; x \in G \}$  is vaguely bounded) then  $N_0(G) < \infty$ .

In fact, since  $v \in E(N)$  there exists a non-zero measure  $\lambda \geqslant 0$  such that  $N \star \lambda \leqslant v$  and then

$$N_0 \star N \star \lambda \leqslant N_0 \star v = N' \leqslant N.$$

The shift-boundedness of  $N$  implies that  $N_0 \star \lambda(G) \leqslant 1$ , hence  $N_0(G) < \infty$ .

If  $N' = N_0 \star \nu$  with  $\nu \neq 0$ , and  $N$  is not shift-bounded,  $N_0$  need not be of finite mass as the following example shows:

$$G = \mathbf{R}, N = (1)_{]0, \infty[}(x) + e^x dx.$$

The regular part of  $N$  is the Heaviside kernel  $(1)_{]0, \infty[}$  and  $N' = \nu = e^x$ .

If  $N$  is shift-bounded and  $N_0(G) < \infty$ , then  $N'$  is a  $N_0$ -potential, because  $I(N_0)$  does not contain any shift-bounded non-zero measures.

#### REFERENCES

- [1] DENY (J.). — Noyaux de convolution de Hunt et noyaux associés à une famille fondamentale, *Ann. Inst. Fourier*, Grenoble, t. 12, 1962, p. 643-667.
- [2] ITÔ (M.). — Sur le principe de domination pour les noyaux de convolution, *Nagoya math. J.*, t. 50, 1973, p. 149-173.
- [3] ITÔ (M.). — Caractérisation du principe de domination pour les noyaux de convolution non-bornés, *Nagoya math. J.*, t. 57, 1975, p. 167-197.
- [4] ITÔ (M.). — Sur le principe relatif de domination pour les noyaux de convolution, *Hiroshima math. J.*, t. 5, 1975, p. 293-350.
- [5] ITÔ (M.). — Une caractérisation du principe de domination pour les noyaux de convolution, *Japan. J. Math.*, t. 1, 1975, p. 5-35.
- [6] KISHI (M.). — Positive idempotents on a locally compact abelian group, *Kodai math. Sem. Rep.*, t. 27, 1976, p. 181-187.
- [7] LAUB (J.). — On unicity of the Riesz decomposition of an excessive measure, *Math. Scand.* t. 43, 1978, p. 141-156.