

# BULLETIN DE LA S. M. F.

W. FULTON

S. KLEIMAN

R. PIENE

H. TAI

## **Some intrinsic and extrinsic characterizations of the projective space**

*Bulletin de la S. M. F.*, tome 113 (1985), p. 205-210

[http://www.numdam.org/item?id=BSMF\\_1985\\_\\_113\\_\\_205\\_0](http://www.numdam.org/item?id=BSMF_1985__113__205_0)

© Bulletin de la S. M. F., 1985, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (<http://smf.emath.fr/Publications/Bulletin/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SOME INTRINSIC  
AND EXTRINSIC CHARACTERIZATIONS  
OF THE PROJECTIVE SPACE

BY

W. FULTON (\*), S. KLEIMAN (\*\*),  
R. PIENE and H. TAI (\*\*\*)

RÉSUMÉ. — On démontre le théorème suivant : Le  $n$ -ième plongement de Veronese de  $\mathbb{P}^r$  est la seule immersion  $f: X \rightarrow \mathbb{P}^N$  où  $X$  est lisse, irréductible, de dimension  $r$  et  $N = \binom{n+r}{r} - 1$ , telle que le  $n$ -ième espace osculateur en tout point  $x$  de  $X$  soit l'espace  $\mathbb{P}^N$  tout entier.

Ensuite on conjecture — et démontre dans quelques cas particuliers — qu'une variété  $X$  lisse, irréductible, de dimension  $r$ , est isomorphe à  $\mathbb{P}^r$  si le diviseur anti-canonique  $-K$  est ample, et si, ou bien  $-K$  est numériquement équivalent à  $(r+1)H$  ( $H$  étant un diviseur positif quelconque), ou bien  $\int c_1(O_X(-K))^r = (r+1)^r$ .

ABSTRACT. — The following theorem is proved: The  $n$ -th Veronese embedding of  $\mathbb{P}^r$  is the one and only immersion  $f: X \rightarrow \mathbb{P}^N$  where  $X$  is a smooth, irreducible  $r$ -fold and  $N = \binom{n+r}{r} - 1$ , such that the  $n$ -th osculating space at every point  $x$  of  $X$  is all of  $\mathbb{P}^N$ .

It is also conjectured — and verified in some cases — that a smooth, irreducible  $r$ -fold  $X$  is isomorphic to  $\mathbb{P}^r$  if the anticanonical divisor  $-K$  is ample and if either  $-K$  is numerically equivalent to  $(r+1)H$  for some divisor  $H$ , or if  $\int c_1(O_X(-K))^r = (r+1)^r$ .

(\*) Partially supported by National Foundation Society Grant MCS 81-03400.

(\*\*) Partially supported by the Norwegian Research Council for Science and the Humanities under grant No. C.00.01-01, Matematisk Seminar, and by the Danish Natural Science Research Council.

(\*\*\*) Texte reçu le 7 septembre 1984, révisé le 13 mars 1985.

William FULTON, Dept. of Math., Brown University, Providence, RI 02912, U.S.A.

Steven L. KLEIMAN, Dept. of Math., M.I.T., Cambridge, MA 02139, U.S.A.

Ragni PIENE, Matematisk institutt, PB 1053 Blindern, Oslo 3, Norway.

Hsin-Sheng TAI, Dept. of Math., University of Notre-Dame, Notre-Dame, IN 46556, U.S.A. and Institute of Mathematics, Academia Sinica, Beijing, China.

It is surprising in view of the work of the Italian school of algebraic geometry 70 years ago that the following theorem is new even in the special case characterizing the Veronese surface in the complex projective 5-space.

**THEOREM.** — *Over an algebraically closed field of any characteristic, the  $n$ -th Veronese embedding of  $\mathbb{P}^r$  is the one and only one immersion  $f: X \rightarrow \mathbb{P}^N$  where  $X$  is a smooth, irreducible  $r$ -fold and  $N = \binom{n+r}{r} - 1$ , such that the  $n$ -th osculating space at every point  $x$  of  $X$  is all of  $\mathbb{P}^N$ .*

*Proof.* — Recall that the  $n$ -th osculating space at  $x$  is, by definition, the linear subspace of  $\mathbb{P}^N$  determined by the first  $N+1$  partial derivatives of  $f$  taken with respect to a system of local parameters for  $X$  at  $x$  and evaluated at  $x$ .

Suppose that  $X = \mathbb{P}^r$ . The  $n$ -th Veronese embedding is given in affine coordinates centered at  $x$  by

$$f(x_1, \dots, x_r) = (x_1, \dots, x_r, x_1^2, x_1 x_2, \dots, x_r^2, x_1^3, \dots, x_r^n).$$

Hence the first  $N+1$  derivatives evaluated at  $x$  form the standard frame for  $\mathbb{P}^N$ . Thus the  $n$ -th osculating space is all of  $\mathbb{P}^N$ .

Conversely, for any smooth, irreducible  $r$ -fold  $X$  and for any immersion  $f: X \rightarrow \mathbb{P}^N$  the  $n$ -th osculating space at  $x$  is, in other words, the space determined by the fiber at  $x$  of the following natural map of sheaves on  $X$ :

$$a: \mathcal{O}_X^{N+1} \rightarrow P_X^n(L),$$

where  $L = f^* \mathcal{O}(1)$  and where the target is the twisted sheaf of principal parts (see PIENE [15], §2 and 6). Hence the osculating space is all of  $\mathbb{P}^N$  if and only if  $a$  is injective at  $x$ . Assume  $N = \binom{n+r}{r} - 1$ . Then  $a$  is injective at  $x$  if and only if  $a$  is an isomorphism at  $x$ .

Recall [EGA IV, 16.10.1] that there is a natural exact sequence:

$$(1) \quad 0 \rightarrow (S^n \Omega_X^1) \otimes L \xrightarrow{b} P_X^n(L) \rightarrow P_X^{n-1}(L) \rightarrow 0.$$

It yields via a straightforward calculation the following relation among the first Chern classes:

$$(2) \quad c_1(P_X^n(L)) = \binom{n+r}{r+1} c_1(\Omega_X^1) + \binom{n+r}{r} c_1(L).$$

Assume now that  $a$  is an isomorphism at every point  $x$ . Form the composition  $a^{-1}b$ , dualize it and tensor with  $L$ . The result is a surjection,

$$L^{N+1} \rightarrow S^n T_x.$$

Since  $L$  is ample, therefore  $S^n T_x$  is ample and hence  $T_x$  is ample, by Propositions 2.2 and 2.4 of HARTSHORNE [5]. Consequently, by Theorem 8 of MORI [13],  $X = \mathbb{P}^r$ .

Finally, since  $a$  is an isomorphism, (2) yields the relation:

$$(3) \quad nc_1(T_x) = (r+1)c_1(L).$$

Since  $X = \mathbb{P}^r$ , therefore  $L = O_x(n)$ . Hence  $f$  is the  $n$ -th Veronese embedding, possibly followed by a projection and then an inclusion. However, each  $n$ -th osculating space is all of  $\mathbb{P}^N$ . So  $f$  is simply the Veronese embedding.

There is another proof that  $X = \mathbb{P}^r$  when  $a$  is an isomorphism, which works in the following three cases:

- (i)  $r = 1, 2$ ;
- (ii)  $r = 3$  and the characteristic is 0;
- (iii)  $n$  and  $r + 1$  are relatively prime and the characteristic is 0.

More generally, if  $r = 1, 2$ , or if the characteristic is 0, or if  $n$  and  $r + 1$  are relatively prime, then this proof reduces the problem to establishing the following conjecture, which may be of interest in its own right (for other, related, conjectures, see FUJITA [2]).

CONJECTURE. — *Over an algebraically closed field of any characteristic, a smooth, irreducible  $r$ -fold  $X$  is isomorphic to  $\mathbb{P}^r$  if an anticanonical divisor  $-K$  is ample and if either one of the following hypotheses is satisfied:*

(a) *there exists a divisor  $H$  such that  $(r+1)H$  is numerically equivalent to  $-K$ :*

$$(b) \quad \int c_1(O_x(-K))^r = (r+1)^r.$$

*Second proof.* — First, (3) says that  $-K$  is ample. So, if  $r = 1$ , then  $X = \mathbb{P}^1$ , and the proof is complete in this case. Secondly, suppose that  $n$  and  $r + 1$  are relatively prime, say  $un + v(r + 1) = 1$ . Say  $L = O_x(D)$  and set  $H = v(-K) + uD$ . Then (3) implies that  $(r + 1)H$  is numerically equivalent to  $-K$  (in fact, linearly equivalent, if the Chern classes are taken modulo linear equivalence). Thus (a) holds. Thirdly, the following argu-

ment shows that, if  $r=2$  or if the characteristic is 0, then (b) holds; in fact, it shows that then all the Chern numbers of  $X$  are the same as those of  $\mathbb{P}^r$ . To complete the alternative proof, it will then suffice to establish the conjecture in the asserted cases, and this is done in the proposition below.

Set  $c_i = c_i(T_X)$ . Since  $a$  is an isomorphism,

$$\int c_1^{i_1} \dots c_j^{i_j} \cdot c_j(P_X^n(L)) = 0 \quad \text{whenever } i_1 + 2i_2 + \dots + ji_j = r.$$

Now, (1) and (3) imply that, for a certain nonzero integer  $q$  and a certain polynomial  $Q$  with rational coefficients,

$$c_j(P_X^n(L)) = qc_j + Q(c_{j-1}, \dots, c_1).$$

Combined, the two equations yield a relation among the Chern numbers

$$c(k_1, \dots, k_r) = \int c_1^{k_1} \dots c_r^{k_r} \quad \text{where } k_1 + 2k_2 + \dots + rk_r = r.$$

Index the relation by the  $r$ -tuple  $(i_1, \dots, i_{j-1}, i_j + 1, 0, \dots, 0)$ . Then, obviously, there is exactly one relation for each  $r$ -tuple  $(m_1, \dots, m_r)$  such that  $m_1 + 2m_2 + \dots + rm_r = r$ . Only one relation is trivial, the one with index  $(r, 0, \dots, 0)$ . The remaining relations are linearly independent; indeed, if the  $r$ -tuples are put in reverse lexicographical order (that is, compare them starting with  $m_r$ ), then the relation with index  $(m_1, \dots, m_r)$  involves a Chern number that does not appear in any of the successive relations, namely,  $c(m_1, \dots, m_r)$ .

The Riemann-Roch theorem gives another relation, which expresses the Euler characteristic  $\chi(O_X)$  in terms of the Chern numbers. It is independent of the other relations if  $\chi(O_X) \neq 0$ , because then it is the only relation with a nonzero constant term. Hence the relations determine the Chern numbers. So the numbers are those of  $\mathbb{P}^r$ , provided  $\chi(O_X) = 1$ .

Suppose  $r=2$ . Then (1) yields the formula

$$c_2(P_X^n(L)) = \binom{n+3}{4} (c_2 + [(n^2 - 1)/3]c_1^2 - 2nc_1 \cdot c_1(L) + 3c_1(L)^2).$$

Hence, in this case, the relations are as follows:

$$\begin{aligned} 3c_2 &= c_1^2, \\ c_2 + c_1^2 &= 12\chi(O_X). \end{aligned}$$

Since  $-K$  is ample,  $c_1^2 \geq 1$ . Hence,  $\chi(O_X) \geq 1$ . On the other hand,

$$h^2(O_X) = h^0(K) \quad \text{and} \quad h^0(K) = 0,$$

because of duality and because  $-K$  is ample. Since  $h^0(O_X) = 1$ , it follows that  $\chi(O_X) = 1$ .

For any  $r$  in characteristic 0, Kodaira's vanishing theorem yields  $h^i(O_X) = 0$  for  $i \geq 1$ . Since  $h^0(O_X) = 1$ , therefore  $\chi(O_X) = 1$ .

It now remains to prove the following proposition.

**PROPOSITION.** — (1) (FUJITA [1]). *Under hypothesis (a), the conjecture holds in characteristic 0.*

(2) *Under hypothesis (b), the conjecture holds for  $r = 1, 2$  in any characteristic, and for  $r = 3$  in characteristic 0.*

Indeed, if  $r = 1$ , then  $X = \mathbb{P}^1$ , just because  $-K$  is ample; in particular, (2) holds. If  $r = 2$  or 3, then (2) holds by the classification of Del Pezzo surfaces (e. g., Theorem 24.4 (i) of MANIN [12]) and by the classification of Fano 3-folds (ISKOVSKIĬ [7], [8], MORI-MUKAI [14], L'VOVSKIĬ [11]). As to (1),  $h^i(K + jH) = 0$  for  $i, j \geq 1$  by Kodaira's vanishing theorem and for  $i = 0$  and  $j \leq r$  because  $-K$  is ample and

$$(K + jH) \cdot (-K)^{r-1} < 0.$$

Furthermore,  $\chi(O_X) = 1$ . Since  $\chi(mH)$  is of degree  $r$ , therefore it must be  $\binom{m+r}{r}$ . Therefore  $H^r = 1$  and  $h^0(H) = r + 1$ . By Theorem 1 of

GOREN [3],  $X = \mathbb{P}^r$ . (While Goren does not assume  $X$  to be Cohen-Macaulay, he uses this hypothesis implicitly in the last line of the proof of Lemma 2. This interesting characterization of  $\mathbb{P}^n$  has been rediscovered at least twice after Goren, by KOBAYASHI and OCHIAI [9] and by FUJITA [1]. Earlier HIRZEBRUCH and KODAIRA [6], Theorem 6, gave a weaker form of Goren's result, which is insufficient for our purposes.) FUJITA [1] proved (1), although he used linear equivalence in place of numerical equivalence. He determined the polynomial  $\chi(mH)$  much as above, and he reproved Goren's theorem. KOLLAR [10] gave a similar proof, using the stronger form of Mori's theorem instead of Goren's theorem.

*Remark.* — The Hilbert polynomial of the  $n$ -th Veronese embedding of  $\mathbb{P}^r$  does not always suffice to characterize the embedding. For example, the surface  $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^5$  embedded by  $O(2,2)$  has the same Hilbert polynomial (in fact the same Hilbert function) as the Veronese surface in

$\mathbb{P}^5$ . Conceivably, the case  $n=2$ ,  $r=2$  is the only case in which the embedding is not characterized by its Hilbert polynomial.

## REFERENCES

- [1] FUJITA (T.). — On the structure of polarized varieties with  $\Delta$ -genera zero, *J. Fac. Sc., Univ. Tokyo, Sect. IA Math.*, Vol. 22, 1975, pp. 103-115.
- [2] FUJITA (T.). — On topological characterizations of complex projective spaces and affine linear spaces, *Proc. Japan Acad.*, Vol. 56, Ser. A, 1980, pp. 231-234.
- [3] GOREN (R.). — Characterization and algebraic deformations of projective space, *J. Math.*, Kyoto Univ., Vol. 8-1, 1968, pp. 41-47.
- [4] GROTHENDIECK (A.) [EGA IV]. — Éléments de géométrie algébrique IV, *Publ. Math. I.H.E.S.*, No. 32, 1967.
- [5] HARTSHORNE (R.). — Ample vector bundles, *Publ. Math. I.H.E.S.*, Vol. 29, 1966, pp. 63-94.
- [6] HIRZEBRUCH (F.) and KODAIRA (K.). — On the complex projective spaces, *J. Math. pures et appl.*, Ser. 9, Vol. 36, 1957, pp. 201-216.
- [7] ISKOVSKIĖ (V. A.). — Fano 3-folds I, *Izv. Akad. Nauk.*, Vol. 41, 1977; *Engl. transl. Math. U.S.S.R., Izv.*, Vol. 11, 1977, pp. 485-527.
- [8] ISKOVSKIĖ (V. A.). — Fano 3-folds II, *Izv. Akad. Nauk.*, Vol. 42, 1978; *Engl. transl. Math. U.S.S.R., Izv.*, Vol. 12, 1978, pp. 469-506.
- [9] KOBAYASHI (S.) and OCHIAI (T.). — Characterizations of complex projective spaces and hyperquadrics, *J. Math. Kyoto Univ.*, Vol. 13-1, 1973, pp. 31-47.
- [10] KOLLAR (J.). — Fano varieties of large dimension and large index, *Vestnik Moskovskogo Univ.*, No. 3, 1981, pp. 31-34.
- [11] L'VOVSKIĖ (S. M.). — Boundedness of the degree of Fano threefolds, *Izv. Akad. Nauk. S.S.S.R., Ser. Mat.*, Vol. 45, 1981; *Engl. transl. Math. U.S.S.R. Izv.*, Vol. 19, 1982, pp. 521-558.
- [12] MANIN (Yu. I.). — *Cubic Forms*, North Holland Publishing Company, Amsterdam-London, 1974.
- [13] MORI (S.). — Projective manifolds with ample tangent bundles, *Ann. Math.*, Vol. 110, 1979, pp. 593-606.
- [14] MORI (S.) and MUKAI (S.). — Classification of Fano 3-folds with  $B_2 \geq 2$ , *Manuscr. math.*, Vol. 36, 1981, pp. 147-162.
- [15] PIENE (R.). — Numerical characters of curves in projective  $n$ -space, *Real and complex singularities*, Oslo 1976, P. Holm, Sijthoff & Noordhoff, Alpen aan den Rijn, 1977, pp. 475-495.