

COMPOSITIO MATHEMATICA

N. J. S. HUGHES

Steinitz' exchange theorem for infinite bases

Compositio Mathematica, tome 15 (1962-1964), p. 113-118

http://www.numdam.org/item?id=CM_1962-1964__15__113_0

© Foundation Compositio Mathematica, 1962-1964, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Steinitz' Exchange Theorem for Infinite Bases

by

N. J. S. Hughes

Given a system in which a suitable relation of dependence is defined, we give a construction (assuming well ordering), by which some of the elements of any basis may be replaced, in a one-one manner, by all the elements of any independent subset to give a new basis.

1. Definitions and notation

We call the set S a *dependence space* if there is defined a set Δ , whose members are finite subsets of S , each containing at least 2 elements, and if the Transitivity Axiom (below) is satisfied.

We shall use a, b, c, x, y (with or without suffixes) to denote elements of S and A, B, C, X for subsets of S and also i, j for suffixes and I, J for sets of suffixes and n will always be a positive integer.

$A + B$ will denote the union of the *disjoint* sets A and B and $A - B$ the set of those elements of A which are not in B .

We call A *directly dependent* if $A \in \Delta$.

If either $x \in A$ or there exist distinct x_0, x_1, \dots, x_n , such that

$$(x_0, x_1, \dots, x_n) \in \Delta, \quad (1)$$

where $x_0 = x$ and $x_1, \dots, x_n \in A$, we call x *dependent* on A , denoted by $x \sim \sum A$, and *directly dependent* on (x) or (x_1, \dots, x_n) respectively.

We say that A is *dependent* if (1) is satisfied for some distinct $x_0, x_1, \dots, x_n \in A$, and otherwise that A is *independent*.

If A is independent and, for any $x \in S$, $x \sim \sum A$, then A is a *basis* of S .

If $A = (a_i)_{i \in I}$ then $\sum A$ and $\sum_{i \in I} a_i$ are equivalent symbols. Also $\sum A + \sum B$ and $\sum(A \cup B)$ are equivalent symbols.

If either $x = a$ or (1) is satisfied for some $n \geq 1$, with $x_0 = x$, $x_1 = a$, and, for $2 \leq m \leq n$, $x_m \in C$, we write

$$x \sim (a) + \sum C. \quad (2)$$

Clearly, (2) implies $a \sim (x) + \sum C$.

We assume the following Transitivity Axiom:

if $x \sim \sum A$ and, for all $a \in A$, $a \sim \sum B$, then $x \sim \sum B$.

In particular, we may take S to be the set of all non-zero elements of a vector space over a field F , and have (1) if and only if

$$\xi_0 x_0 + \dots + \xi_n x_n = 0$$

for some non-zero ξ_0, \dots, ξ_n in F .

2. Well ordered subsets

We now assume that $A = (a_i)_{i \in I}$ is well ordered, I being also well ordered, and assume the Principle of Transfinite Induction in the form:

$(i \in I)$, $P(i)$, (i.e. $P(i)$ is true for all $i \in I$), if

$$(i \in I), (j < i) \Rightarrow P(j) \cdot \Rightarrow P(i).$$

LEMMA 1

If $(i \in I)$, $a_i \sim \sum_{j < i} a_j + \sum C$, then $(i \in I)$, $a_i \sim \sum C$.

This is easily proved by Transfinite Induction.

LEMMA 2

If $A + C$ is a basis of S and

$$(i \in I), x_i \sim (a_i) + \sum_{j < i} a_j + \sum C, \quad (1)$$

then the x_i are distinct and not in C , $X + C$ is a basis of S , where $X = (x_i)_{i \in I}$, and

$$(i \in I), a_i \sim (x_i) + \sum_{j < i} x_j + \sum C. \quad (2)$$

Also, if

$$y \sim (a_i) + \sum_{j < i} a_j + \sum C, \quad (3)$$

then

$$y \sim (x_i) + \sum_{j < i} x_j + \sum C. \quad (4)$$

From (1), we have

$$(i \in I), a_i \sim (x_i) + \sum_{j < i} a_j + \sum C, \quad (5)$$

and hence, by Transfinite Induction,

$$(i \in I), a_i \sim \sum_{j \leq i} x_j + \sum C. \quad (6)$$

From (3) and (6), we have

$$y \sim \sum_{j \leq i} x_j + \sum C. \quad (7)$$

If

$$y \sim \sum_{j < i} x_j + \sum C, \quad (8)$$

then, by (1),

$$y \sim \sum_{j < i} a_j + \sum C,$$

and hence, since, by (3),

$$\begin{aligned} a_i &\sim (y) + \sum_{j < i} a_j + \sum C, \\ a_i &\sim \sum_{j < i} a_j + \sum C, \end{aligned} \quad (9)$$

which is a contradiction, since $A + C$ is independent.

From (7) and the falsity of (8), we have (4), and then, putting $y = a_i$, also (2).

If 2 of the x_i were equal, or if an x_i were in C , or if $X + C$ were dependent, we would have (since C is independent) a relation of the form:

$$x_i \sim \sum_{j < i} x_j + \sum C.$$

Then, by (1), we would have

$$x_i \sim \sum_{j < i} a_j + \sum C,$$

and, by (5), again (9).

Thus $X + C$ is independent and, by (6), is a basis of S .

3. Proof of Steinitz' exchange theorem

THEOREM

If A is a basis and B an independent subset (both being well ordered) of the dependence space S , then there is a definite subset A' of A , such that $B + (A - A')$ is also a basis of S , and a definite one-one correspondence between A' and B .

If B is a basis of S , then $A' = A$.

We shall suppose that $A = (a_i)_{i \in I}$ where I is well ordered and shall define successively disjoint subsets $I(1), I(2), \dots$ and, for all i in their union, distinct elements b_i of B .

We suppose that $I(1), \dots, I(p)$ have been defined and also, for all $i \in I(1) + \dots + I(p)$, distinct $b_i \in B$.

We let

$$J(p) = I - (I(1) + \dots + I(p)), \quad (1)$$

$$A^p = (a_i^p)_{i \in J(p)}, \text{ where, } (i \in J(p)), a_i^p = a_i, \quad (2)$$

$$(q = 1, \dots, p), B^q = (b_i)_{i \in I(q)}. \quad (3)$$

We shall further suppose that A_p , defined by

$$A_p = A^p + B^1 + \dots + B^p \quad (4)$$

is a basis of S .

If $p = 0$, we define $J(0) = I$, $A^0 = A_0 = A$.

If $b \in B - (B^1 + \dots + B^p)$, since A_p is a basis of S and B is independent, we have a relation of the form:

$$b \sim (a_i^p) + \sum_{j < i} a_j^p + \sum B^1 + \dots + \sum B^p. \quad (5)$$

In (5), $i = i(p+1, b)$ may, by the well ordering of $J(p)$, be supposed the least possible, but it follows easily from the independence of A_p that the set of elements, on which b is directly dependent, is in fact unique.

We now define $I(p+1)$ to be the set of all i in $J(p)$, such that $i = i(p+1, b)$, for some $b \in B - (B^1 + \dots + B^p)$, and b_i to be the first such b (in the well ordering of B) and may replace p by $p+1$ in the definitions (1) to (4).

We then have

$$(i \in I(p+1)), b_i \sim (a_i^p) + \sum_{j < i} a_j^p + \sum B^1 + \dots + \sum B^p. \quad (6)$$

By Lemma 2, with A^p for A , $B^1 + \dots + B^p$ for C and

$$(i \in I(p+1)), x_i = b_i, (i \in J(p+1)), x_i = a_i, \quad (7)$$

A_{p+1} is a basis of S .

By the last part of Lemma 2, (with $i = i(p+1, b)$), and (7), we have

$$(b \in B - (B^1 + \dots + B^{p+1}), i(p+2, b) < i(p+1, b). \quad (8)$$

The process of successively defining the subsets $I(1), I(2), \dots$ of I and the corresponding disjoint subsets B^1, B^2, \dots of B may be continued either until, for some p , $B^1 + \dots + B^p = B$ or to give an infinite sequence of subsets.

In the latter case $B = B^1 + B^2 + \dots$, for, by (8), if $b \in B - (B^1 + B^2 + \dots)$,

$$i(1, b), i(2, b), \dots$$

would be an infinite, strictly descending sequence of members of I .

In each case we take $A' = (a_i)_{i \in I(1) + I(2) + \dots}$ and the correspondence $a_i \leftrightarrow b_i$ is one-one between A' and B .

In the former case, $A - A' = A^p$ and, by (4), $B + (A - A') = A_p$, and is therefore a basis of S .

In the latter case, $A - A' \subseteq A^p$, for all $p \geq 0$, and we see, by (4), that any finite subset of $B + (A - A')$ is contained in A_p , for sufficiently large p . Thus $B + (A - A')$ is independent.

Since

$$a_i \sim \sum_{j < i} a_j + \sum(A - A') + \sum B$$

is trivial if $i \in I - (I(1) + I(2) + \dots)$ and follows from (6) if $i \in I(p+1)$, for any $p \geq 0$, by Lemma 1,

$$(i \in I), a_i \sim \sum(A - A') + \sum B.$$

Thus, being independent, $B + (A - A')$ is a basis of S .

Finally, since a basis is a maximal independent subset, if B is a basis of S , $A - A'$ is empty and $A' = A$.

4. Rank

Since the bases of S coincide with its maximal independent subsets, S , assumed to be well ordered, has at least one basis, and by the last part of the Theorem, any 2 bases have the same cardinal number, which may be called the rank of S (with respect to Δ).

From the example at the end of § 1, we see that a vector space over a field has a unique rank.

If G is an additive Abelian group, we let S be the set of elements of infinite order and $(x_0, \dots, x_n) \in \Delta$ if and only if, for some non-zero integers N_0, \dots, N_n ,

$$N_0 x_0 + \dots + N_n x_n = 0.$$

It now follows that the rank of G is unique (Kurosh, p. 140).

Now let G be a p -primary additive Abelian group and r be a positive integer. Let H be the subset of G generated by the union of the set of all $g \in G$, such that $p^{r-1}g = 0$ and the set of all g , such that $g = pg'$, for some $g' \in G$.

We take S to be the set of all elements of G , whose orders are exactly p^r and which are not in H , and $(x_0, \dots, x_n) \in \mathcal{A}$, if and only if, for some integers N_0, \dots, N_n prime to p ,

$$N_0 x_0 + \dots + N_n x_n \in H.$$

If G can be expressed as a direct sum of cyclic groups, we see easily that the set of generators of the cyclic groups of order p^r is a basis of S and hence that the cardinal number of such summands is a group invariant (Kurosh, p. 174).

REFERENCE

A. G. KUROSH

Theory of Groups, Vol. 1, Chelsea Publishing Co., New York, 1955.

University College,
Cathays Park, Cardiff

(Oblatum 30-3-62).