

# COMPOSITIO MATHEMATICA

FU CHENG HSIANG

**On the absolute summability factors of  
Fourier series at a given point**

*Compositio Mathematica*, tome 17 (1965-1966), p. 156-160

[http://www.numdam.org/item?id=CM\\_1965-1966\\_\\_17\\_\\_156\\_0](http://www.numdam.org/item?id=CM_1965-1966__17__156_0)

© Foundation Compositio Mathematica, 1965-1966, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# On the absolute summability factors of Fourier series at a given point

by

Fu Cheng Hsiang

## 1.

A series  $\sum a_n$  is said to be absolutely summable ( $A$ ) or summable  $|A|$  if the function

$$f(x) = \sum_{n=0}^{\infty} a_n x^n$$

is of bounded variation in the interval  $(0,1)$  Let  $\sigma_n^\alpha$  denote the  $n$ -th Cesàro mean of order  $\alpha$  of the series  $\sum a_n$ , i.e.,

$$\sigma_n^\alpha = \frac{1}{(\alpha)_n} \sum_{k=0}^n (\alpha)_k a_{n-k}, \quad (\alpha)_k = \Gamma(k+\alpha+1)/\Gamma(k+1)\Gamma(\alpha+1).$$

If the series

$$\sum |\sigma_n^\alpha - \sigma_{n-1}^\alpha|$$

converges, then, we say that the series  $\sum a_n$  is absolutely summable  $(C, \alpha)$  or summable  $|C, \alpha|$ . It is known that [2] if a series is summable  $|C|$ , then it is also summable  $|A|$ .

## 2.

Suppose now that  $f(x)$  is a function integrable in the sense of Lebesgue and periodic with period  $2\pi$ . Let

$$f(x) \sim \frac{1}{2}a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) \equiv \sum A_n(x).$$

Whittaker [5] proved that the series

$$\sum_{n=1}^{\infty} A_n(x)/n^\alpha \quad (\alpha > 0)$$

is summable  $|A|$  almost everywhere. Prasad [5] improved this result by showing that the series  $\sum A_n(x)$  when multiplied by one of the factors:

$$1/(\log n)^{1+\varepsilon}, 1/\log n(\log^2 n)^{1+\varepsilon}, \dots, \\ 1/(\log n) \cdot (\log^2 n) \dots (\log^{k-1} n)(\log^k n)^{1+\varepsilon},$$

where  $\varepsilon$  is any positive quantity and  $\log^1 n = \log n$ ,  $\log^k n = \log(\log^{k-1} n)$ , is summable  $|A|$  at the point  $x$ .

Let  $(\lambda_n)$  be a convex and bounded sequence [3], Chow [1] has demonstrated that the series

$$\sum A_n(x)\lambda_n$$

is summable  $|C, 1|$  almost everywhere, if the series  $\sum n^{-1}\lambda_n$  converges.

In the present note, we are interested particularly in the case of  $|C, 1|$  summability. For a fixed point of  $x$ , we write

$$\varphi(t) = \varphi_x(t) = f(x+t) + f(x-t) - 2f(x).$$

We are going to establish at first the following

**THEOREM 1.** *If*

$$\Phi(t) = \int_0^t |\varphi(u)| du = O\left(\frac{t}{\log 1/t}\right)$$

as  $t \rightarrow +0$ , then the series

$$\sum_{n=2}^{\infty} \frac{A_n(x)}{(\log n)^{1+\varepsilon}}$$

is summable  $|C, 1|$  for every  $\varepsilon > 0$ .

### 3.

In the proof of the theorem, we require the following lemmas.

**LEMMA 1** [4]. *Let  $\alpha > -1$  and let  $\tau_n^\alpha$  be the  $n$ -th Cesàro mean of order  $\alpha$  of the sequence  $(na_n)$ , then*

$$\tau_n^\alpha = n(\sigma_n^\alpha - \sigma_{n-1}^\alpha),$$

where  $\sigma_n^\alpha$  is the  $n$ -th Cesàro mean of order  $\alpha$  of the series  $\sum a_n$ .

**LEMMA 2.** *Write*

$$S_n(t) = \sum_{k=0}^n (n+2-k) \cos(n+2-k)t,$$

then

$$S_n(t) = O \begin{cases} nt^{-1} & (nt \geq 1), \\ n^2 & (\text{for all } t). \end{cases}$$

In fact, we have

$$\begin{aligned} S_n(t) &= \mathcal{J} \left\{ \frac{d}{dt} \left( \overline{e^{t(n+2)t} \sum_{k=0}^n e^{-kt}} \right) \right\} \\ &= \mathcal{J} \left\{ \frac{d}{dt} \left( \frac{e^{t(n+2)t}}{1-e^{-t}} - \frac{e^{tt}}{1-e^{-t}} \right) \right\} \\ &= O(nt^{-1}) + O(t^{-2}) \\ &= O(nt^{-1}), \end{aligned}$$

if  $nt \geq 1$ . This proves the first part of the lemma. And the second part of the lemma is evident.

#### 4.

Now, we are in a position to prove the theorem. We use

$$A_n(x) = \frac{2}{\pi} \int_0^\pi \varphi(t) \cos nt \, dt.$$

Let  $\tau_n(x)$  be the  $n$ -th Cesàro mean of first order of the sequence  $\{nA_n(x)(\log n)^{-(1+\varepsilon)}\}$ , then

$$\frac{\pi}{2} \tau_n(x) = \int_0^\pi \varphi(t) \frac{1}{n+1} \sum_{\nu=0}^n \frac{(\nu+2) \cos(\nu+2)t}{(\log(\nu+2))^{1+\varepsilon}} \, dt.$$

Abel's transformation gives

$$\begin{aligned} \frac{\pi}{2} \tau_n(x) &= \int_0^\pi \varphi(t) \frac{1}{n+1} \left\{ \sum_{\nu=0}^n S_\nu(t) \Delta \frac{1}{(\log(\nu+2))^{1+\varepsilon}} \right\} \, dt \\ &\quad + \int_0^\pi \varphi(t) \frac{1}{n+1} \cdot \frac{S_n(t)}{(\log(n+3))^{1+\varepsilon}} \, dt \\ &= I_1 + I_2, \end{aligned}$$

say. By Lemma 2, we have

$$\frac{1}{n+1} \left\{ \sum_{\nu=0}^n S_\nu(t) \Delta \frac{1}{(\log(\nu+2))^{1+\varepsilon}} \right\} = O \left\{ \begin{array}{l} \frac{1}{t(\log n)^{2+\varepsilon}} \quad (nt \geq 1), \\ \frac{n}{(\log n)^{2+\varepsilon}} \quad (\text{for all } t). \end{array} \right.$$

Thus, on writing

$$I_1 = \int_0^{1/n} + \int_{1/n}^\pi = I_3 + I_4,$$

say, we see that

$$I_3 = O \left\{ \frac{n}{(\log n)^{2+\varepsilon}} \int_0^{1/n} |\varphi| dt \right\} = O \left\{ \frac{1}{(\log n)^{2+\varepsilon}} \right\},$$

$$I_4 = O \left\{ \frac{1}{(\log n)^{2+\varepsilon}} \int_{1/n}^{\pi} \frac{|\varphi|}{t} dt \right\}.$$

Now,

$$\int_{1/n}^{\pi} \frac{|\varphi|}{t} dt = \left( \frac{\Phi}{t} \right)_{1/n}^{\pi} + \int_{1/n}^{\pi} \frac{\Phi}{t^2} dt$$

$$= O(\log n).$$

It follows that  $I_4 = O\{(\log n)^{-(1+\varepsilon)}\}$ . As before, we write

$$I_2 = \int_0^{1/n} + \int_{1/n}^{\pi} = I_5 + I_6,$$

say. Then,

$$I_5 = O \left\{ \frac{n}{(\log n)^{1+\varepsilon}} \int_0^{1/n} |\varphi| dt \right\}$$

$$= O \left\{ \frac{1}{(\log n)^{1+\varepsilon}} \right\}.$$

And

$$I_6 = O \left\{ \frac{1}{(\log n)^{1+\varepsilon}} \int_{1/n}^{\pi} \frac{|\varphi|}{t} dt \right\}.$$

Now,

$$\int_{1/n}^{\pi} \frac{|\varphi|}{t} dt = \left( \frac{\Phi}{t} \right)_{1/n}^{\pi} + \int_{1/n}^{\pi} \frac{\Phi}{t^2} dt$$

$$= O(1) + O \left( \int_{1/n}^{\pi} \frac{dt}{1/n + \log 1/t} \right)$$

$$= O(\log^2 n).$$

It follows that

$$I_6 = O \left( \frac{\log^2 n}{(\log n)^{1+\varepsilon}} \right).$$

By Lemma 1, we need only to prove that  $\sum |\tau_n(x)|/n$  converges. And from the above analysis, it concludes that

$$\sum |\tau_n(x)|/n = O \left\{ \sum_{n=2}^{\infty} \frac{\log^2 n}{n(\log n)^{1+\varepsilon}} \right\}$$

$$= O(1).$$

This completes the proof of Theorem 1.

## 5.

For the conjugate series

$$\sum_{n=1}^{\infty} (b_n \cos nx - a_n \sin nx) \equiv \sum B_n(x),$$

we can derive an analogous theorem. Write

$$\bar{\Psi}(t) = \int_0^t |\psi(u)| du \equiv \int_0^t |f(x+u) - f(x-u)| du.$$

Then, we get the following

**THEOREM 2.** *If*

$$\bar{\Psi}(t) = O\left(\frac{t}{\log 1/t}\right)$$

*as*  $t \rightarrow +0$ , *then the series*

$$\sum \frac{B_n(x)}{(\log n)^{1+\varepsilon}}$$

*is summable*  $|C, 1|$  *at*  $x$ .

## REFERENCES

- H. C. CHOW,  
 [1] On the summability factors of Fourier series, Journ. London Math. Soc., 16 (1941), 215—220.
- M. FEKETE,  
 [2] On the absolute summability (A) of infinite series, Proc. Edinburgh Math. Soc. (2), 3 (1932), 132—134.
- S. IZUMI and T. KAWATA,  
 [3] Notes on Fourier series III, absolute summability, Proc. Imperial Academy of Japan, 14(1938), 32—35.
- E. KOGBETLIANTZ,  
 [4] Sur les séries absolument sommables par la methode des moyennes arithmétiques, Bull. Sciences Math. (2), 49 (1925), 234—256.
- B. N. PRASAD,  
 [5] On the summability of Fourier series and the bounded variation of power series, Proc. London Math. Soc., 14 (1939), 162—168.
- T. PATI,  
 [6] On the absolute summability factors of Fourier series, Indian Journ. Math., 1 (1958), 41—54.