

COMPOSITIO MATHEMATICA

J. DE VRIES

Equivalence of almost periodic compactifications

Compositio Mathematica, tome 22, n° 4 (1970), p. 453-456

<http://www.numdam.org/item?id=CM_1970__22_4_453_0>

© Foundation Compositio Mathematica, 1970, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

EQUIVALENCE OF ALMOST PERIODIC COMPACTIFICATIONS

by

J. de Vries

1. Introduction

1.1. Purpose of this note is to give a simple proof of [1], Theorem 5.5 (see Corollary 2.4 below).

Let S, T be semitopological semigroups ([2], p. 1). We do not assume S and T to have an identity. The Banach space of almost periodic (a.p.) functions on S is denoted by $A(S)$ and the Banach space of weakly almost periodic functions on S is denoted by $W(S)$. If $\phi : S \rightarrow T$ is a continuous homomorphism, then the induced mapping $f \mapsto f \circ \phi$ of $C(T)$ into $C(S)$ is denoted by $\tilde{\phi}$.

Recall that $\tilde{\phi}[C(T)] \subset W(S)$ if T is a compact semitopological semigroup, and that $\tilde{\phi}[C(T)] \subset A(S)$ if T is a compact topological semigroup ([1], lemma 5.2).

1.2. An ordered pair (ϕ, T) is called an *almost periodic compactification* (a *weakly almost periodic compactification*) of S , if the following conditions are satisfied:

- (i) T is a compact topological (semitopological) Hausdorff semigroup.
- (ii) $\phi : S \rightarrow T$ is a continuous homomorphism such that for each $f \in A(S)$ ($f \in W(S)$) there is a unique $\tilde{f} \in C(T)$ with $f = \tilde{f} \circ \phi$.

Note that (ii) is equivalent with

- (ii)' $\phi : S \rightarrow T$ is a continuous homomorphism with dense image in T , and $\tilde{\phi}[C(T)] = A(S)$ ($\tilde{\phi}[C(T)] = W(S)$).

(The proof depends on 1.1 and the fact, that T is a completely regular topological space.)

1.3. Let S be a semitopological semigroup. Two a.p. compactifications (w.a.p. compactifications) (ϕ_1, T_1) and (ϕ_2, T_2) are called *equivalent*, if there is a topological isomorphism ψ of T_1 onto T_2 such that $\psi \circ \phi_1 = \phi_2$.

We shall prove, that two a.p. compactifications (w.a.p. compactifications) of a semitopological semigroup are equivalent ([1], Corollary 5.6); in fact, this is an easy corollary of the 'universal property' of the a.p. and

w.a.p. compactifications, explained in Theorem 2.2 of this note. From this, Theorem 5.5 of [1] is also easily derived.

All semitopological semigroups are not supposed to have an identity (so the existence of the compactifications is not a priori guaranteed).

2. The main theorem

2.1. LEMMA. *Let X be a uniform space, Y a compact Hausdorff topological space. A function $f : X \rightarrow Y$ is uniformly continuous (Y with its unique uniformity) if and only if $g \circ f : X \rightarrow [0, 1]$ is uniformly continuous for all continuous $g : Y \rightarrow [0, 1]$.*

PROOF: It is known that Y may be regarded as a (closed) subset of a topological product of copies of the interval $[0, 1]$ such that the restrictions to Y of the canonical projections are the continuous functions of Y into $[0, 1]$. The lemma now follows from [3], Ch. 6, Theorem 10.

2.2. THEOREM. *Let (ϕ, S^a) be an a.p. compactification of S and (ψ, S^w) a weakly a.p. compactification. Then (ϕ, S^a) and (ψ, S^w) have the following 'universal' property:*

If T is a topological (resp. semitopological) compact Hausdorff semigroup and $\xi : S \rightarrow T$ a continuous homomorphism, then there exists a unique continuous homomorphism $\xi^a : S^a \rightarrow T$ (resp. $\xi^w : S^w \rightarrow T$) such that $\xi = \xi^a \circ \phi$ (resp. $\xi = \xi^w \circ \psi$).

PROOF: Unicity follows from the fact that T is a Hausdorff topological space and that ϕ and ψ have dense images.

We now prove the existence of ξ^w in the case that T is a semitopological compact Hausdorff semigroup (the existence of ξ^a in the case that T is a topological compact Hausdorff semigroup can be proved in a similar way). First, we note that

$$(1) \quad \forall s, t \in S : \psi(s) = \psi(t) \Rightarrow \xi(s) = \xi(t)$$

Indeed: $C(T)$ separates the points of T , so

$$\xi(s) \neq \xi(t) \Rightarrow \exists g \in C(T) : g(\xi(s)) \neq g(\xi(t)).$$

Since $g \circ \xi \in W(S)$ it follows from 1.2 that in this case $\psi(s) \neq \psi(t)$.

Defining $\psi' : S \rightarrow \text{Im } \psi$ by $\psi = i \circ \psi'$ ($i : \text{Im } \psi \rightarrow S^w$ is the inclusion map) it follows from (1) that there is a homomorphism $\xi' : \text{Im } \psi \rightarrow T$ such that $\xi = \xi' \circ \psi$.


Now it is sufficient to show that ξ' is uniformly continuous ($\text{Im } \psi$ provided with the relative uniformity of S^w): T is complete as a uniform space ([3], Ch. 6, Theorem 32), so in this case ξ' has a uniformly con-

tinuous extension $\xi^w : S^w \rightarrow T$ ([3], Ch. 6, Theorem 26). It is easy to see that this continuous extension ξ^w of the homomorphism ξ' is itself a homomorphism, using separate continuity of multiplication in S^w and T .

Uniform continuity of ξ' will follow from 2.1 if we succeed in proving the following assertion: if $g : T \rightarrow [0, 1]$ is any continuous map, then $g \circ \xi' : \text{Im } \psi \rightarrow [0, 1]$ is uniformly continuous.

By 1.1, $g \circ \xi \in W(S)$, so there is a $\bar{g} \in C(S^w)$ such that $g \circ \xi = \bar{g} \circ \psi$, that is

$$g \circ \xi' \circ \psi' = \bar{g} \circ i \circ \psi'$$



Since S is mapped onto $\text{Im } \psi$ by ψ' , it follows that

$$g \circ \xi' = \bar{g} \circ i.$$

Now uniform continuity of $g \circ \xi'$ follows from uniform continuity of i and \bar{g} .

2.3. COROLLARY ([1], Corollary 5.6). *Two a.p. compactifications (weakly a.p. compactifications) of a semitopological semigroup are equivalent.*

PROOF: Let (ϕ_1, T_1) and (ϕ_2, T_2) be weakly a.p. compactifications of S . By 2.2 there are continuous homomorphisms $\phi_1^w : T_2 \rightarrow T_1$ and $\phi_2^w : T_1 \rightarrow T_2$ such that

$$\phi_2^w \circ \phi_1 = \phi_2 \quad \text{and} \quad \phi_1^w \circ \phi_2 = \phi_1.$$

We have to prove, that ϕ_1 (or ϕ_2) is a topological isomorphism. It is easy to see, that

$$(\phi_1^w \circ \phi_2^w) \circ \phi_1 = \phi_1 = I_1 \circ \phi_1$$

so by unicity it follows that $\phi_1^w \circ \phi_2^w = I_1$.

Similarly $\phi_2^w \circ \phi_1^w = I_2$ (I_i denotes the identity mapping of T_i for $i = 1, 2$). From this the result follows.

2.4. COROLLARY ([1], Theorem 5.5). *Let S and S_1 be semitopological semigroups, (ϕ, S') and (ϕ_1, S'_1) a.p. compactifications (resp. w.a.p.*

compactifications) of S and S_1 and $\psi : S \rightarrow S_1$ a continuous homomorphism. Then there is a unique continuous homomorphism $\psi' : S' \rightarrow S'_1$ for which $\psi' \circ \phi = \phi_1 \circ \psi$.

PROOF: Existence: take $\psi' = (\phi_1 \circ \psi)^a$ (respectively $\psi' = (\phi_1 \circ \psi)^w$; notation as in 2.2).

Unicity follows from unicity in 2.2: if $\psi'' : S' \rightarrow S'_1$ satisfies $\psi'' \circ \phi = \phi_1 \circ \psi$, then $\psi'' = (\phi_1 \circ \psi)^a$ (respectively $\psi'' = (\phi_1 \circ \psi)^w$).

REFERENCES

K. DE LEEUW AND I. GLICKSBERG

[1] Applications of almost periodic compactifications, *Acta Math.* 105 (1961), 63–97.

J. F. BERGLUND AND K. H. HOFMANN

[2] Compact Semitopological Semigroups and Weakly Almost Periodic Functions, *Lecture Notes in Mathematics* 42, Springer, Berlin, 1967.

J. L. KELLEY

[3] *General Topology*, Van Nostrand, New York, 1955.

(Oblatum 30–IX–1969)

Free University, Amsterdam.