

COMPOSITIO MATHEMATICA

T. A. CHAPMAN

Homotopic homeomorphisms of infinite-dimensional manifolds

Compositio Mathematica, tome 27, n° 2 (1973), p. 135-140

http://www.numdam.org/item?id=CM_1973__27_2_135_0

© Foundation Compositio Mathematica, 1973, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

HOMOTOPIC HOMEOMORPHISMS OF INFINITE-DIMENSIONAL MANIFOLDS

T. A. Chapman¹

1. Introduction and preliminaries

We use F to represent any Fréchet space which is homeomorphic (\cong) to its own countable infinite product F^ω , and by a *Fréchet manifold* (or *F-manifold*) we mean a paracompact manifold modelled on F . The condition $F \cong F^\omega$ is known to be satisfied by any infinite-dimensional separable Fréchet space (as they are all homeomorphic [1]), by any infinite-dimensional Hilbert or reflexive Banach space [2], and by the space of bounded sequences, l_∞ [3]. More generally there is no known example of an infinite-dimensional Fréchet space for which this condition is not satisfied.

In this note we establish the following two results on homotopic homeomorphisms of F -manifolds. (All our homeomorphisms will be onto.)

THEOREM 1: *Let M be an F -manifold and let $f, g : M \rightarrow M$ be homotopic homeomorphisms. Then f is ambient isotopic to g (i.e. each level of the isotopy is onto).*

We remark that previously this result was known for separable l^2 -manifolds, where l^2 is separable infinite-dimensional real Hilbert space. The first proof was given by Burghelea and Henderson, with an argument that mixed differential and point-set techniques [4]. Later Wong gave another proof of the separable case (and some substantial generalizations) by using some recent results on fiber-preserving homeomorphisms on product bundles, with base space a locally compact polyhedron and fiber a separable l^2 -manifold [11]. We give here a short proof of Theorem 1 that uses pointset techniques and a simple modification of the Alexander trick, [7, p. 321].

THEOREM 2: *Let M, N be F -manifolds and let $f, g : M \rightarrow N$ be homo-*

¹ Supported in part by NSF Grant GP 14429.

topic open embeddings. Then f is isotopic to g , with each level of the isotopy being an open embedding.

(We will call an isotopy for which each level is an open embedding an *open isotopy*.)

Previously this result was not known even for l^2 -manifolds and it answers affirmatively a question that was raised at a problem seminar held in January (1969) at Cornell University.

There is one special definition that we will need. A closed subset K of a space X is said to be a *Z-set* in X provided that given any non-null, homotopically trivial open set U in X , $U \setminus K$ is non-null and homotopically trivial. There are several places we will need the following homeomorphism extension theorem of [6], and we henceforth refer to it as the HET: *Let K_1, K_2 be Z-sets in an F-manifold M and let $h : K_1 \rightarrow K_2$ be a homeomorphism which is homotopic to id_{K_1} (the identity on K_1). There is an invertible ambient isotopy $G : M \times I \rightarrow M$ such that $G_0 = id_M$ and $G_1|_{K_1} = h$.*

(Here $I = [0, 1]$ and an ambient isotopy $H : X \times I \rightarrow X$ is said to be *invertible* provided that $H^* : X \times I \rightarrow X$, defined by $H^*(x, t) = H_t^{-1}(x)$, is continuous.)

2. Proof of Theorem 1

Without loss of generality we may assume that $g = id_M$ and we may replace M by $M \times (0, 1]$. (See [6] for references to papers which show, among other things, that $M \cong M \times [0, 1] = M \times (0, 1] \cong M \times (0, 1)$.) Thus we can reduce the problem to showing that a homeomorphism $f : M \times (0, 1] \rightarrow M \times (0, 1]$ which is homotopic to id is ambient isotopic to id (we suppress the subscript on id when the meaning is clear).

Note that $M \times \{1\}$ is a Z-set in $M \times (0, 1]$ and $f|_{M \times \{1\}}$ is a homeomorphism of $M \times \{1\}$ onto a Z-set in $M \times (0, 1]$ which is homotopic to id . Using the HET we see that f is ambient isotopic to a homeomorphism $f' : M \times (0, 1] \rightarrow M \times (0, 1]$ which satisfies $f'|_{M \times \{1\}} = id$.

Using the Alexander trick let $G : (M \times (0, 1]) \times I \rightarrow M \times (0, 1]$ be defined by

$$G_t(x, s) = \begin{cases} (x, s), & \text{for } t \leq s \\ tf'(x, s/t), & \text{for } s < t \end{cases}$$

(where $t(x, u) = (x, tu)$). Then G is an ambient isotopy satisfying $G_0 = id$ and $G_1 = f'$.

3. Some technical lemmas

Since the proof of Theorem 2 is a bit more involved it will be convenient to describe some of the apparatus needed there.

LEMMA 3.1: *Let $M \subset F$ be a connected F -manifold which is a Z -set in F and let U be an open subset of F containing M . Then there exists a closed embedding $h : M \times I \rightarrow F$ such that $h(x, 0) = x$, for all $x \in M$, $h(M \times I) \subset U$, and $h(M \times \{1\}) = Bd_F(h(M \times I))$ (Bd_F is the topological boundary operator in F).*

PROOF. Let $f : M \rightarrow F$ be an open embedding (which exists by [8]) and let $f' : M \times (-1, 1) \rightarrow F \times (-1, 1)$ be defined by $f'(x, t) = (f(x), t)$. By using motions only in the $(-1, 1)$ - direction we can easily construct a homeomorphism $\alpha : M \times (-1, 1) \rightarrow M \times (-1, 1)$ so that $f' \circ \alpha(M \times [0, 1])$ is closed in $F \times (-1, 1)$, thus $Bd_{F \times (-1, 1)}(f' \circ \alpha(M \times [0, 1])) = f' \circ \alpha(M \times \{0\})$. Let $\beta : F \times (-1, 1) \rightarrow F$ be a homeomorphism and let $\gamma : M \times [0, 1] \rightarrow M \times [0, 1]$ be a homeomorphism which satisfies $\gamma(M \times \{0\}) = M \times \{1\}$. (To see how to construct γ let $\gamma_1 : M \rightarrow M \times [0, 1]$ be a homeomorphism and let $\gamma_2 : [0, 1] \times [0, 1] \rightarrow [0, 1] \times [0, 1]$ be a homeomorphism satisfying $\gamma_2([0, 1] \times \{0\}) = [0, 1] \times \{1\}$. Then let $\gamma(x, t) = (\gamma_1^{-1}(p_1 \circ \gamma_1(x), p_1 \circ \gamma_2(p_2 \circ \gamma_1(x), t)), p_2 \circ \gamma_2(p_2 \circ \gamma_1(x), t))$, where we adopt the convention that p_i represents projection onto the i th factor, for $i = 1, 2$.) Then $\beta \circ f' \circ \alpha \circ \gamma^{-1} : M \times [0, 1] \rightarrow F$ is a closed embedding satisfying $Bd_F(\beta \circ f' \circ \alpha \circ \gamma^{-1}(M \times [0, 1])) = \beta \circ f' \circ \alpha \circ \gamma^{-1}(M \times \{1\})$. Since $\beta \circ f' \circ \alpha \circ \gamma^{-1}(M \times \{0\})$ is a Z -set in F we can use the HET to get a homeomorphism $\delta : F \rightarrow F$ so that $\delta \circ \beta \circ f' \circ \alpha \circ \gamma^{-1}(x, 0) = x$, for all $x \in M$. Put $g = \delta \circ \beta \circ f' \circ \alpha \circ \gamma^{-1}$.

Now $g^{-1}(U)$ is an open subset of $M \times [0, 1]$ containing $M \times \{0\}$, and we can clearly construct a closed embedding $\phi : M \times [0, 1] \rightarrow M \times [0, 1]$ such that $\phi(x, 0) = (x, 0)$, for all $x \in M$, $\phi(M \times [0, 1]) \subset g^{-1}(U)$, and $Bd(\phi(M \times [0, 1])) = \phi(M \times \{1\})$. Then $h = g \circ \phi$ fulfills our requirements.

LEMMA 3.2: *Let U be an open subset of a connected F -manifold M and let $K \subset U$ be a Z -set in U . Then there exists an open embedding $f : U \rightarrow M$ such that $f(K)$ is a Z -set in M and f is openly isotopic to id_U .*

PROOF: We may once more use [8] to assume that M is an open subset of F . Theorem 2.2 of [10] shows that $U \times F \cong U$, and it is clear that the proof given there can be modified slightly to get a homotopy $G : U \times F \times I \rightarrow U$ such that $G_0 = \pi_U$ (projection onto U), $G_t : U \times F \rightarrow U$ is a homeomorphism for all $t \in (0, 1]$, and $\lim_{t \rightarrow 0} d(G_t, \pi_U) = 0$, where $d(G_t, \pi_U) = \sup \{d(G_t(x), \pi_U(x)) \mid x \in U \times F\}$ and d is a suitably chosen

metric for F . Similarly let $H : M \times F \times I \rightarrow M$ be a homotopy such that $H_0 = \pi_M$, $H_t : M \times F \rightarrow M$ is a homeomorphism, for all $t \in (0, 1]$, and $\lim_{t \rightarrow 0} d(H_t, \pi_M) = 0$. Then define $f' = H_1 \circ G_1^{-1}$, which is an open embedding of U in M . We now show that f' can be modified to fulfill our requirements.

Observe that $H_t \circ G_t^{-1}$, $0 < t \leq 1$, gives an isotopy of open embeddings of U in M . It is also clear that $\lim_{t \rightarrow 0} d(H_t \circ G_t^{-1}, \text{id}_U) = 0$. Thus f' is openly isotopic to id_U . Using Theorem 1 of [6] there exists a homeomorphism $\alpha : U \times F \rightarrow U \times F$ so that $\alpha \circ G_1^{-1}(K) \subset U \times \{0\}$. It is also clear from the construction given in [6] that α may be chosen to be homotopic to id , thus by our Theorem 1 ambient isotopic to id . We can use motions only in the F -direction to construct a homeomorphism $\beta : U \times F \rightarrow U \times F$ so that $\beta \circ \alpha \circ G_1^{-1}(K)$ is a Z -set in $M \times F$ and β is ambient isotopic to id . Thus $f = H_1 \circ \beta \circ \alpha \circ G_1^{-1}$ gives an open embedding of U in M which is openly isotopic to f' and for which $f(K)$ is a Z -set.

LEMMA 3.3: *Let M be an F -manifold and let $f, g : M \rightarrow M$ be closed bicollared embeddings of M into itself which are homotopy equivalences. Then there is a homeomorphism $h : M \rightarrow M$ such that $h|f(M) = g \circ f^{-1}$. (A set A in a space X is bicollared provided that there exists an open embedding $\phi : A \times (-1, 1) \rightarrow X$ such that $\phi(a, 0) = a$, for all $a \in A$.)*

PROOF: Since M is an ANR [9, page 3] we can use Lemma 11.3 of [5] to conclude that $M \setminus f(M) = M_1 \cup M_2$ and $M \setminus g(M) = N_1 \cup N_2$, where M_1, M_2 and N_1, N_2 are disjoint pairs of subsets of M such that $C1(M_1) \cong C1(M_2) \cong C1(N_1) \cong C1(N_2)$, $f(M)$ is a strong deformation retract of each of $C1(M_1)$ and $C1(M_2)$, $g(M)$ is a strong deformation retract of each of $C1(N_1)$ and $C1(N_2)$, $f(M)$ is collared in each of $C1(M_1)$ and $C1(M_2)$, and $g(M)$ is collared in each of $C1(N_1)$ and $C1(N_2)$.

To finish the proof we can imitate the proof of Theorem 9 of [5] provided that we use the HET and Theorem 6 of [8], which implies that every homotopy equivalence between F -manifolds is homotopic to a homeomorphism.

4. Proof of Theorem 2

Since each component of N is open we may assume, without loss of generality, that N is connected. We may also replace M by $M \times [0, 1)$. Using Lemma 3.2 we can additionally assume that $f(M \times \{0\})$ and $g(M \times \{0\})$ are Z -sets in N . Thus $f|M \times \{0\}$ and $g|M \times \{0\}$ are homotopic homeomorphisms of $M \times \{0\}$ onto Z -sets in N . We can use the HET to obtain an ambient invertible isotopy $G : N \times I \rightarrow N$ such that $G_0 = \text{id}$

and $G_1 \circ f = g$ (on $M \times \{0\}$). Thus $f' = G_1 \circ f : M \times [0, 1) \rightarrow N$ is an open embedding which satisfies $f'|M \times \{0\} = g|M \times \{0\}$ and f' is openly isotopic to f . Then $(f')^{-1}(g(M \times [0, 1)))$ is an open subset of $M \times [0, 1)$ containing $M \times \{0\}$, and we can find an open embedding $\theta : M \times [0, 1) \rightarrow M \times [0, 1)$ such that $\theta(M \times [0, 1)) \subset (f')^{-1}(g(M \times [0, 1)))$, $\theta|M \times \{0\} = \text{id}$, and θ is openly isotopic to id . Thus $f_1 = f' \circ \theta : M \times [0, 1) \rightarrow g(M \times [0, 1))$ is an open embedding which satisfies $f_1|M \times \{0\} = g|M \times \{0\}$ and for which f_1 is openly isotopic to f .

Using Lemma 3.1 and the fact that N can be embedded as an open subset of F , there is a closed embedding $h : M \times [0, 1] \rightarrow g(M \times [0, 1))$ such that $h(M \times [0, 1]) \subset f_1(M \times [0, 1))$, $h(x, 0) = g(x, 0)$, for all $x \in M$, and $Bd_N(h(M \times [0, 1])) = h(M \times \{1\})$. Using Lemma 11.3 of [5] as applied to our Lemma 3.3, it follows that for $A = g(M \times [0, 1)) \setminus h(M \times [0, \frac{1}{2}))$, $h(M \times \{\frac{1}{2}\})$ is collared in A and $h(M \times \{\frac{1}{2}\})$ is a strong deformation retract of A . Using the HET we can construct a homeomorphism $\alpha : M \times [0, 1) \rightarrow g(M \times [0, 1))$ such that $\alpha|M \times [0, \frac{1}{2}] = h|M \times [0, \frac{1}{2}]$. Note that there is a homeomorphism $\beta : M \times [0, \frac{1}{2}) \rightarrow M \times [0, 1)$ such that $\beta(x, 0) = (x, 0)$, for all $x \in M$ and β is openly isotopic to id .

Then $\alpha \circ \beta \circ \alpha^{-1} : h(M \times [0, \frac{1}{2})) \rightarrow g(M \times [0, 1))$ is a homeomorphism such that $\alpha \circ \beta \circ \alpha^{-1} \circ h(x, 0) = g(x, 0)$, for all $x \in M$, and $\alpha \circ \beta \circ \alpha^{-1}$ is openly isotopic to id . Let $\gamma = \alpha \circ \beta \circ \alpha^{-1}$ and similarly let $\delta : f_1(M \times [0, 1)) \rightarrow h(M \times [0, \frac{1}{2}))$ be a homeomorphism which satisfies $\delta \circ f_1(x, 0) = h(x, 0)$, for all $x \in M$, and δ is openly isotopic to id .

Then $g^{-1} \circ \gamma \circ \delta \circ f_1 : M \times [0, 1) \rightarrow M \times [0, 1)$ is a homeomorphism which satisfies $g^{-1} \circ \gamma \circ \delta \circ f_1(x, 0) = (x, 0)$, for all $x \in M$, and it is therefore homotopic to id . Using Theorem 1 there exists an ambient isotopy $\Phi : (M \times [0, 1)) \times I \rightarrow M \times [0, 1)$ such that $\Phi_0 = g^{-1} \circ \gamma \circ \delta \circ f_1$ and $\Phi_1 = \text{id}$. Then $g \circ \Phi_t : M \times [0, 1) \rightarrow g(M \times [0, 1))$ is an isotopy of open embeddings such that $g \circ \Phi_0 = \gamma \circ \delta \circ f_1$ and $g \circ \Phi_1 = g$. Since γ and δ were constructed to be openly isotopic to id , it follows that f_1 is openly isotopic to g . As f_1 is openly isotopic to f we are done.

REFERENCES

- [1] R. D. ANDERSON, Hilbert space is homeomorphic to the countable infinite product of lines, *Bull. Amer. Math. Soc.*, 72 (1966) 515-519.
- [2] C. BESSAGA and M. I. KADEC, On topological classification of non-separable Banach spaces, (to appear).
- [3] C. BESSAGA and A. PELCZYNSKI, Some remarks on homeomorphisms of Banach spaces, *Bull. Acad. Polon. Sci. Ser. Sci. Math. Astro. Phys.*, 8 (1960) 757-761.
- [4] D. BURGHELEA and D. W. HENDERSON, Smoothings and homeomorphisms for Hilbert manifolds, *Bull. Amer. Math. Soc.*, 76 (1970) 1261-1265.
- [5] T. A. CHAPMAN, On the structure of Hilbert cube manifolds, *Compositio Math.*, 24 (1972) 329-353.

- [6] T. A. CHAPMAN, Deficiency in infinite-dimensional manifolds, *General Topology and its Applications*, 1 (1971) 263–272.
- [7] J. DUGUNDJI, *Topology*, Allyn and Bacon, Inc., Boston, 1966.
- [8] D. W. HENDERSON, *Corrections and extensions of two papers about infinite-dimensional manifolds*, (preprint).
- [9] R. S. PALAIS, *Homotopy theory of infinite-dimensional manifolds*. *Topology*, 5 (1966) 1–16.
- [10] R. M. SCHORI, Topological stability for infinite-dimensional manifolds. *Compositio Math.* (to appear).
- [11] RAYMOND Y. T. WONG, *Parametric extension of homeomorphisms for Hilbert manifolds* (preprint).

(Oblatum 2–III–1971 & 7–V–1973)

Louisiana State University
Baton Rouge, Louisiana
Mathematical Center
Amsterdam, Holland