

COMPOSITIO MATHEMATICA

H. G. MEIJER

R. TIJDEMAN

On additive functions

Compositio Mathematica, tome 29, n° 3 (1974), p. 265-271

http://www.numdam.org/item?id=CM_1974__29_3_265_0

© Foundation Compositio Mathematica, 1974, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON ADDITIVE FUNCTIONS

H. G. Meijer and R. Tijdeman

1. Introduction

A function F defined on the set of positive integers is said to be additive if

$$(1) \quad F(nm) = F(n) + F(m)$$

whenever $(n, m) = 1$; F is called completely additive if (1) holds for every pair of positive integers n and m . Obviously a completely additive function is determined by its values on the set of primes. The function $F(n) = c \log n$ (c constant) is an example of a completely additive function. On the other hand Erdős [1] proved that if an additive function F satisfies one of the following two conditions

$$(2) \quad F(n+1) \geq F(n) \quad \text{for all } n$$

or

$$(3) \quad F(n+1) - F(n) = o(1) \quad \text{as } n \rightarrow \infty$$

then F is necessarily of the form $F(n) = c \log n$ for some constant c . The same results have been found by several other authors.

Pisot and Schoenberg [2] considered the following situation. Let p_1, \dots, p_r denote r different primes, let A be the multiplicative semigroup generated by p_1, \dots, p_r and let F be an additive function defined on A . Pisot and Schoenberg proved that if F is non decreasing on A , then $F(n) = c \log n$ on A provided that $r \geq 3$. (For $r = 2$ monotonic additive functions exist which are not of the form $c \log n$). Obviously this result is a generalisation of the first mentioned result of Erdős. Now one may ask if a condition similar to condition (3) would imply that $F(n) = c \log n$ on A . The present paper deals with this question.

We write $A = \{1 = n_1 < n_2 < n_3 < \dots\}$ and restrict ourselves to

functions F on A which are completely additive. In section 2 we give some results on the structure of A . For the proofs we refer to [3], [4] and [5]. In section 3 we prove that

$$F(n_{i+1}) - F(n_i) = o(1) \quad \text{as } i \rightarrow \infty$$

implies that $F(n) = c \log n$ on A . Subsequently we prove in section 4 that every completely additive function F on A satisfies

$$F(n_{i+1}) - F(n_i) = O(\log(n_{i+1} - n_i)) \quad \text{as } i \rightarrow \infty.$$

We conjecture that the result of section 3 can be improved considerably and that the condition

$$F(n_{i+1}) - F(n_i) = o(\log(n_{i+1} - n_i)) \quad \text{as } i \rightarrow \infty$$

is already sufficient to imply that $F(n) = c \log n$ on A . We are able to prove the conjecture for $r \leq 5$; see section 5.

2. The structure of A

In the sequel we shall use the following notations. By p_1, \dots, p_r we denote r different primes ($r \geq 2$). The multiplicative semigroup generated by them will be denoted by A ; $1 = n_1 < n_2 < n_3 < \dots$ are the elements of A in increasing order.

LEMMA 1: *There exist positive constants C_1, C_2 and N such that*

$$\frac{n_i}{(\log n_i)^{C_1}} < n_{i+1} - n_i < \frac{n_i}{(\log n_i)^{C_2}} \quad \text{for } n_i \geq N.$$

PROOF: The first inequality is the corollary of [3] Theorem 1. The second can be found in [4].

We shall use the following easy consequences of lemma 1:

$$(4) \quad n_{i+1} - n_i \geq \sqrt{n_i} \quad \text{for } n_i \geq N_1,$$

$$(5) \quad n_{i+1} \leq 2n_i \quad \text{for } n_i \geq N_2,$$

$$(6) \quad \frac{\log n_{i+1}}{\log n_i} \rightarrow 1 \quad \text{for } i \rightarrow \infty,$$

$$(7) \quad \frac{n_{i+1}}{n_i} \rightarrow 1 \quad \text{for } i \rightarrow \infty.$$

LEMMA 2: Let p be one of the primes $\{p_1, \dots, p_r\}$. Then there exists an infinite number of pairs n_i, n_{i+1} such that n_i is a pure power of p and n_{i+1} is not divisible by p .

PROOF: See [5] Theorem 2.

LEMMA 3: Let p and q be two different primes from $\{p_1, \dots, p_r\}$. There exist infinitely many pairs n_i, n_{i+1} such that one of the numbers n_i, n_{i+1} is composed of p and q and the other is neither divisible by p nor by q .

PROOF: See [5] Theorem 3.

3

In this section we prove the following theorem.

THEOREM 1: Let F be a completely additive function defined on A satisfying

$$(8) \quad F(n_{i+1}) - F(n_i) = o(1) \quad \text{as } i \rightarrow \infty,$$

then $F(n) = c \log n$ on A for some constant c .

PROOF: Put $F(p_\rho) = c_\rho \log p_\rho$ ($\rho = 1, \dots, r$). Without loss of generality we may assume that the p_1, \dots, p_r are arranged in such a way that $c_1 \geq c_2 \geq \dots \geq c_r$. We shall prove $c_1 = \dots = c_r$, which obviously implies the assertion of the theorem.

Suppose $c_1 = c_2 = \dots = c_s > c_{s+1} \geq \dots \geq c_r$ for some $s \in \{1, \dots, r-1\}$. Let n_i, n_{i+1} be a pair of consecutive elements of A such that n_i is composed of primes p_1, \dots, p_s only and n_{i+1} contains at least one prime from $\{p_{s+1}, \dots, p_r\}$. It is evident that an infinite sequence of such pairs exists.

Put $n_{i+1} = p_1^{k_1} \dots p_s^{k_s} p_{s+1}^{k_{s+1}} \dots p_r^{k_r}$, then

$$F(n_{i+1}) = c_1 k_1 \log p_1 + \dots + c_s k_s \log p_s + c_{s+1} k_{s+1} \log p_{s+1} + \dots + c_r k_r \log p_r = c_1 (k_1 \log p_1 + \dots + k_r \log p_r) - R = c_1 \log n_{i+1} - R,$$

where

$$(9) \quad R \geq (c_s - c_{s+1}) \log 2.$$

Since $F(n_i) = c_1 \log n_i$ we obtain

$$F(n_i) - F(n_{i+1}) = c_1 \log \frac{n_i}{n_{i+1}} + R.$$

In view of (7) and (9) this is a contradiction with (8).

4

Using the estimates of $n_{i+1} - n_i$ we can easily deduce the following theorem.

THEOREM 2: *Let F be a completely additive function on A . Then*

$$F(n_{i+1}) - F(n_i) = O(\log(n_{i+1} - n_i)) \quad \text{as } i \rightarrow \infty.$$

PROOF: Put $F(p_\rho) = c_\rho \log p_\rho$ ($\rho = 1, \dots, r$) and $M = \max_\rho |c_\rho|$. Then we obtain for $n_i = p_1^{t_1} \cdots p_r^{t_r}$ that

$$|F(n_i)| = |t_1 c_1 \log p_1 + \cdots + t_r c_r \log p_r| \leq M \log n_i.$$

Hence by (5),

$$|F(n_{i+1}) - F(n_i)| \leq 2M \log n_{i+1} \leq 2M \log 2n_i \quad \text{for } n_i \geq N_2.$$

Therefore using (4) we obtain

$$\left| \frac{F(n_{i+1}) - F(n_i)}{\log(n_{i+1} - n_i)} \right| \leq \frac{4M \log 2n_i}{\log n_i} = O(1) \quad \text{for } n_i \geq \max(N_1, N_2)$$

from which the theorem follows.

5

For $r \leq 5$ we can improve theorem 1. To that purpose we prove the following theorem.

THEOREM 3: *Let F be a completely additive function on A . Put $F(p_\rho) = c_\rho \log p_\rho$ ($\rho = 1, \dots, r$). We may suppose that p_1, \dots, p_r are arranged in such a way that $c_1 \geq c_2 \geq \cdots \geq c_r$. If*

$$(10) \quad F(n_{i+1}) - F(n_i) = o(\log(n_{i+1} - n_i)) \quad \text{as } i \rightarrow \infty,$$

then it follows that $c_1 = c_2$ if $r = 2$ and $c_1 = c_2 = c_3$ and $c_{r-2} = c_{r-1} = c_r$ if $r \geq 3$.

COROLLARY: *If $r = 2, 3, 4$ or 5 then condition (10) is a sufficient one for a completely additive function on A to imply $F(n) = c \log n$ on A . Obviously this is an improvement of theorem 1.*

PROOF OF THEOREM 3: Suppose $c_1 > c_2$. Choose $\varepsilon > 0$ such that

$$(11) \quad c_2 + \varepsilon < c_1.$$

We obtain from (10) that $F(n_i) \leq F(n_{i+1}) + \varepsilon \log n_{i+1}$, if i is sufficiently large.

By lemma 2 there exists an infinite number of pairs n_i, n_{i+1} such that n_i is a pure power of p_1 and n_{i+1} is not divisible by p_1 . Then $F(n_i) = c_1 \log n_i$ and $c_2 \geq \dots \geq c_r$ implies $F(n_{i+1}) \leq c_2 \log n_{i+1}$. Hence

$$c_1 \log n_i \leq (c_2 + \varepsilon) \log n_{i+1}$$

from which it follows by (6),

$$c_1 \leq c_2 + \varepsilon.$$

This is a contradiction with (11). Thus $c_1 = c_2$. Applying this result to the function $-F$ we obtain $c_r = c_{r-1}$.

Suppose $r \geq 3$ and $c_1 = c_2 > c_3$. Choose $\varepsilon > 0$ such that

$$(12) \quad c_3 + \varepsilon < c_1.$$

We obtain from (10) that

$$(13) \quad F(n_i) \leq F(n_{i+1}) + \varepsilon \log n_{i+1}$$

and

$$(14) \quad F(n_{i+1}) \leq F(n_i) + \varepsilon \log n_{i+1}$$

if i is sufficiently large. By lemma 3 there exist infinitely many pairs n_i, n_{i+1} such that one of them is composed of p_1 and p_2 and the other is composed of p_3, \dots, p_r .

Suppose first that n_i is composed of p_1, p_2 . Then $F(n_i) = c_1 \log n_i$ and since $c_3 \geq \dots \geq c_r$, we have $F(n_{i+1}) \leq c_3 \log n_{i+1}$. Hence, by (13),

$$(15) \quad c_1 \log n_i \leq c_3 \log n_{i+1} + \varepsilon \log n_{i+1} \quad (i \geq i_0).$$

Suppose on the other hand that n_{i+1} is composed of p_1, p_2 . Then

$$F(n_{i+1}) = c_1 \log n_{i+1} \quad \text{and} \quad F(n_i) \leq c_3 \log n_i.$$

Therefore by (14)

$$(16) \quad c_1 \log n_{i+1} \leq c_3 \log n_i + \varepsilon \log n_{i+1} \quad (i \geq i_0).$$

Since there is an infinite number of integers i for which (15) or (16) holds we obtain by (6)

$$c_1 \leq c_3 + \varepsilon.$$

This is a contradiction with (12). Therefore $c_1 = c_2 = c_3$. Applying this result to the function $-F$ we obtain $c_r = c_{r-1} = c_{r-2}$.

REMARK: In [5] the following conjecture is made:

Let p_1, \dots, p_r be different primes and $\{n_i\}$ the increasing sequence of integers composed of these primes. Let t be fixed, $1 \leq t \leq r-1$. Then there exist infinitely many pairs n_i, n_{i+1} such that one of the numbers n_i, n_{i+1} is composed of p_1, \dots, p_t and the other is composed of p_{t+1}, \dots, p_r .

Under the assumption that this conjecture is true, we can prove, for all r , that the condition

$$F(n_{i+1}) - F(n_i) = o(\log(n_{i+1} - n_i)) \quad \text{as } i \rightarrow \infty$$

implies that $F(n) = c \log n$ on A for some constant c .

REFERENCES

- [1] P. ERDÖS: On the distribution function of additive functions. *Ann. of Math.* 47 (1946) 1–20.
- [2] C. PISOT and I. J. SCHOENBERG: Arithmetic problems concerning Cauchy's functional equation. *Illinois J. Math.* 8 (1964) 40–56.
- [3] R. TIJDEMAN: On integers with many small prime factors. *Compositio Math.* 26 (1973) 319–330.
- [4] R. TIJDEMAN: On the maximal distance between integers composed of small primes. *Compositio Math.* 28 (1974) 159–162.
- [5] R. TIJDEMAN and H. G. MEIJER: On integers generated by a finite number of fixed primes. *Compositio Math.* 29 (1974) 273–286.

(Oblatum 19-III-1974)

University of Technology
Department of Mathematics
Delft, Netherlands
University of Leiden,
Mathematical Institute
Leiden, Netherlands