

COMPOSITIO MATHEMATICA

D. R. LEWIS

An isomorphic characterization of the Schmidt class

Compositio Mathematica, tome 30, n° 3 (1975), p. 293-297

http://www.numdam.org/item?id=CM_1975__30_3_293_0

© Foundation Compositio Mathematica, 1975, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

AN ISOMORPHIC CHARACTERIZATION OF THE SCHMIDT CLASS

D. R. Lewis¹

This note gives an isomorphic characterization of the Hilbert-Schmidt norm within the class of unitarily invariant crossnorms on the space R of finite rank operators on l_2 . Specifically if the completion of R under α has an unconditional basis (or more generally local unconditional structure) than α is equivalent to the Hilbert-Schmidt norm.

For convenience only real Banach spaces are considered. Notation and terminology is standard, with the following possible exceptions.

For E finite dimensional the unconditional basis constant of E is denoted by $\chi(E)$, and the parameter η is defined by $\eta(E) = \inf \|u\| \|v\| \chi(F)$, where the infimum is taken over all spaces F and pairs of operators $u : E \rightarrow F, v : F \rightarrow E$ satisfying $vu = \text{identity}$. The definition of η is extended to infinite dimensional spaces by setting $\eta(G) = \inf_{\mathcal{Q}} \sup_{E \in \mathcal{Q}} \eta(E)$, with the infimum taken over all confinal collections of finite dimensional subspaces of G . G has *local unconditional structure* if $\eta(G) < \infty$.

R (respectively, $R(n)$) is the space of finite rank operators on l_2 (resp., l_2^n). A norm α on R (or $R(n)$) is called a *unitarily invariant crossnorm* if, for each $u \in R$, $\|u\| \leq \alpha(u) \leq i_1(u)$ and $\alpha(guh) = \alpha(u)$ for all isometries g and h . This coincides with Schatten's definition [7]. The completion of R under α is written $R(\alpha)$, and $R(\alpha, n)$ is $R(n)$ with norm α . Finally, π_p and i_p denote the p -absolutely summing and p -integral norms, respectively ([6]).

THEOREM: *If α is a unitarily invariant crossnorm on l_2 and if $R(\alpha)$ has local unconditional structure, then α is equivalent to the Hilbert-Schmidt norm.*

PROOF: Fix n and let P (respectively, Q) be the natural inclusion of $R(\alpha, n)$ (resp., $R(\alpha, n)$) into $R(\pi_2, n)$. G denotes the group of isometries of l_2^n and dg is the normalized Haar measure on G . Let $\omega \in R(n)$ with $\omega \neq 0$, and write

¹ The author was supported in part by NSF GP-42666.

$(\omega^*\omega)^{\frac{1}{2}} = \sum_{i \leq n} \lambda_i e_i \otimes e_i$, where (e_i) is some orthonormal basis for l_2^n and each $\lambda_i \geq 0$. The dual of $R(\alpha, n)$ is naturally identified with $R(\alpha', n)$, where α' is the associate crossnorm of α ([7]) and the action is the trace of the composition.

Define a probability measure μ on the closed unit ball of $R(\alpha, n)$ by setting

$$\mu(f) = \int_G \int_G f(\alpha(\omega)^{-1}g\omega h^*)dgdh.$$

Then for $u \in R(\alpha, n)$

$$\begin{aligned} \alpha(\omega)\mu(|\langle u, \cdot \rangle|) &= \iint |\text{trace}(ug(\omega^*\omega)^{\frac{1}{2}}h^*)|dgdh \\ &= \iint \left| \sum_{i \leq n} \lambda_i(h(e_i), ug(e_i)) \right|dgdh, \end{aligned}$$

the first since ω may be translated to $(\omega^*\omega)^{\frac{1}{2}}$ by an isometry. Now in the last integral replace g by gg_ε , where $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$ is an n -tuple of signs and $g_\varepsilon(e_i) = \varepsilon_i e_i$. Averaging over all such n -tuples ε and applying Khinchin's inequality yields

$$\begin{aligned} \alpha(\omega)\mu(|\langle u, \cdot \rangle|) &\geq 3^{-\frac{1}{2}} \iint \left(\sum_{i \leq n} \lambda_i^2 |(h(e_i), ug(e_i))|^2 \right)^{\frac{1}{2}} dgdh \\ &\geq 3^{-\frac{1}{2}} \left[\sum_{i \leq n} \lambda_i^2 \left(\iint |h(e_i), ug(e_i)| dgdh \right)^2 \right]^{\frac{1}{2}} \\ &\geq 4^{-1} \pi_2(\omega) \iint |(h(e), ug(e))| dgdh \end{aligned}$$

for any vector $e \in l_2^n$ with norm one. Let dm be the normalized $(n-1)$ -dimensional rotational invariant measure on the unit sphere of l_2^n and c_n the constant satisfying

$$\|x\| = c_n \int |(x, z)|m(dz), \quad x \in l_2^n.$$

This equality shows that for $\|e\| = 1$

$$\iint |(h(e), ug(e))|dgdh = \int c_n^{-1} \|u(x)\|m(dx)$$

$$\begin{aligned}
 &= c_n^{-1} \int (\sum_{i \leq n} |(x, u^*(e_i))|^2)^{\frac{1}{2}} m(dx) \\
 &\geq c_n^{-1} \left[\sum_{i \leq n} \left(\int |(x, u^*(e_i))| m(dx) \right)^2 \right]^{\frac{1}{2}} \\
 &= c_n^{-2} (\sum_{i \leq n} \|u^*(e_i)\|^2)^{\frac{1}{2}},
 \end{aligned}$$

and the last is nothing but $c_n^{-2} \pi_2(u)$. Combining inequalities now gives

$$\pi_2(u) \leq 4c_n^2 \alpha(\omega) \pi_2(\omega)^{-1} \mu(|\langle u, \cdot \rangle|)$$

so that

$$\pi_1(Q) \leq 4c_n^2 \alpha(\omega) \pi_2(\omega)^{-1}.$$

To estimate $\pi_1(P)$ let γ be the probability measure on the closed unit ball of $R(\alpha, n)$ defined by

$$\gamma(f) = \int_G \int_G f(g(e) \otimes h(e)) dg dh,$$

where e is some fixed vector of norm one. The previous inequality shows

$$\pi_2(u) \leq c_n^2 \gamma(|\langle \cdot, u \rangle|)$$

for all $u \in R(n)$, and consequently $\pi_1(P) \leq c_n^2$. By the proof of Theorem 3.5 of [2]

$$n^2 \leq \eta(R(\alpha, n)) \pi_1(P) \pi_1(Q) \leq \eta(R(\alpha, n)) 4c_n^4 \alpha(\omega) \pi_2(\omega)^{-1}$$

and by [1] $c_n \leq (\pi n/2)^{\frac{1}{2}}$, so

$$\pi_2(\omega) \leq \pi^2 \eta(R(\alpha, n)) \alpha(\omega).$$

This first inequality is also valid if α is replaced by its associate α' , and hence by duality (using $\pi'_2 = \pi_2$)

$$\alpha(\omega) \leq \pi^2 \eta(R(\alpha, n)) \pi_2(\omega).$$

The constants in the last two inequalities are both dominated by $\pi^2 \eta(R(\alpha))$, so that α and π_2 are $\pi^4 \eta(R(\alpha))^2$ -equivalent on R . This completes the proof.

REMARK 1: An absolutely summing operator on a subspace of an $L_1(\mu)$ -space or a quotient of a $C(K)$ -space must factor through an L_1 -space [2]. Thus the same proof as above shows that α is equivalent to the Hilbert-Schmidt norm if $R(\alpha)$ is isomorphic to a subspace of an L_1 -space or to a quotient of a $C(K)$ -space. In addition, by applying a basis selection theorem of Kadec-Pelczynski [3] and the result of McCarthy [5] that $R(c_p) \not\subset L_p$, it can be shown that α is equivalent to π_2 if $R(\alpha) \subset L_p$ for some $p > 2$. It seems a reasonable conjecture that α is equivalent to π_2 whenever $R(\alpha)$ embeds in a space with an unconditional basis.

REMARK 2: Given $\lambda \geq 1$ there is a unitarily invariant crossnorm α on R which is equivalent to π_2 and such that the unconditional basis constant of $R(\alpha)$ is at least λ . To see this fix n and let ϕ_n be the symmetric gauge function ([7], Chapter V) on the space of finitely non-zero scalar sequences defined by

$$\phi_n(x) = \max \left\{ \left(\sum_{k \geq 1} |x_k|^2 \right)^{\frac{1}{2}}, \max_{|\sigma|=n} \sum_{k \in \sigma} |x_k| \right\}.$$

The crossnorm α on R induced by ϕ_n is clearly $n^{\frac{1}{2}}$ -equivalent to π_2 and by the inequalities of the proof $\eta(R(\alpha)) \geq \pi^{-2} n^{\frac{1}{2}}$. Similarly it is possible to produce unitary crossnorms β for which $\eta(R(\beta, n))$ increases very slowly. For instance, the crossnorm β induced by the symmetric gauge function

$$\psi(x) = \max_{\sigma} |\sigma|^{-\frac{1}{2}} \sum_{k \in \sigma} |x_k|$$

is, asymptotically, $(\log n)^{\frac{1}{2}}$ -equivalent to π_2 on l_2^n .

REMARK 3: Suppose $R(\alpha, n)$ has a basis with the property that each permutation of the basis vectors naturally defines an isomorphism of norm at most λ . By [4] $\eta(R(\alpha, n)) \leq 3\lambda$, so α is $9\pi^4 \lambda^2$ -equivalent to π_2 on l_2^n .

REFERENCES

- [1] Y. GORDON: On p -absolutely summing constants of Banach spaces. *Israel J. Math.* 7 (1969) 151–162.
- [2] Y. GORDON and D. R. LEWIS: Absolutely summing operators and local unconditional structures. *Acta Math.* (to appear).
- [3] M. I. KADEC and A. OELCZYNSKI: Bases, lacunary sequences and complemented subspaces in the space L_p . *Studia Math.* 21 (1962) 161–176.
- [4] D. R. LEWIS: A relation between diagonal and unconditional basis constants (to appear)
- [5] Ch. A. MCCARTHY: Cp. *Israel J. Math.* 5 (1967) 249–271.
- [6] A. PLETSCH: *Theorie der operatorenideale*. Jena, 1972.
- [7] R. SCHATTEN: *Norm ideals of completely continuous operators*. Berlin–Göttingen–Heidelberg, 1960.

(Oblatum 11–XI–1974)

Department of Mathematics
University of Florida
Gainesville, Florida 32601 U.S.A.