

COMPOSITIO MATHEMATICA

ROBERT E. DRESSLER

A property of the φ and σ_j functions

Compositio Mathematica, tome 31, n° 2 (1975), p. 115-118

http://www.numdam.org/item?id=CM_1975__31_2_115_0

© Foundation Compositio Mathematica, 1975, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A PROPERTY OF THE φ AND σ_j FUNCTIONS

Robert E. Dressler

1. Introduction

As usual, φ stands for Euler's function and σ_j stands for the sum of the j^{th} powers of the divisors function. The purpose of this note is to answer the following very natural question: If t is a positive integer and f is φ or σ_j , when does t divide $f(n)$ for almost all positive integers n ? We also answer this question for the Jordan totient function, φ_j , a generalization of the φ function.

We will use the well known formulas:

$$(1) \quad \varphi(n) = n \prod_{p|n} \frac{p-1}{p}$$

$$(2) \quad \sigma_j(n) = \prod_{p^e||n} (p^{ej} + p^{(e-1)j} + \dots + p^j + 1).$$

Here $p^e||n$ means $p^e|n$ and $p^{e+1} \nmid n$.

2. Results

Our first theorem concerns the φ function.

THEOREM (1): *For any prime p_0 and any positive integer k we have $p_0^k|\varphi(n)$ for almost all n . That is, the set of integers n for which $p_0^k \nmid \varphi(n)$ has natural density zero.*

PROOF: If $p_0^k \nmid \varphi(n)$ then by (1), no prime divisor p of n satisfies $p \equiv 1$

¹ In the case of φ_j , j arbitrary, and σ_j , j odd, somewhat stronger results than the ones we give may be obtained by much deeper methods, cf. [2, pg. 167] In the case of σ_j , j even, our results are new. In all cases, our methods appear to be much simpler than those of [2].

(mod p_0^k). Now, if N and M satisfy $N > p'_1 p'_2 \cdots p'_M$ where the p'_i ($i = 1, \dots, M$) are the first M primes congruent to 1 (mod p_0^k), then the number of positive integers not exceeding N , none of whose prime divisors is congruent to 1 (mod p_0^k) is

$$\leq 2N \prod_{i=1}^M \frac{p'_i - 1}{p'_i}.$$

If we let N and M vary together to infinity, then we have, by a strong form of Dirichlet's theorem, that

$$\left(2N \prod_{i=1}^M \frac{p'_i - 1}{p'_i} \right) / N \rightarrow 0.$$

This establishes our result.

Since the finite union of sets of natural density zero is a set of natural density zero we may state the following:

COROLLARY (1): *Let t be any positive integer. Then $t|\varphi(n)$ for almost all n .*

It is also worth noting that the φ_j function where

$$\varphi_j(n) = n^j \prod_{p|n} (1 - p^{-j})$$

also satisfies the conclusions of Theorem 1 and Corollary 1. To see this, observe that if $p \equiv 1 \pmod{p_0^k}$ then also $p^j \equiv 1 \pmod{p_0^k}$.

The situation for the σ_j functions is more complicated. We first need the following two lemmas:

LEMMA (1): [3, pg. 58]. *Let $(c, q) = 1$ where q is any integer having primitive roots. The congruence $x^j \equiv c \pmod{q}$ is solvable if and only if*

$$c^{\varphi(q)/(\varphi(q), j)} \equiv 1 \pmod{q}.$$

LEMMA (2): *Given any prime p_0 and r such that $(r, p_0) = 1$ and any positive integer k , then almost all n are such that n is divisible by only the first power of some prime congruent to $r \pmod{p_0^k}$.*

PROOF: Let p'_1, p'_2, \dots, p'_M be the first M primes congruent to $r \pmod{p_0^k}$.

Let N be greater than $(p'_1 p'_2 \cdots p'_M)^2$. Now for any subset

$$\{p'_{i_1}, p'_{i_2}, \dots, p'_{i_T}\} \text{ of } \{p'_1, p'_2, \dots, p'_M\}$$

the number of integers $\leq N$ which are not divisible by any

$$q \in \{p'_1, \dots, p'_M\} \setminus \{p'_{i_1}, \dots, p'_{i_T}\} = \{q_1, \dots, q_{M-T}\}$$

and are divisible by $p_{i_1}^2 \cdots p_{i_T}^2$ is less than

$$2N \cdot \frac{1}{p_{i_1}^2 \cdots p_{i_T}^2} \left(\frac{q_1 - 1}{q_1} \right) \cdots \left(\frac{q_{M-T} - 1}{q_{M-T}} \right).$$

Thus, the number of integers $\leq N$ which are divisible by some p'_i ($i = 1, \dots, M$) only to the first power is greater than

$$\begin{aligned} N - 2N \sum_{\substack{\text{all subsets} \\ \{p'_{i_1}, \dots, p'_{i_T}\} \text{ of } \{p'_1, \dots, p'_M\}}} \frac{1}{p_{i_1}^2 \cdots p_{i_T}^2} \left(\frac{q_1 - 1}{q_1} \right) \cdots \left(\frac{q_{M-T} - 1}{q_{M-T}} \right) \\ = N - 2N \prod_{i=1}^M \left(\frac{1}{p_i^2} + \frac{p_i - 1}{p_i} \right). \end{aligned}$$

If we now let $M, N \rightarrow \infty$ then by a strong form of Dirichlet's theorem we have

$$(N - 2N) \prod_{i=1}^M \left(\frac{1}{p_i^2} + \frac{p_i - 1}{p_i} \right) / N \rightarrow 1.$$

This completes the proof.

THEOREM (2): Let p_0 be an odd prime and let k and j be any positive integers. Then $p_0^k | \sigma_j(n)$ for almost all n if and only if $\varphi(p_0^k) / (\varphi(p_0^k), j)$ is even.

PROOF: Since p_0 is odd, p_0^k has primitive roots. If $\varphi(p_0^k) / (\varphi(p_0^k), j)$ is even then, by Lemma 1, $x^j \equiv -1 \pmod{p_0^k}$ is solvable. Thus we can find an x_0 such that $x_0^j \equiv -1 \pmod{p_0^k}$. If a prime p satisfies $p \equiv x_0 \pmod{p_0^k}$ and if $p | n$, then by (2) we have $p_0^k | \sigma_j(n)$. To complete this half of the proof we apply Lemma 2 with $r = x_0$.

Now, suppose $\varphi(p_0^k) / (\varphi(p_0^k), j)$ is odd. Since $\varphi(p_0^k) = p_0^{k-1}(p_0 - 1)$, it follows that $\varphi(p_0^k) / (\varphi(p_0^k), j)$ is odd, for an odd prime p_0 , if and only if

$$\frac{p_0 - 1}{(p_0 - 1, j)} = \frac{\varphi(p_0)}{(\varphi(p_0), j)}$$

is odd. Thus, by Lemma 1, $x^j \equiv -1 \pmod{p_0}$ is not solvable. Thus for any square-free integer n (since $\sigma_j(n) = \prod_{p|n}(p^j + 1)$) we have $p_0 \nmid \sigma_j(n)$. Since the square-free integers have natural density $6/\pi^2 > 0$ we are done.

In addition, we have

THEOREM (3): *For any positive integers k and j , $2^k | \sigma_j(n)$ for almost all n .*

PROOF: It is known [1] that for any positive integer k , almost all integers n have the property that they are divisible only to the first degree by at least k distinct odd primes. For these integers n it follows, from (2), that $2^k | \sigma_j(n)$ and the proof is complete.

We may now capsulize Theorems 2 and 3 with

THEOREM (4): *Let p_0 be any prime and let k and j be any positive integers. Then $p_0^k | \sigma_j(n)$ for almost all integers n if and only if*

$$\frac{p_0(p_0 - 1)}{(p_0 - 1, j)}$$

is even.

Finally, we state

COROLLARY (2): *Let t and j be any positive integers. Then $t | \sigma_j(n)$ for almost all n if and only if for each prime divisor p of t we have $p(p-1)/(p-1, j)$ is even.*

REFERENCES

- [1] DRESSLER, R. E.: On a Theorem of Niven, *Can. Math. Bull.*, Vol. 17 (1974) 109–110.
- [2] HARDY, G. H.: *Ramanujan-Twelve Lectures on Subjects Suggested by His Life and Work*. Chelsea (1940).
- [3] LEVEQUE, W. J.: *Topics in Number Theory*, Vol. I. Addison-Wesley (1958).

(Oblatum 22–XI–1974)

Kansas State University
Manhattan, Kansas 66506