

COMPOSITIO MATHEMATICA

M. S. RAMANUJAN

B. ROSENBERGER

On $\lambda(\phi, P)$ -nuclearity

Compositio Mathematica, tome 34, n° 2 (1977), p. 113-125

http://www.numdam.org/item?id=CM_1977__34_2_113_0

© Foundation Compositio Mathematica, 1977, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON $\lambda(\phi, P)$ -NUCLEARITY

M. S. Ramanujan* and B. Rosenberger

Introduction

We consider sequence spaces $\lambda(\phi, P, \infty)$ and $\lambda(\phi, P, k)$ and define $\lambda(\phi, P, \infty)$ -nuclear resp. $\lambda(\phi, P, \mathbb{N})$ -nuclear spaces in order to unify the concepts of $\lambda(P)$ -nuclearity, $\Lambda_n(\alpha)$ -nuclearity, and ϕ -nuclearity considered in [7, 8, 9]; we are especially interested in obtaining universal generators for the varieties of $\lambda(\phi, P, \infty)$ -nuclear and $\lambda(\phi, P, \mathbb{N})$ -nuclear spaces. Section 1 of the paper contains various definitions and some remarks on the operator ideal of $\lambda(P, \infty)$ -nuclear maps, P a stable, nuclear G_∞ -set; in section 2 we consider $\lambda(\phi, P, \infty)$ -nuclear spaces and show that $\lambda(\phi, P, \infty)$ -nuclearity is the same as $\lambda(Q, \infty)$ -nuclearity, Q a suitably chosen G_∞ -set. In section 3 we introduce the concept of $\lambda(\phi, P, \mathbb{N})$ -nuclearity and extend a result in [8] by showing that $\lambda(Q, \infty) - Q$ suitably chosen G_∞ -set – is a universal generator for the variety of $\lambda(\phi, P, \mathbb{N})$ -nuclear spaces whenever P is a countable, monotone, stable, nuclear G_∞ -set.

1. Definitions, notations, and some remarks on $\lambda(P, \infty)$ -nuclear maps

For terminology and notations not explained here we refer to Köthe [3], Pietsch [4], Dubinsky and Ramanujan [1], and Terzioglu [12].

Let X and Y be Banach spaces, λ a normal sequence space, and λ^\times

* The first author acknowledges with pleasure the support of this work under SFB 72 at the University of Bonn and the hospitality of Professor E. Schock.

its Köthe-dual. A continuous linear map $T \in L(X, Y)$ is said to be

(i) λ -nuclear (written $T \in N_\lambda(X, Y)$) if there exists a representation

$$Tx = \sum_{n=0}^{\infty} \gamma_n \langle x, a_n \rangle y_n \quad \text{for } x \in X$$

with $\{\gamma_n\}_n \in \lambda$, $a_n \in X'$, $\|a_n\| \leq 1$, and $y_n \in Y$, $\{\langle y_n, b \rangle\}_n \in \lambda^\times$ for each $b \in Y'$;

(ii) pseudo- λ -nuclear or $\tilde{\lambda}$ -nuclear (written $T \in \tilde{N}_\lambda(X, Y)$) if T has a representation

$$Tx = \sum_{n=0}^{\infty} \gamma_n \langle x, a_n \rangle y_n \quad \text{for } x \in X$$

with $\{\gamma_n\}_n \in \lambda$, $a_n \in X'$, $\|a_n\| \leq 1$, $y_n \in Y$, and $\|y_n\| \leq 1$;

(iii) of type λ if $\{s_n^{\text{app}}(T)\}_n \in \lambda$ where $s_n^{\text{app}}(T)$ denotes the n -th approximation number of T .

For a locally convex Hausdorff space (l.c.s.) E , $\mathcal{U}(E)$ will denote a neighbourhood base of 0 of absolutely convex, closed sets; E_U will denote the completion of the normed space $E/p_U^{-1}(0)$ $U \in \mathcal{U}(E)$; $\delta_n(V, U)$ denotes the n -th Kolmogorov-diameter of $V \in \mathcal{U}(E)$ with respect to $U \in \mathcal{U}(E)$; $\Delta(E)$ denotes the Δ -diametral dimension of E , viz., the sequences $\{\gamma_n\}_n$ such that given $U \in \mathcal{U}(E)$ there exists a $V \in \mathcal{U}(E)$ with $\gamma_n \delta_n(V, U) \rightarrow 0$.

Let $\lambda(P)$ be a Köthe space with its generating Köthe set P . The Köthe set P is called a *power set of infinite type* if it satisfies the following additional conditions:

(i) for each $a \in P$, $0 < a_n \leq a_{n+1}$, $n \in \mathbb{N}$;

(ii) for each $a \in P$, there exists a $b \in P$ with $a_n^2 \leq b_n$, $n \in \mathbb{N}$.

The corresponding space $\lambda(P, \infty)$ is called a *smooth sequence space of infinite type* or a G_∞ -space. For an example of a G_∞ -space which is not a power series space of infinite type see [1; theorem 2.25].

Throughout, $\lambda(P, \infty)$ is assumed to be a G_∞ -space. The nuclearity and related concepts of such spaces are discussed in [1, 12, 13]; we only need the following result.

1.1 LEMMA: $\lambda(P, \infty)$ is nuclear if and only if there exists a sequence $\{p_n\}_n \in P$ such that $\{1/p_n\}_n \in l_1$.

We shall frequently say “ P is a nuclear G_∞ -set” to mean that the corresponding $\lambda(P, \infty)$ is a nuclear G_∞ -space; P is said to be a countable, monotone, nuclear G_∞ -set if $P = \{p_n^i\}_n$: $i = 1, 2, \dots$ with

$p_n^i \leq p_n^{i+1}$ for each $i, n \in \mathbb{N}$ and P is a nuclear G_∞ -set; note that P is a G_∞ -set already implies $0 < p_n^i \leq p_{n+1}^i$.

Let $\lambda(P, \infty)$ be nuclear; then a l.c.s. E is said to be $\lambda(P, \infty)$ -nuclear if for each $U \in \mathcal{U}(E)$ there exists a $V \in \mathcal{U}(E)$ such that V is absorbed by U and the canonical map $K(V, U)$ on E_V to E_U is a $\lambda(P, \infty)$ -nuclear map. In [1] it is shown that a l.c.s. E is $\lambda(P, \infty)$ -nuclear if and only if for each $U \in \mathcal{U}(E)$ there exists $V \in \mathcal{U}(E)$ such that $\{\delta_n(V, U)\}_n \in \lambda(P, \infty)$. We shall denote the class of all $\lambda(P, \infty)$ -nuclear spaces by $\mathcal{N}_{\lambda(P, \infty)}$.

A l.c.s. E is said to be *stable* if $E \times E$ is isomorphic to E . It is known that a nuclear G_∞ -space is stable if and only if for each $p \in P$ there exists a $q \in P$ such that $\{p_{2n}/q_n\}_n \in l_\infty$ [14]. We say “ P is a stable G_∞ -set” to mean that $\lambda(P, \infty)$ is a stable G_∞ -space. Stability plays an important role in the study of various permanence properties of $\lambda(P, \infty)$ -nuclear spaces; the following result can be found in [7; Proposition 4.5].

1.2 PROPOSITION: *Let P be a countable, monotone, nuclear G_∞ -set, $P = \{p_n^i : i = 1, 2, \dots\}$. Then the following statements are equivalent:*

- (i) *For each $j \in \mathbb{N}$ there exists a $\gamma(j) \in \mathbb{N}$ such that $\{p_{2n}^j/p_n^{\gamma(j)}\}_n \in l_\infty$.*
- (ii) *If $\xi^k \in \lambda(P, \infty)$ for each $k \in \mathbb{N}$ and $\beta: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ is a bijection defined by $\beta^{-1}(k, m) = 2^{k-1}(2m - 1)$ then there exist $t_k > 0, k \in \mathbb{N}$, such that the sequence $\{t_{\beta_1(n)} \xi_{\beta_2(n)}^{\beta_1(n)}\}_n \in \lambda(P, \infty)$ where $\beta(n) = (\beta_1(n), \beta_2(n))$.*
- (iii) *If $\xi, \eta \in \lambda(P, \infty)$ and $\zeta = \xi * \eta = (\xi_1, \eta_1, \xi_2, \eta_2, \dots)$ then there exists a bijection $\pi: \mathbb{N} \rightarrow \mathbb{N}$ such that $\{\xi_{\pi(n)}\}_n \in \lambda(P, \infty)$.*
- (iv) *$\mathcal{N}_{\lambda(P, \infty)}$ is closed under the operation of forming countable direct sums.*
- (v) *$\mathcal{N}_{\lambda(P, \infty)}$ is closed under the operation of forming finite Cartesian products.*
- (vi) *$\mathcal{N}_{\lambda(P, \infty)}$ is closed under the operation of forming arbitrary Cartesian products.*
- (vii) *The sum of two $\lambda(P, \infty)$ -nuclear maps is a $\lambda(P, \infty)$ -nuclear map.*
- (viii) *$\lambda(P, \infty)$ is stable.*

An operator ideal A is said to be

- (i) *surjective* if for each closed subspace N of a Banach space X and each $T \in L(X/N, Y)$, Y a Banach space, $TQ_N^X \in A(X, Y)$ implies $T \in A(X/N, Y)$, $Q_N^X: X \rightarrow X/N$ denotes the canonical map onto X/N ;

(ii) *injective* if for each closed subspace M of a Banach space Y and each $T \in L(X, M)$, X a Banach space, $J_M^X T \in A(X, Y)$ implies $T \in A(X, Y)$, $J_M^X: M \rightarrow Y$ denotes the injection.

The following result shows that the operator ideal of $\lambda(P, \infty)$ -nuclear maps – P a stable, countable, monotone, nuclear G_∞ -set – is injective and surjective.

1.3 PROPOSITION: *Let P be a stable, countable, monotone, nuclear G_∞ -set, X and Y be Banach spaces with closed unit balls U_X and U_Y . Then the following statements are equivalent.*

- (i) $T \in L(X, Y)$ is $\lambda(P, \infty)$ -nuclear.
- (ii) $T \in L(X, Y)$ is of type $\lambda(P, \infty)$.
- (iii) $\{s_n^{\text{gel}}(T)\}_n \in \lambda(P, \infty)$ where $s_n^{\text{gel}}(T) := \inf \{\|TJ_M^X\| : \text{codim } M < n, M \text{ closed subspace of } X\}$ denotes the n -th Gelfand-number.
- (iv) $\{s_n^{\text{kol}}(T)\}_n \in \lambda(P, \infty)$ where $s_n^{\text{kol}}(T) := \delta_n(TU_X, U_Y)$.

PROOF: (i) \Leftrightarrow (ii) is shown in [1] and [7]. (ii) \Rightarrow (iii) and (ii) \Rightarrow (iv) are consequences of the facts that $s_n^{\text{gel}}(T) \leq s_n^{\text{app}}(T)$ and $s_n^{\text{kol}}(T) \leq s_n^{\text{app}}(T)$ for each $n \in \mathbb{N}$ [5]. (iii) \Rightarrow (ii) and (iv) \Rightarrow (ii) are consequences of the facts that $s_n^{\text{app}}(T) \leq (n+1) s_n^{\text{gel}}(T)$ and $s_n^{\text{app}}(T) \leq (n+1) s_n^{\text{kol}}(T)$ for each $n \in \mathbb{N}$ [5] and the well known fact that $\{(n+1)\xi_n\}_n \in \lambda(P, \infty)$ for $\{\xi_n\}_n \in \lambda(P, \infty)$.

1.4 COROLLARY: *Let P be a stable, countable, monotone, nuclear G_∞ -set. Then the operator ideal of $\lambda(P, \infty)$ -nuclear maps is injective and surjective.*

PROOF: Let X and Y be Banach spaces, M a closed subspace of Y , and X/N a quotient space of X . Then $s_n^{\text{gel}}(T) = s_n^{\text{gel}}(J_M^Y T)$ for all $T \in L(X, M)$ and $s_n^{\text{kol}}(SQ_N^X) = s_n^{\text{kol}}(S)$ for all $S \in L(X/N, Y)$ [5]. Now the proof easily follows from Proposition 1.3.

1.5 COROLLARY: *Let P be a stable, countable, monotone, nuclear G_∞ -set, X and Y be Banach spaces such that each map $T \in L(X, Y)$ is $\lambda(P, \infty)$ -nuclear. Then X or Y must be finite-dimensional.*

PROOF: By Proposition 1.3 every operator $T \in L(X, Y)$ is of type $\lambda(P, \infty)$, hence of type l_1 . By a result of Pietsch [6], X or Y must be finite-dimensional.

REMARK: With methods used in [11], it can be also shown that X or Y is finite-dimensional if each nuclear map $T \in L(X, Y)$ is $\lambda(P, \infty)$ -nuclear, P as in Corollary 1.5.

2. On $\lambda(\phi, P, \infty)$ -Nuclearity

Throughout, let Φ denote the set of all functions $\phi: [0, \infty] \rightarrow [0, \infty]$ which are continuous, strictly increasing, subadditive with $\phi(0) = 0$ and $\phi(1) = 1$, and satisfy the additional condition (+) there exist constants $M \geq 1$ and $t_\phi \in [0, \infty]$ such that $\phi(\sqrt{t}) \leq \sqrt{\phi(Mt)}$ for $t \in [0, t_\phi]$.

Note that all examples given in [10;2.6] do have the additional property (+). So far we have been unable to find an example of a continuous, strictly increasing, subadditive function $\phi: [0, \infty] \rightarrow [0, \infty]$ with $\phi(0) = 0$ which does not fulfill condition (+).

2.1 DEFINITION: Let $\lambda(P, \infty)$ be nuclear. For $\phi \in \Phi$ define the sequence space $\lambda(\phi, P, \infty)$ by

$$\lambda(\phi, P, \infty) = \{ \{ \xi_n \}_n : \{ \phi(|\xi_n|) \}_n \in \lambda(P, \infty) \}.$$

A l.c.s. E is called $\lambda(\phi, P, \infty)$ -nuclear if for each $U \in \mathcal{U}(E)$ there exists a $V \in \mathcal{U}(E)$ such that $\{ \delta_n(V, U) \}_n \in \lambda(\phi, P, \infty)$.

For $\phi = \text{id}$ this definition agrees with the definition of a $\lambda(P, \infty)$ -nuclear space, so we write $\lambda(P, \infty)$ -nuclear instead of $\lambda(\text{id}, P, \infty)$ -nuclear.

We now show that the concept of $\lambda(\phi, P, \infty)$ -nuclearity is exactly the same as the concept of $\lambda(Q, \infty)$ -nuclearity where Q is suitably chosen depending on ϕ .

2.2 LEMMA: Let P a countable, monotone, nuclear G_∞ -set, $P = \{ \{ p_n^i \}_n : i = 1, 2, \dots \}$. Take $\phi \in \Phi$; define q_n^i by $1/q_n^i := \phi^{-1}(1/p_n^i)$, $i \in \mathbb{N}$. Then $Q := \{ \{ q_n^i \}_n : i = 1, 2, \dots \}$ is a countable, monotone, nuclear G_∞ -set.

PROOF: (i) It is obvious that Q is countable and monotone and that for each n , $i \in \mathbb{N}$ $q_n^i > 0$ and $q_n^i \leq q_{n+1}^i$.

(ii) Given $q^k, q^l \in Q$, we have to show the existence of a $q^r \in Q$ such that $q^k q^l < q^r$, i.e. $q_n^k q_n^l \leq M q_n^r$ for a constant $M > 0$ and each $n \in \mathbb{N}$. Since P is a Köthe set we find $p^m \in P$ and $M_1 > 0$ such that $p_n^k \leq M_1 p_n^m$ and $p_n^l \leq M_1 p_n^m$ for each $n \in \mathbb{N}$. Since $\lambda(P, \infty)$ is nuclear,

there exists a $p^s \in P$ with $\{p_n^m/p_n^s\}_n \in l_1$. This can be seen as follows. By Lemma 1.1 there exists a $p^j \in P$ with $\{1/p_n^j\}_n \in l_1$; we choose $p^h \in P$ with $p_n^j \leq M_2 p_n^h$ and $p_n^m \leq M_2 p_n^h$ for $n \in \mathbb{N}$, we also choose $p^s \in P$ with $(p_n^h)^2 \leq p_n^s$ for all $n \in \mathbb{N}$. (This can be done by condition (ii) of a G_∞ -set.) From the inequalities

$$p_n^m/p_n^s \leq p_n^m/(p_n^h)^2 \leq M_2/p_n^h \leq M_2^2/p_n^j$$

we get $\{p_n^m/p_n^s\}_n \in l_1$. We also find an integer n_0 such that $p_n^k/p_n^s \leq 1$, $p_n^l/p_n^s \leq 1$, and $1/p_n^s < t_\phi$ for each $n \geq n_0$ (here t_ϕ is determined by condition (+)). Because of condition (+) for the function ϕ we then have

$$M_\phi^{-1}(1/p_n^k)\phi^{-1}(1/p_n^l) \geq M\phi^{-1}(1/p_n^s)\phi^{-1}(1/p_n^s) \geq \phi^{-1}(1/p_n^s p_n^s)$$

for each $n \geq n_0$. Again because of the nuclearity of $\lambda(P, \infty)$ and the second additional condition for a G_∞ -set we find $p^r \in P$, $n_1 \in \mathbb{N}$, $n_1 \geq n_0$, such that $p_n^s p_n^s \leq p_n^r$ for each $n \geq n_1$. Therefore there exists $K > 0$ such that $\phi^{-1}(1/p_n^r) \leq K\phi^{-1}(1/p_n^k)\phi^{-1}(1/p_n^l)$ for each $n \in \mathbb{N}$, i.e. $q^k q^l < q^r$.

(iii) To prove the nuclearity of $\lambda(Q, \infty)$ we use Lemma 1.1 and show the existence of a sequence $\{q_n^k\}_n \in Q$ such that $\{1/q_n^k\}_n \in l_1$. Since $\lambda(P, \infty)$ is nuclear we can find $p^k \in P$ such that $\{1/p_n^k\}_n \in l_1$. Since ϕ is subadditive we have $1/q_n^k = \phi^{-1}(1/p_n^k) \leq M/p_n^k$ for each $n \in \mathbb{N}$, and $\lambda(Q, \infty)$ is nuclear.

If $\lambda(P, \infty)$ is a power series space of infinite type then $\lambda(Q, \infty)$ is in general not a power series space as the following example shows.

EXAMPLE: Define $p^i \in P$ by $p_n^i := n^i$, $n, i \in \mathbb{N}$. Let $\phi \in \Phi$ be the function ϕ_{\log} as given in [10], i.e.

$$\phi_{\log}(t) := \begin{cases} 0 & \text{for } t = 0 \\ -\alpha/\log t & \text{for } t \in (0, t_0], t_0 \text{ sufficiently small} \\ \beta t + \gamma & \text{for } t \geq t_0 \end{cases}$$

where $\alpha, \beta, \gamma \geq 0$ are chosen in such a way that ϕ_{\log} is continuous with $\phi(1) = 1$. Then $\phi_{\log} \in \Phi$ and $\phi_{\log}^{-1}(t) = \exp(-\alpha/t)$ for $t \in (0, t_0]$. Therefore $q_n^i = \exp(\alpha n^i)$ for $n \geq n_0, i \in \mathbb{N}$, and $\lambda(Q, \infty)$ is not a power series space [1].

The following lemma is well known and can be found in [4].

2.3 LEMMA: Let P be a countable, monotone, nuclear G_∞ -set. Then $x = \{\xi_n\}_n \in \lambda(P, \infty)$ if and only if $\{\xi_n p_n^i\}_n \in l_\infty$ for each $i \in \mathbb{N}$. The

topologies determined by the semi-norms $\pi_i(x) := \sum_n |\xi_n| p_n^i$ resp. $\sigma_i(x) := \sup |\xi_n| p_n^i$, $i \in \mathbb{N}$, are equivalent.

2.4 PROPOSITION: *Let P be a countable, nuclear G_∞ -set, $\phi \in \Phi$. Then the following statements are true.*

- (i) $\lambda(\phi, P, \infty) = \lambda(Q, \infty)$
- (ii) *On $\lambda(\phi, P, \infty)$ the two topologies given by the F -norms $\tilde{\pi}_i(x) := \sum_n \phi(|\xi_n|) p_n^i$ resp. $\tilde{\sigma}_i(x) := \sup \phi(|\xi_n|) p_n^i$, $i \in \mathbb{N}$, are equivalent.*
- (iii) $\lambda(\phi, P, \infty)$ with the above topologies is topologically equivalent to $\lambda(Q, \infty)$.

PROOF: (ii) For $x = \{\xi_n\}_n \in \lambda(\phi, P, \infty)$ we always have $\tilde{\sigma}_i(x) \leq \tilde{\pi}_i(x)$, $i \in \mathbb{N}$. On the other hand for $i \in \mathbb{N}$ there exists $p^r \in P$ and $p^l \in P$ such that $\{1/p_n^r\} \in l_1$ and $p^i p^r < p^l$; therefore

$$\begin{aligned} \tilde{\pi}_i(x) &= \sum_n \phi(|\xi_n|) p_n^i = \sum_n \phi(|\xi_n|) p_n^i p_n^r / p_n^r \leq M \sum_n \phi(|\xi_n|) p_n^l / p_n^r \\ &\leq M_0 \tilde{\sigma}_i(x) \quad \text{for } x \in \lambda(\phi, P, \infty). \end{aligned}$$

To prove (i) and (iii) notice that because of Lemma 2.3 $x = \{\xi_n\}_n \in \lambda(\phi, P, \infty) \Leftrightarrow \phi(|\xi_n|) p_n^k \leq 1$ for $k \in \mathbb{N}$ and $n \geq n_k \Leftrightarrow |\xi_n| \leq \phi^{-1}(1/p_n^k) = 1/q_n^k$ for $k \in \mathbb{N}$ and $n \geq n_k \Leftrightarrow x \in \lambda(Q, \infty)$.

2.5 COROLLARY: *Let P be a countable, nuclear G_∞ -set. $\phi \in \Phi$. Then $[\lambda(Q, \infty)]'_b = [\lambda(\phi, P, \infty)]'_b$.*

As an immediate consequence of Proposition 4.6 in [7] we finally get

2.6 PROPOSITION: *Let P be a stable, countable, monotone, nuclear G_∞ -set; $\phi \in \Phi$. Let E be a l.c.s. Then the following statements are equivalent.*

- (i) E is $\lambda(\phi, P, \infty)$ -nuclear.
- (ii) E is $\lambda(Q, \infty)$ -nuclear.
- (iii) E is isomorphic to a subspace of a suitable I -fold product $[\lambda(Q, \infty)]'_b$.

3. On $\lambda(\phi, P, \mathbb{N})$ -nuclearity

In this section we study the concepts of $\lambda(\phi, P, \mathbb{N})$ -nuclearity, $\phi \in \Phi$, and $\lambda(P, \mathbb{N})$ -nuclearity, P a countable, monotone G_∞ -set. As in

the case of $\lambda(\phi, P, \infty)$ -nuclearity we obtain that both concepts are closely related. The main result in this section is that whenever P is a countable, monotone, stable, nuclear G_∞ -set $\lambda(P, \infty)$ is a universal generator for the variety of $\lambda(P, \mathbb{N})$ -nuclear spaces. For $\lambda(P, \infty) = \Lambda_\infty(\alpha)$, α a stable exponent sequence, this result has been established in [8].

Let $P = \{\{p_n^i\}_n : i = 1, 2, 3, \dots\}$ be a countable, monotone G_∞ -set. Fix $k \in \mathbb{N}$. We define the sequence space $\lambda(P, k)$ by

$$\lambda(P, k) := \{\{\xi_n\}_n : \sum_n |\xi_n| p_n^k < \infty\}.$$

It is obvious that $\lambda(P, k+1) \subset \lambda(P, k)$ and $\lambda(P, \infty) = \bigcap_k \lambda(P, k)$. For $\phi \in \Phi$ define $\lambda(\phi, P, k)$ by $\lambda(\phi, P, k) := \{\{\xi_n\}_n : \{\phi(|\xi_n|)\}_n \in \lambda(P, k)\}$.

3.1 DEFINITION: Let P be a countable, monotone, nuclear G_∞ -set, $\phi \in \Phi$. A l.c.s. E is said to be

- (i) $\tilde{\lambda}(\phi, P, k)$ -nuclear if for each $U \in \mathcal{U}(E)$ there exists a $V \in \mathcal{U}(E)$ such that $\{\delta_n(V, U)\}_n \in \lambda(\phi, P, k)$;
- (ii) $\lambda(\phi, P, \mathbb{N})$ -nuclear if E is $\tilde{\lambda}(\phi, P, k)$ -nuclear for each $k \geq k_0$, where $\{1/p_n^{k_0}\}_n \in l_1$ because of the nuclearity of $\lambda(P, \infty)$.

For $\phi = \text{id}$ we write $\tilde{\lambda}(P, k)$ -nuclear resp. $\lambda(P, \mathbb{N})$ -nuclear instead of $\tilde{\lambda}(\text{id}, P, K)$ -nuclear resp. $\lambda(\text{id}, P, \mathbb{N})$ -nuclear.

If α is an exponent sequence and if $P = \{\{k^{\alpha_n}\}_n : k \in \mathbb{N}\}$, then a l.c.s. E is $\lambda(P, \mathbb{N})$ -nuclear iff E is $\Lambda_\infty(\alpha)$ -nuclear, i.e. $\tilde{\Lambda}_k(\alpha)$ -nuclear for each $k > 1$. So we indeed generalize the concept of $\Lambda_\infty(\alpha)$ -nuclearity considered in [8].

As in section 2 we now show that the concepts of $\lambda(\phi, P, \mathbb{N})$ -nuclearity and $\lambda(Q, \mathbb{N})$ -nuclearity are the same if Q is suitably chosen.

3.2 PROPOSITION: Let P be a countable, monotone, nuclear G_∞ -set, $\phi \in \Phi$. Define $\{q_n^i\}_n$ by $1/q_n^i := \phi^{-1}(1/p_n^i)$ and $Q := \{\{q_n^i\}_n\}$. Let E be a l.c.s. Then the following statements are equivalent.

- (i) E is $\lambda(\phi, P, \mathbb{N})$ -nuclear.
- (ii) E is $\lambda(Q, \mathbb{N})$ -nuclear.

PROOF: Note that E is $\lambda(\phi, P, \mathbb{N})$ -nuclear (resp. $\lambda(Q, \mathbb{N})$ -nuclear) will follow if E is shown to be $\tilde{\lambda}(\phi, P, k)$ -nuclear (resp. $\tilde{\lambda}(Q, k)$ -nuclear) for each $k \geq k_0$, k_0 as in 3.1. Therefore we will show (1) given $k \geq k_0$ there exists $l \in \mathbb{N}$ such that $\lambda(\phi, P, l+1) \subset \lambda(Q, k)$; (2) given $r \geq r_0$ there exists $s \in \mathbb{N}$ such that $\lambda(Q, s+1) \subset \lambda(\phi, P, r)$. Given k , by nuclearity of $\lambda(Q, \infty)$ (Lemma 2.2) there exists $l \in \mathbb{N}$ such that

$\{q_n^k/q_n^l\}_n \in l_1$. If $x = \{\xi_n\}_n \in \lambda(\phi, P, l + 1)$ then $\phi(|\xi_n|)p_n^l \leq 1$ for $n \geq n_l$ and therefore $\sum_n |\xi_n|q_n^k = \sum_n |\xi_n|q_n^k q_n^l/q_n^l < \infty$. On the other hand given r , by nuclearity of $\lambda(P, \infty)$, find $s \in \mathbb{N}$ such that $\{p_n^r/p_n^s\}_n \in l_1$. So if $x = \{\xi_n\}_n \in \lambda(Q, s + 1)$ we have $|\xi_n| \leq 1/q_n^s$ for $n \geq n_s$ and therefore $\sum_n \phi(|\xi_n|)p_n^r = \sum_n \phi(|\xi_n|)p_n^s p_n^r/p_n^s < \infty$; this shows (1) and (2).

To prove (i) \Rightarrow (ii) fix $k \geq k_0$, find $l \in \mathbb{N}$ such that $\lambda(\phi, P, l + 1) \subset \lambda(Q, k)$. Since E is $\lambda(\phi, P, \mathbb{N})$ -nuclear, E is $\tilde{\lambda}(\phi, P, l + 1)$ -nuclear and therefore $\lambda(Q, \mathbb{N})$ -nuclear. The implication (ii) \Rightarrow (i) can be proved in the same way.

REMARK: It is easily seen that for $\phi \in \Phi$ $\lambda(\phi, P, \mathbb{N})$ -nuclearity implies ϕ -nuclearity [9]. But the reverse of this implication is not true in general as the following example shows.

Take $\phi = \phi_{\log}$; consider the power series space $\Lambda_\infty(\alpha)$ with $\alpha_n := (n + 1)^2$. We show that $\Lambda_\infty(\alpha)$ is ϕ_{\log} -nuclear but not $\lambda(\phi_{\log}, \mathbb{R}, \mathbb{N})$ -nuclear if $R := \{(n + 1)^k\}_n : k = 1, 2, \dots\}$. The ϕ_{\log} -nuclearity of $\Lambda_\infty(\alpha)$ immediately follows from Korollar 3.6 in [9]. But $\Lambda_\infty(\alpha)$ is $\lambda(\phi_{\log}, \mathbb{R}, \mathbb{N})$ -nuclear if and only if there exists $M > 1$ such that $\{\phi_{\log}(1/M^{\alpha_n})\}_n \in \lambda(R, k)$ for each k [8]. Therefore $\Lambda_\infty(\alpha)$ is $\lambda(\phi_{\log}, \mathbb{R}, \mathbb{N})$ -nuclear if and only if for each $k \geq k_0$ the sum $\sum_n (n + 1)^k/\alpha_n \log M = \sum_n (n + 1)^{k-2}/\log M$ is finite which obviously is not true.

In order to answer the question what the model of a universal $\lambda(\phi, P, \mathbb{N})$ -nuclear space is, it is enough to describe the model of a universal $\lambda(Q, \mathbb{N})$ -nuclear space; so from now on P is always supposed to be a countable, monotone, nuclear G -set.

Since $\lambda(P, \mathbb{N})$ -nuclearity implies nuclearity one easily obtains following permanence properties.

3.3 PROPOSITION: *Subspaces, quotient spaces by closed subspaces, and completions of $\lambda(P, \mathbb{N})$ -nuclear spaces are $\lambda(P, \mathbb{N})$ -nuclear.*

3.4 LEMMA: *For each $k \in \mathbb{N}$ there exists $r \in \mathbb{N}$ such that $\{(n + 1)\xi_n\}_n \in \lambda(P, k)$ whenever $\{\xi_n\}_n \in \lambda(P, r + 1)$.*

PROOF: Fix $k \in \mathbb{N}$. Since $\lambda(P, \infty)$ is nuclear there exists $l \in \mathbb{N}$ such that $\{p_n^k/p_n^l\}_n \in l_1$. Therefore for each $n \in \mathbb{N}$ $(p_n^k/p_n^l) + \dots + (p_n^k/p_n^l) \leq M$ and $n + 1 \leq p_n^l M/p_n^k =: M_1 p_n^l$. This implies

$$\sum_n (n + 1) |\xi_n| p_n^k \leq M_1 \sum_n |\xi_n| p_n^k p_n^l \leq M_2 \sum_n |\xi_n| p_n^r < \infty.$$

3.5 LEMMA: *Let H_1, H_2 be Hilbert spaces and $T \in L(H_1, H_2)$. Then (i) given k , T is $\tilde{\lambda}(P, k)$ -nuclear if T is of type $\lambda(P, k)$;*

(ii) given k , there exists r so that T is of type $\lambda(P, k)$ if T is $\tilde{\lambda}(P, r + 1)$ -nuclear.

PROOF: (i): We recall that for a compact operator $T \in L(H_1, H_2)$ T has a representation $Tx = \sum_n \lambda_n(x, e_n)f_n$ for suitable orthonormal sequences $\{e_n\}, \{f_n\}$ in H_1 resp. H_2 ; also $\lambda_n = \delta_n(TU_{H_1}, U_{H_2}) = s_n^{\text{app}}(T)$.

(ii) Fix k . Lemma 3.4 guarantees the existence of $r \in \mathbb{N}$ such that $\{(n + 1)\xi_n\}_n \in \lambda(P, k)$ if $\{\xi_n\}_n \in \lambda(P, r + 1)$. Now let T be $\tilde{\lambda}(P, r + 1)$ -nuclear, then T has a representation $Tx = \sum_n \gamma_n(x, a_n)y_n, \{\gamma_n\}_n \in \lambda(P, r + 1), \|a_n\| = \|y_n\| = 1, a_n \in H'_1, y_n \in H'_2$. Since $s_n^{\text{app}}(T) \leq \sum_{i=n}^\infty \gamma_i$, we have

$$\begin{aligned} \sum_{n=0}^\infty s_n^{\text{app}}(T)p_n^k &\leq \sum_{n=0}^\infty p_n^k \sum_{i=n}^\infty \gamma_i = \sum_{i=0}^\infty \sum_{n=0}^i \gamma_i p_n^k \\ &\leq \sum_{i=0}^\infty (i + 1)\gamma_i p_i^k \leq M \sum_{i=0}^\infty \gamma_i p_i^r < \infty. \end{aligned}$$

3.6 PROPOSITION: Let P be a countable, monotone, stable, nuclear G_∞ -set. Then countable direct sums of $\lambda(P, \mathbb{N})$ -nuclear spaces are $\lambda(P, \mathbb{N})$ -nuclear.

PROOF: We only indicate a partial proof and refer the reader to [1, Theorem 2.8] for the rest of the proof.

In $E = \bigoplus_{i=1}^\infty E_i$ a typical fundamental system of neighbourhoods of 0 is of the form $U = \Gamma((U_i)_i)$ where each U_k is an absolutely convex, closed, and absorbing neighbourhood of 0 in E_k and Γ represents the closed convex hull of the union. Stability of $\lambda(P, \infty)$ now gives a single valued, increasing map $\gamma: \mathbb{N} \rightarrow \mathbb{N}$ such that $\gamma(j) > j$ and $p_{2^n}^j = 0(p_n^{\gamma(j)})$. Then

- (1) $p_{2^{j-1}(2m-1)}^k \leq p_{2^j m}^k \leq M_{k,j} p_m^{\gamma^j(k)}$ for each j, k, m where $\gamma^j := \gamma \circ \gamma \circ \dots \circ \gamma$ (j -times)
- (2) $p_{2^{j-1}(2m-1)}^k \leq p_{2^{j-1}(2m-1)}^j \leq M_j p_m^{\gamma^j(j)}$ for each $k \leq j$.

Define $\beta: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ by $\beta^{-1}(j, m) := 2^{j-1}(2m - 1)$; β is a bijection and $\beta(n) = (\beta_1(n), \beta_2(n))$.

We have to show that $E = \bigoplus_i E_i$ is $\tilde{\lambda}(P, k)$ -nuclear for each $k \geq k_0$. Fix k , assume $i < k$. Using the fact that E_i is $\lambda(P, \mathbb{N})$ -nuclear and therefore nuclear for each neighbourhood $U_i \in \mathcal{U}(E_i)$ we find a neighbourhood $W_i \in \mathcal{U}(E_i)$ so that the canonical map $K_i: (E_i)_{W_i} \rightarrow (E_i)_{U_i}$ is represented by

$$K_i(x_i) = \sum_{m=0}^\infty \xi_m^i(x, a_m^i)y_m^i$$

where $\{\xi_m^i\}_m \in \lambda(P, \gamma^i(k) + 1), \|a_m^i\| \leq 1, \|y_m^i\| \leq 1$; then we have $\sum_{m=0}^\infty \xi_m^i p_m^{\gamma^i(k)} < \infty$. Now pick $t_i > 0$ so that $\sum_{m=0}^\infty t_i \xi_m^i p_m^{\gamma^i(k)} \leq 1/2^i M_{k,i}$. For

$i \geq k$ we get a representation

$$K_i(x_i) = \sum_{m=0}^{\infty} \xi_m^i \langle x, a_m^i \rangle y_m^i$$

where $\{\xi_m^i\}_m \in \lambda(P, \gamma^i(i) + 1)$, $\|a_m^i\| \leq 1$, $\|y_m^i\| \leq 1$; then we have $\sum_{m=0}^{\infty} \xi_m^i p_m^{\gamma^i(i)} < \infty$. Pick $t_i > 0$ so that $\sum_{m=0}^{\infty} t_i \xi_m^i p_m^{\gamma^i(i)} \leq 1/2^i M_i$. We then get

$$\begin{aligned} \sum_{n=1}^{\infty} t_{\beta_1(n)} |\xi_{\beta_2(n)}^{\beta_1(n)}| p_n^k &= \sum_{j=1}^{\infty} \sum_{m=1}^{\infty} t_j |\xi_m^j| p_{2^{j-1}(j,m)}^k \\ &= \sum_{j=1}^{\infty} \sum_{m=1}^{\infty} t_j |\xi_m^j| p_{2^{j-1}(2m-1)}^k \\ &= \sum_{j=1}^{k-1} \sum_{m=1}^{\infty} t_j |\xi_m^j| p_{2^{j-1}(2m-1)}^k + \sum_{j=k}^{\infty} \sum_{m=1}^{\infty} t_j |\xi_m^j| p_{2^{j-1}(2m-1)}^k \\ &\leq \sum_{j=1}^{k-1} \sum_{m=1}^{\infty} t_j |\xi_m^j| M_{k,j} p_m^{\gamma^j(k)} + \sum_{j=k}^{\infty} \sum_{m=1}^{\infty} t_j |\xi_m^j| M_j p_m^{\gamma^j(j)} \\ &\leq \sum_{j=1}^{\infty} 2^{-j} < \infty. \end{aligned}$$

Thus we have shown that $\{t_{\beta(n)} \xi_{\beta(n)}^{\beta(n)}\} \in \lambda(P, k)$. As in the proof of Theorem 2.8 in [1], we now can show that E is $\tilde{\lambda}(P, k)$ -nuclear and since this can be done for each $k \geq k_0$, E is $\lambda(P, \mathbb{N})$ -nuclear.

3.7 COROLLARY: *Arbitrary products of $\lambda(P, \mathbb{N})$ -nuclear spaces again are $\lambda(P, \mathbb{N})$ -nuclear whenever P is a countable, monotone, stable, nuclear G_{∞} -set.*

3.8 PROPOSITION: *Let P be a countable, monotone, nuclear G_{∞} -set. Then $\lambda(P, \infty)$ is $\lambda(P, \mathbb{N})$ -nuclear.*

PROOF: Since $\lambda(P, \infty)$ is a nuclear G_{∞} -space, there exists a $p^i \in P$ such that $\{1/p_n^i\} \in l_1$. Given $k, r \in \mathbb{N}$, there exists a $p^r \in P$ such that $p^r p^k < p^r$; we also find a $p^s \in P$ so that $p^r p^i < p^s$. We then have

$$\sum_n p_n^k p_n^r / p_n^s \leq M \sum_n p_n^i / p_n^s = M \sum_n (p_n^r p_n^i) / (p_n^s p_n^i) \leq M_0 \sum_n 1/p_n^i < \infty,$$

so $\lambda(P, \infty)$ is $\tilde{\lambda}(P, k)$ -nuclear for each k .

Our next result shows that $\lambda(P, \infty)$ is a universal generator for the variety of $\lambda(P, \mathbb{N})$ -nuclear spaces if P is stable.

3.9 PROPOSITION: *Let P be a countable, monotone, stable, nuclear G_{∞} -set; then each $\lambda(P, \mathbb{N})$ -nuclear space E is isomorphic to a subspace of $[\lambda(P, \infty)]^I$ for a suitable I .*

PROOF: Let $k \in \mathbb{N}$ be fixed. By stability let $\gamma: \mathbb{N} \rightarrow \mathbb{N}$ be a single valued, increasing function such that $\gamma(j) > j$ and $p_{2n}^j = 0(p_n^{\gamma(j)})$. Write $\bar{k} := \gamma^k(k)$. Since E is $\lambda(P, \mathbb{N})$ -nuclear we have $p^{\bar{k}} \in \Delta(E)$, and there exists an absolutely convex, closed, and absorbing neighbourhood $U \in \mathcal{U}(E)$ so that E'_{U^0} (U^0 is the polar of U) is a Hilbert space; now by Proposition IV.1 of [12] there exists an orthonormal basis $\{e_n^k\}_n$ in E'_{U^0} so that the set

$$A_k := \left\{ \sum_{n=0}^{\infty} \xi_n p_n^{\bar{k}} e_n^k : \sum_n |\xi_n|^2 \leq 1 \right\}$$

is equicontinuous in E' . Rearrange the set $\{e_n^k: k, n = 1, 2, \dots\}$ into a single sequence by using the bijection map $\beta: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ with $\beta^{-1}(k, n) := 2^{k-1}(2n - 1)$; use the Gram-Schmidt process to obtain a new orthonormal basis $\{e_m\}$ for E'_{U^0} . We then have $e_m = \sum_{n \geq m/2^k} (e_m, e_n^k) e_n^k$ and therefore

$$\sum_{n \geq m/2^k} |(e_m, e_n^k)|^2 / (p_n^{\bar{k}})^2 \leq \sum_{n \geq m/2^k} |(e_m, e_n^k)|^2 / (p_{m/2^k}^{\bar{k}})^2.$$

Since $p_{2^k m}^k \leq M_k p_m^{\gamma^k(k)} = M_k p_m^{\bar{k}}$ and since without loss of generality we may assume $M_k = 1$, we get

$$\sum_{n \geq m/2^k} |(e_m, e_n^k)|^2 / (p_{m/2^k}^{\bar{k}})^2 \leq \sum_{n \geq m/2^k} |(e_m, e_n^k)|^2 / (p_m^k)^2 \leq 1 / (p_m^k)^2$$

and therefore $p_m^k e_k \in A_k$.

So we have shown that there exists an orthonormal basis $\{e_m\}$ of E'_{U_0} such that $\{p_m^k e_m: m = 1, 2, \dots\}$ is equicontinuous in E'_{U_0} , for each fixed k .

Let $\mathcal{U} = (U_i: i \in I)$ be a base of neighbourhoods of 0 in E so that each U_i is absolutely convex, closed, and absorbing and E'_{U_i} is a Hilbert space. For each $i \in I$ we can get an orthonormal basis $\{e_m^i: m = 1, 2, \dots\}$ of E'_{U_i} such that the sets $B_{i,k} := \{p_m^k e_m^i: m = 1, 2, \dots\}$ are equicontinuous for each fixed k ; for each $i \in I$, define the map $T_i: E \rightarrow \lambda(P, \infty)$ by $T_i x := \{(x, e_m^i)\}_m$; T_i goes into $\lambda(P, \infty)$ and is continuous. Define $T: E \rightarrow [\lambda(P, \infty)]^I$ by $Tx := \{T_i x\}_i$. Then T is continuous and one-to-one. With obvious changes, the rest of the proof is exactly the same as of Proposition 3.4 in [8].

In [2], Fenske and Schock consider the class Ω of all l.c.s. E such that the set ω of all strictly positive, non-decreasing sequences of reals is contained in $\Delta(E)$. They prove Ω to be a stability class of nuclear spaces and therefore closed under the operations of forming completions, subspaces, quotients by closed subspaces, arbitrary products, countable direct sums, tensor products, and isomorphic

