

COMPOSITIO MATHEMATICA

B. MAUREY

G. SCHECHTMAN

Some remarks on symmetric basic sequences in L1

Compositio Mathematica, tome 38, n° 1 (1979), p. 67-76

http://www.numdam.org/item?id=CM_1979__38_1_67_0

© Foundation Compositio Mathematica, 1979, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SOME REMARKS ON SYMMETRIC BASIC SEQUENCES IN L_1

B. Maurey and G. Schechtman*

Summary

Every subspace of L_1 with an unconditional basis is isomorphic to a complemented subspace of a subspace of L_1 with a symmetric basis.

1. Introduction

The purpose of this paper is to show that, in spite of the quite simple representation of subspaces of L_1 with symmetric bases, given in [2], such spaces can be very complicated from the point of view of the structure of their complemented subspaces.

THE MAIN THEOREM: *Let $(x_i)_{i=1}^{\infty}$ be an unconditional basic sequence in L_p , $1 \leq p \leq 2$. There exists a symmetric basic sequence $(s_i)_{i=1}^{\infty}$ in L_p such that $(x_i)_{i=1}^{\infty}$ is equivalent to a block basis with constant coefficients of $(s_i)_{i=1}^{\infty}$. In particular $[x_i]_{i=1}^{\infty}$ is isomorphic to a complemented subspace of $[s_i]_{i=1}^{\infty}$.*

Theorems of such a nature were previously discovered by Lindenstrauss [8], Szankowski [15] and Davis [4]. The proof of the main theorem, given in §2, uses the technique of Davis' proof as well as some results concerning the space X_p of Rosenthal [12].

Section 3 is devoted mainly to an alternative proof of the following reduction (cf. Dacunha-Castelle and Krivine [3]): every subspace of L_1 contains some l_p if and only if every subspace of L_1 with a symmetric basis contains one.

Notions which are not explained here can be found in [9] or [10].

* Supported in part by the U.S.–Israeli Binational Science Foundation.

2. The main result

We begin with two known lemmas:

LEMMA 1 (Rosenthal [12]): *Let $1 < p < 2$. There exists a constant K_p such that if $m \geq 1$ is a real number and $(y_{i,m})_{i=1}^{\infty}$ is a sequence of independent, identically distributed, symmetric random variables (on $[0, 1]$) each taking three values in such a manner that $\|y_{i,m}\|_{L_p} = m^{-1}$ and $\|y_{i,m}\|_{L_2} = m$, then:*

$$K_p^{-1} \left\| \sum_{i=1}^{\infty} c_i y_{i,m} \right\|_{L_p} \leq \inf \left\{ \left(\sum_{i=1}^{\infty} |a_i|^2 + \left(\sum_{i=1}^{\infty} |b_i|^p \right)^{2/p} \right)^{1/2}; \forall i \geq 1, \right. \\ \left. m^{-1} a_i + m b_i = c_i \right\} \leq K_p \left\| \sum_{i=1}^{\infty} c_i y_{i,m} \right\|_{L_p}$$

PROOF: Let $q = p/(p-1)$ and let $z_{i,m} = m^{-2} y_{i,m}$, $i = 1, 2, \dots$. It is easily checked that $\|z_{i,m}\|_{L_2} = m^{-1}$ and $\|z_{i,m}\|_{L_q} = m$, $i = 1, 2, \dots$. Thus, by the main result of [12]:

$$A_p^{-1} \left\| \sum_{i=1}^{\infty} c_i z_{i,m} \right\|_{L_q} \leq \max \left\{ m^{-1} \left(\sum_{i=1}^{\infty} |c_i|^2 \right)^{1/2}, m \left(\sum_{i=1}^{\infty} |c_i|^q \right)^{1/q} \right\} \leq A_p \left\| \sum_{i=1}^{\infty} c_i z_{i,m} \right\|_{L_q}$$

and $P: L_q \rightarrow L_q$ defined by:

$$Pf = \sum_{i=1}^{\infty} \left(\int_0^1 f \frac{z_{i,m}}{\|z_{i,m}\|_{L_2}} \right) z_{i,m} = \sum_{i=1}^{\infty} \left(\int_0^1 f y_{i,m} \right) z_{i,m}$$

is a projection of norm $\|P\| \leq A_p$, where A_p depends only on p . It follows that $(y_{i,m})_{i=1}^{\infty}$ is equivalent, with constant depending on p only, to sequence of the functionals biorthogonal to the unit vector basis in the space

$$X_{q,m} = \left\{ (c_i)_{i=1}^{\infty}; \|(c_i)_{i=1}^{\infty}\| = \max \left\{ m^{-1} \left(\sum_{i=1}^{\infty} |c_i|^2 \right)^{1/2}, m \left(\sum_{i=1}^{\infty} |c_i|^q \right)^{1/q} \right\} < \infty \right\}$$

and the result follows. ■

Let $1 \leq p < 2$, $m \geq 1$ and define a norm on ℓ_2 by

$$\|x\|_m = \inf \{ (\|y\|_{\ell_2}^2 + \|z\|_{\ell_p}^2)^{1/2}; m^{-1}y + mz = x \}.$$

The unit vectors $(e_i)_{i=1}^{\infty}$ in ℓ_2 clearly form a 1-symmetric basis for

$(\ell_2, |\cdot|_m)$. We define

$$\lambda_m(n) = \left| \sum_{i=1}^n e_i \right|_m \quad n = 1, 2, \dots$$

In the following lemma we collect some properties of the $\lambda_m(n)$'s. For a proof see [4] or [10, lemma 3.b.3 and the proof of proposition 3.b.4].

LEMMA 2: (1) $\lambda_m(n) = (m^{-2}n^{-1} + m^2n^{-2/p})^{-1/2}$ for every integer n and every $m \geq 1$.

(2) For a fixed n :

$$\max_{m \geq 1} \lambda_m(n) = 2^{-1/2} n^{(1/2+1/p)(1/2)}$$

and the maximum is attained at $m = m(n) = n^{(1/p-1/2)(1/2)}$

(3) For any sequence of positive numbers $(\epsilon_i)_{i=1}^\infty$ there exists a subsequence $(n_i)_{i=1}^\infty$ of the integers such that, putting $m_i = m(n_i)$, $i = 1, 2, \dots$, we get:

$$\sum_{j=1, j \neq i}^\infty \lambda_{m_i}(n_j) / \lambda_{m_j}(n_j) < \epsilon_k$$

and

$$\sum_{i=1}^\infty \frac{1}{m_i} \leq 1$$

We are ready now for the proof of the main theorem. The proof is written for the case $p = 1$ only, the case $1 < p < 2$ requires only minor changes (and the case $p = 2$ is trivial).

PROOF OF THE MAIN THEOREM: Let $(x_k)_{k=1}^\infty$ be a normalized unconditional basic sequence in L_1 and let $1 < p < 2$. $(\epsilon_i)_{i=1}^\infty$ will denote a sequence of positive numbers to be specified later and $(n_i)_{i=1}^\infty$ will denote subsequence of the integers with the properties stated in Lemma 2.3.

Let $(y_{i,k})_{i=1, k=1}^\infty$ be a double sequence of independent, symmetric, three-valued random variables on $[0, 1]$ such that

$$\|y_{i,k}\|_{L_p} = m_k^{-1} \text{ and } \|y_{i,k}\|_{L_2} = m_k \quad i, k = 1, 2, \dots$$

By Lemma 1:

$$(1) \quad K_p^{-1} |(c_i)_{i=1}^\infty|_{m_k} \leq \left\| \sum_{i=1}^\infty c_i y_{i,k} \right\| \leq K_p |(c_i)_{i=1}^\infty|_{m_k}$$

for all scalars $(c_i)_{i=1}^\infty$, where K_p depends only on p .

Define $s_i \in L_p([0, 1]^2)$, $i = 1, 2, \dots$ by:

$$s_i(r, t) = \sum_{k=1}^{\infty} y_{i,k}(r)x_k(t)$$

Then:

$$\|s_i\|_{L_1(L_p)} = \int \left(\int |s_i(r, t)|^p dr \right)^{1/p} dt \leq \sum_{k=1}^{\infty} \|y_{i,k}\|_{L_p} \leq 1$$

and $s_i(r, t)$, $i = 1, 2, \dots$ clearly constitutes a 1-symmetric basic sequence in $L_1(L_p)$.

Choose a sequence $(\Gamma_\ell)_{\ell=1}^\infty$ of disjoint subsets of the integers with $\bar{\Gamma}_\ell = n_\ell$ (\bar{A} denotes the cardinality of the set A), and put

$$w_\ell = \sum_{i \in \Gamma_\ell} s_i / \lambda_{m_\ell}(n_\ell) \quad \ell = 1, 2, \dots$$

$(w_\ell)_{\ell=1}^\infty$ is a block basis with constant coefficients of $(s_i)_{i=1}^\infty$; we are going to show that, if the ϵ_i 's are chosen properly, $(w_\ell)_{\ell=1}^\infty$ is equivalent to $(x_\ell)_{\ell=1}^\infty$.

Let

$$v_\ell(n, t) = \sum_{i \in \Gamma_\ell} y_{i,\ell}(r) \cdot x_\ell(t) / \lambda_{m_\ell}(n_\ell) \quad \ell = 1, 2, \dots$$

and note that, by the triangle inequality, (1) and Lemma 2:

$$\begin{aligned} (2) \quad \left\| \sum_{\ell=1}^{\infty} a_\ell w_\ell \right\| &= \left\| \sum_{\ell=1}^{\infty} a_\ell \sum_{k=1}^{\infty} \sum_{i \in \Gamma_\ell} y_{i,k}(r)x_k(t) / \lambda_{m_\ell}(n_\ell) \right\|_{L_1(L_p)} \\ &\leq \left\| \sum_{\ell=1}^{\infty} a_\ell v_\ell(n, t) \right\|_{L_1(L_p)} \\ &\quad + \sum_{\ell=1}^{\infty} |a_\ell| \sum_{k \neq \ell} \left\| \sum_{i \in \Gamma_\ell} y_{i,k} \right\|_{L_p} / \lambda_{m_\ell}(n_\ell) \\ &\leq \left\| \sum_{\ell=1}^{\infty} a_\ell v_\ell(r, t) \right\|_{L_1(L_p)} \\ &\quad + K_p \sum_{\ell=1}^{\infty} |a_\ell| \sum_{k \neq \ell} \lambda_{m_k}(n_\ell) / \lambda_{m_\ell}(n_\ell) \\ &\leq \left\| \sum_{\ell=1}^{\infty} a_\ell v_\ell(r, t) \right\|_{L_1(L_p)} + K_p \left(\sum_{\ell=1}^{\infty} \epsilon_\ell \right) \max_{i \leq \ell < \infty} |a_\ell| \end{aligned}$$

for all scalars $(a_\ell)_{\ell=1}^\infty$. And similarly:

$$(3) \quad \left\| \sum_{\ell=1}^{\infty} a_\ell w_\ell \right\| \geq \left\| \sum_{\ell=1}^{\infty} a_\ell v_\ell(r, t) \right\|_{L_1(L_p)} - K_p \left(\sum_{i=1}^{\infty} \epsilon_i \right) \max_{1 \leq \ell < \infty} |a_\ell|$$

Now

$$\left(\sum_{i \in \Gamma_\ell} y_{i,\ell} \right)_{\ell=1}^\infty$$

is a sequence of symmetric independent random variables and thus forms a 1-unconditional basic sequence in L_p [cf. 12]; it follows that $(v_\ell)_{\ell=1}^\infty$ is a 1-unconditional basic sequence in $L_1(L_p)$, thus

$$(4) \quad \left\| \sum a_\ell v_\ell(r, t) \right\|_{L_1(L_p)} \geq \max_{1 \leq \ell < \infty} |a_\ell| \cdot \min_{1 \leq \ell < \infty} \|v_\ell(r, t)\|_{L_1(L_p)} \geq K_p^{-1} \max_{1 \leq \ell < \infty} |a_\ell|$$

Hence, if the ϵ_i 's satisfy

$$\sum_{i=1}^{\infty} \epsilon_i < \frac{1}{2} K_p^{-2}$$

we get from (2), (3) and (4) that

$$\frac{1}{2} \left\| \sum a_\ell v_\ell \right\| \leq \left\| \sum a_\ell w_\ell \right\| \leq \frac{3}{2} \left\| \sum a_\ell v_\ell \right\|$$

for all choices of scalars $(a_\ell)_{\ell=1}^\infty$, so it is enough to prove that $(v_\ell)_{\ell=1}^\infty$ is equivalent to $(x_\ell)_{\ell=1}^\infty$.

The sequence

$$\left(\sum_{i \in \Gamma_\ell} y_{i,\ell} / \lambda_{m_\ell}(n_\ell) \right)_{\ell=1}^\infty$$

is, as was already mentioned, a bounded away from zero 1-unconditional orthogonal sequence and, for $\ell = 1, 2, \dots$

$$\left\| \sum_{i \in \Gamma_\ell} y_{i,\ell} / \lambda_{m_\ell}(n_\ell) \right\|_{L_2} = \|y_{1,\ell}\|_{L_2} \cdot n_\ell^{1/2} / \lambda_{m_\ell}(n_\ell) = m_\ell \cdot n_\ell^{1/2} / \lambda_{m_\ell}(n_\ell) = 2^{1/2}$$

by Lemma 2.

It follows easily that

$$\left(\sum_{i \in I_\ell} y_{i,d} \lambda_{m_\ell}(n_\ell) \right)_{\ell=1}^\infty$$

is equivalent, in L_p , to the unit vector basis of ℓ_2 and thus there exists a constant K_p^1 such that

$$(K_p^1)^{-1} \left\| \left(\sum_{\ell=1}^\infty a_\ell^2 x_\ell^2 \right)^{1/2} \right\|_{L_1} \leq \left\| \sum_{\ell=1}^\infty a_\ell v_\ell(r, t) \right\|_{L_1(L_p)} \leq K_p^1 \left\| \left(\sum_{\ell=1}^\infty a_\ell^2 x_\ell^2 \right)^{1/2} \right\|_{L_1}$$

Finally, by the unconditionality of $(x_\ell)_{\ell=1}^\infty$ and by Khinchine's inequality, the expression

$$\left\| \left(\sum_{\ell=1}^\infty a_\ell^2 x_\ell^2 \right)^{1/2} \right\|_{L_1} \text{ is equivalent to } \left\| \sum_{\ell=1}^\infty a_\ell x_\ell \right\|_{L_1}. \quad \blacksquare$$

CONCLUSION: *Let $1 \leq p \leq 2$ there exists a 1-symmetric sequence $(s_i)_{i=1}^\infty$ in L_p such that every unconditional basic sequence $(x_i)_{i=1}^\infty$ in L_p is equivalent, with constant depending only on the unconditionality constant of $(x_i)_{i=1}^\infty$, to a block basis with constant coefficients of $(s_i)_{i=1}^\infty$.*

PROOF: By [14] there exists, in L_p , a 1-unconditional basic sequence such that any other unconditional basic sequence in L_p is equivalent, with constant depending on the unconditionality constant only, to a subsequence of it. Apply the main theorem to this universal basic sequence. \blacksquare

3. Dacunha-Castelle and Krivine's reduction

This section is devoted to the proof of the following proposition.

PROPOSITION 3: *If every subspace of L_1 with a symmetric basis contains an isomorphic copy of ℓ_p for some $1 \leq p \leq 2$ then every subspace of L_1 contains such a subspace.*

We begin by recalling two definitions.

DEFINITION 1 ([6], [7]): An unconditional basis $(x_i)_{i=1}^\infty$ is said to be q -concave ($1 \leq q < \infty$) if there exists a constant $c > 0$ such that for

every N and every N elements $y_n = \sum_{i=1}^{\infty} a_{i,n} x_i$, $n = 1, \dots, N$ in $[x_i]_{i=1}^{\infty}$ we have:

$$\left\| \sum_{i=1}^{\infty} \left(\sum_{n=1}^N |a_{i,n}|^q \right)^{1/q} x_i \right\| \geq c \left(\sum_{n=1}^N \|y_n\|^q \right)^{1/q}$$

DEFINITION 2 ([5]): Given $1 \leq p \leq 2$, an 1-unconditional normalized basis $(x_i)_{i=1}^{\infty}$ and a sequence $m_n \nearrow \infty$ with $\sum_{n=1}^{\infty} m_n^{-1} < \infty$, let $Y = Y(\ell_2, \ell_p, (x_i)_{i=1}^{\infty}, (m_n)_{n=1}^{\infty})$ be the space of all sequences of scalars $\alpha \in \ell_2$ such that

$$\|\alpha\|_Y = \left\| \sum_{n=1}^{\infty} |\alpha|_{m_n} x_n \right\|_X < \infty$$

($|\cdot|_{m_n}$ is defined in the preceding section). $(e_i)_{i=1}^{\infty}$ will denote the unit vectors in this space: note that they form a 1-symmetric basis for Y .

PROPOSITION 4: Let $q < 2$ and let $(x_i)_{i=1}^{\infty}$ be a q -concave 1-unconditional basic sequence in L_1 , let $q < p < 2$, and let $(m_n)_{n=1}^{\infty}$ and $(s_i)_{i=1}^{\infty}$ be as in the proof of the main theorem. Then $(s_i)_{i=1}^{\infty}$ is equivalent to $(e_i)_{i=1}^{\infty}$.

PROOF: We use the same notation as in the proof of the main theorem. By Khinchine's inequality and the triangle inequality in $\ell_{2/p}$ we get that:

$$\begin{aligned} \left\| \sum_{i=1}^{\infty} a_i s_i \right\|_{L_1(L_p)} &= \int \left(\int \left| \sum_{k=1}^{\infty} \sum_{i=1}^{\infty} a_i y_{i,k}(r) x_k(t) \right|^p dr \right)^{1/p} dt \\ &\geq A_p \int \left(\int \left(\sum_{k=1}^{\infty} \left| \sum_{i=1}^{\infty} a_i y_{i,k}(r) x_k(t) \right|^2 \right)^{p/2} dr \right)^{1/2} dt \\ &\geq A_p \int \left(\sum_{k=1}^{\infty} \left(\int \left| \sum_{i=1}^{\infty} a_i y_{i,k}(r) \right|^p dr \right)^{2/p} |x_k(t)|^2 \right)^{1/2} dt \\ &\geq A_p K_p^{-1} \|(a_i)_{i=1}^{\infty}\|_Y \end{aligned}$$

A_p denotes here the Khinchine's constant of L_p : note that we have used the fact that for a fixed t , $(\sum_{i=1}^{\infty} a_i y_{i,k}(r) x_k(t))_{k=1}^{\infty}$ constitutes a 1-unconditional basic sequence in L_p .

For the other side inequality we need the fact [cf. 1] that there exists an isometry (into) $T: L_p \rightarrow L_1$ and a constant C such that

$$C\|f\|_{L_p} = C\|Tf\|_{L_1} \geq \|Tf\|_{L_q} \geq \|Tf\|_{L_1} = \|f\|_{L_p}$$

for all $f \in L_p$. Using this fact and the q -concavity of $(x_i)_{i=1}^\infty$ (with constant c , say) we get that,

$$\begin{aligned}
\left\| \sum_{i=1}^\infty a_i s_i \right\|_{L_1(L_p)} &= \iint \left| \sum_{k=1}^\infty \sum_{i=1}^\infty a_i (Ty_{i,k})(r) x_k(t) \right| dt dr \\
&\leq \left(\iint \left\| \sum_{k=1}^\infty \sum_{i=1}^\infty a_i (Ty_{i,k})(r) x_k \right\|_{L_1}^q dr \right)^{1/q} \\
&\leq c^{-1} \left\| \sum_{k=1}^\infty \left(\int \left| \sum_{i=1}^\infty a_i Ty_{i,k}(r) \right|^q dr \right)^{1/q} x_k \right\|_{L_1} \\
&\leq c^{-1} C \left\| \sum_{k=1}^\infty \left\| \sum_{i=1}^\infty a_i y_{i,k} \right\|_{L_p} x_k \right\|_{L_1} \\
&\leq c^{-1} CK_p \|(a_i)_{i=1}^\infty\|_Y. \quad \blacksquare
\end{aligned}$$

We shall need also the following standard.

LEMMA 5: *Let U be an infinite dimensional subspace of $Y = Y(\ell_2, \ell_p, (x_i)_{i=1}^\infty, (m_n)_{n=1}^\infty)$. Then U contains a subspace which embeds isomorphically in $[x_i]_{i=1}^\infty$.*

PROOF: Using standard arguments (compare for example the proof of proposition 3.b.4 in [10]) the proof reduces to showing that the identity map from Y to ℓ_2 is strictly singular.

Now any infinite dimensional subspace of ℓ_2 contains a norm one vector with arbitrarily small ℓ_∞ norm. Fix n and let x be a norm one vector in ℓ_2 such that $\|x\|_{\ell_\infty} \leq m_n^{-2p/(2-p)}$. Let y and z be elements of ℓ_2 such that

$$|x|_{m_n} \geq (\|y\|_2^2 + \|z\|_p^2)^{1/2} - 1 \quad \text{and} \quad m_n^{-1}y + m_n z = x$$

We may clearly assume also that $\|m_n z\|_{\ell_\infty} \leq \|x\|_{\ell_\infty}$. If $\|m_n^{-1}y\|_{\ell_2} \geq \frac{1}{2}$ then $\|x\|_Y \geq |x|_{m_n} \geq m_n/2 - 1$. If $\|m_n^{-1}y\|_{\ell_2} < \frac{1}{2}$ then $\|m_n z\|_{\ell_2} \geq \frac{1}{2}$ and we get:

$$\frac{1}{2} \leq \|m_n z\|_{\ell_2} \leq \|m_n z\|_{\ell_p}^{p/2} \|m_n z\|_{\ell_\infty}^{(2-p)/2}$$

and

$$\|m_n z\|_{\ell_p} \geq m_n^2 / 2^{2/p}$$

so that

$$\|x\|_Y \geq |x|_{m_n} \geq m_n / 2^{2/p} - 1$$

Hence every infinite dimensional subspace of Y contains vectors with norm one in ℓ_2 and with arbitrarily big norm in Y . ■

PROOF OF PROPOSITION 3: Assume that every subspace of L_1 with a symmetric basis contains some ℓ_p . Notice first that if $(x_i)_{i=1}^\infty$ is a 1-unconditional basic sequence in L_1 which is q -concave for some $q < 2$ then, by the main theorem, Proposition 4, Lemma 5 and the fact [cf. 11] that every infinite dimensional subspace of ℓ_p contains an isomorph of ℓ_p , $[x_i]_{i=1}^\infty$ contains ℓ_p isomorphically for some p (necessarily $1 \leq p \leq q$).

Now let X be a subspace of L_1 . If X does not contain ℓ_1 then by Rosenthal's theorem [13] X is isomorphic to a subspace of L_r for some $1 < r \leq 2$ and since L_r has an unconditional basis the image of X in L_r contains an unconditional basic sequence $(y_i)_{i=1}^\infty$. It is clearly enough to show that $[y_i]_{i=1}^\infty$ contains some ℓ_p .

Let $(g_i)_{i=1}^\infty$ be a sequence in L_1 isometrically equivalent to the unit vector basis of $\ell_{2/r}$ (for instance a sequence of L_1 -norm one, independent $2/r$ stable random variables), and define, for $i = 1, 2, \dots$ and $0 \leq s, t \leq 1$

$$x_i(s, t) = g_i(s)|y_i(t)|^r$$

Then $(x_i)_{i=1}^\infty$ is 1-unconditional in $L_1((0, 1)^2)$ and by the triangle inequality in $\ell_{2/r}$

$$\begin{aligned} \left\| \sum_{i=1}^\infty \left(\sum_{j=1}^\infty |a_{i,j}|^{2/r} \right)^{r/2} x_i \right\| &= \int_0^1 \left(\sum_{j=1}^\infty \sum_{i=1}^\infty |a_{i,j}|^{2/r} |y_i(t)|^2 \right)^{r/2} dt \\ &\geq \left(\sum_{j=1}^\infty \left(\int_0^1 \left(\sum_{i=1}^\infty |a_{i,j}|^{2/r} |y_i(t)|^2 \right)^{r/2} dt \right)^{2/r} \right)^{r/2} \\ &= \left(\sum_{j=1}^\infty \left\| \sum_{i=1}^\infty a_{i,j} x_i \right\|^{2/r} \right)^{r/2} \end{aligned}$$

i.e. $(x_i)_{i=1}^\infty$ is $2/r$ concave. Thus, it follows from the first part of the proof that $[x_i]_{i=1}^\infty$ contains a subspace isomorphic to ℓ_p for some $1 \leq p \leq 2/r$. Passing to a smaller subspace we may assume that there exist disjoint blocks

$$b_n = \sum_{i \in E_n} a_i x_i \qquad n = 1, 2, \dots$$

of $(x_i)_{i=1}^\infty$ which are equivalent to the usual basis of ℓ_p . It is easily

checked that the blocks

$$c_n = \sum_{i \in E_n} |a_i|^{1/r} y_i \quad n = 1, 2, \dots$$

of $(y_i)_{i=1}^{\infty}$ are equivalent to the usual basis of ℓ_p . ■

REFERENCES

- [1] J. BRETAGNOLLE, D. DACUNHA-CASTELLE and J.L. KRIVINE: Lois stable et espaces L^p , *Ann. Inst. Henri Poincaré, série B*, 2 (1966) 231–263.
- [2] D. DACUNHA-CASTELLE: “Variables aléatoires échangeable et espaces d’Orlicz.” Séminaire Maurey-Schwartz, École Polytechnique (1974–1975).
- [3] D. DACUNHA-CASTELLE and J.L. KRIVINE: Sous-espaces de L^1 , *Israel J. Math.*, 26 (1977) 320–351.
- [4] W.J. DAVIS: Embedding spaces with unconditional bases, *Israel J. Math.*, 20 (1975) 189–191.
- [5] W.J. DAVIS, T. FIGIEL, W.B. JOHNSON and A. PELCZYNSKI: Factoring weakly compact operators, *J. Funct. Anal.*, 17 (1974) 311–327.
- [6] E. DUBINSKY, A. PELCZYNSKI and H.P. ROSENTHAL: On Banach spaces X for which $\pi(\mathcal{L}_x, X) = B(\mathcal{L}_x, X)$, *Studia Math.*, 44 (1972) 617–648.
- [7] T. FIGIEL and W.B. JOHNSON: A uniformly convex Banach space which contains no ℓ_p , *Compositio Math.*, 29 (1974) 179–190.
- [8] J. LINDENSTRAUSS: A remark on symmetric bases, *Israel J. Math.*, 13 (1972) 317–320.
- [9] J. LINDENSTRAUSS and L. TZAFRIRI: Classical Banach spaces. *Lecture Notes in Mathematics* 338. Springer, Berlin etc. (1973).
- [10] J. LINDENSTRAUSS and L. TZAFRIRI: Classical Banach Spaces, sequence, spaces. Springer, Berlin.
- [11] A. PELCZYNSKI: Projections in certain Banach spaces, *Studia Math.*, 19 (1960) 209–228.
- [12] H.P. ROSENTHAL: On the subspaces of L^p ($p > 2$) spanned by sequences of independent random variables, *Israel J. Math.*, 8 (1970) 273–303.
- [13] H.P. ROSENTHAL: On subspaces of L^p , *Ann. Math.*, 97 (1973) 344–373.
- [14] G. SCHECHTMAN: On Pelczynski’s paper “Universal bases”, *Israel J. Math.*, 22 (1975) 181–184.
- [15] A. SZANKOWSKI: Embedding Banach spaces with unconditional bases into spaces with symmetric bases, *Israel J. Math.*, 15 (1973) 53–59.

(Oblatum 1–VI–1977)

B. Maurey
 École Polytechnique
 Paris, France
 and
 The Institute for Advanced Studies
 The Hebrew University
 Jerusalem
 Israel

G. Schechtman
 The Institute for Advanced Studies
 The Hebrew University
 Jerusalem
 Israel