

# COMPOSITIO MATHEMATICA

LORENZO ROBBIANO

## **A theorem on normal flatness**

*Compositio Mathematica*, tome 38, n° 3 (1979), p. 293-298

[http://www.numdam.org/item?id=CM\\_1979\\_\\_38\\_3\\_293\\_0](http://www.numdam.org/item?id=CM_1979__38_3_293_0)

© Foundation Compositio Mathematica, 1979, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## A THEOREM ON NORMAL FLATNESS

Lorenzo Robbiano\*

### Introduction

The beginning of the story is in [4] where the first theorem of transitivity of normal flatness is given.

More or less the problem is the following. Let  $R$  be a local ring,  $I, J, \Lambda = I + J$  ideals of  $R$  and assume that  $J$  is generated by a regular sequence mod  $I$ ; then how is it possible to relate the properties that  $G(I)$  (the graded ring associated to  $I$ ) is a free  $R/I$ -module and  $G(\Lambda)$  is a free  $R/\Lambda$ -module?

As I said, the first answer was in [4] and it was the starting point for successive improvements (see for instance [1], [2], [5], [8]); a theory of normal flatness was constructed ([3]) and also some questions on normal torsionfreeness were solved (see for instance [1], [6], [7], [8]). In this kind of problems it always happened that the authors assumed  $J$  to be in particular position with respect to  $I$  (essentially, as I said,  $J$  is generated by a regular sequence mod  $I$ ). The purpose of the present paper is to produce a new theorem (Theorem 10) of transitivity of normal flatness in the case that  $I, J$  are in “symmetric position”; in order to prove it, I firstly achieve some results which give a connection between the graded rings  $G(I), G(J), G(\Lambda)$ ; in this way it is also possible to give a new insight in the proofs of some known theorems.

In this paper all rings are supposed to be commutative, Noetherian and with identity.

Let  $R$  be a ring,  $I, J, \Lambda = I + J$  ideals of  $R$ ; we denote by  $R(I; J)$  the “Rees algebra” associated with the pair  $(I; J)$  i.e. the graded ring  $\bigoplus_n R_n(I; J)$  where  $R_n(I, J) = \bigoplus_{r+s=n} I^r J^s$  and the multiplication is the obvious one.

\*This work was supported by C.N.R. (Consiglio Nazionale delle Ricerche).

Of course  $R(I; R) = R(I)$  the usual Rees algebra.

We shall denote by  $G(I; J)$  the graded ring associated with the pair  $(I; J)$  i.e. the graded ring  $R(I; J) \otimes_R R/\Lambda$ .

It is clear that  $G(I; J) = \bigoplus G_n(I; J)$  where  $G_n(I; J) = \bigoplus_{r+s=n} G_{r,s}(I; J)$  and  $G_{r,s}(I; J) = I^r J^s \otimes_R R/\Lambda \simeq I^r J^s / I^r J^s \Lambda$ .

It is also clear that we have canonical epimorphisms

$$G(I) \otimes_R G(J) \longrightarrow G(I; J) \longrightarrow G(\Lambda)$$

LEMMA 1: Let  $r, n$  be integers such that  $0 \leq r < n$  and call  $s = n - r$ ; let  $R$  be a ring,  $I, J, \Lambda = I + J$  ideals of  $R$  and assume that  $I^{r+1} \cap J^{s-1} = I^{r+1} J^{s-1}$ . Then the canonical epimorphism

$$I^{r+1} \Lambda^{s-2} / I^{r+1} \Lambda^{s-1} \oplus I^r J^{s-1} / I^r J^{s-1} \Lambda \longrightarrow I^r \Lambda^{s-1} / I^r \Lambda^s$$

is an isomorphism.

PROOF: An element of the direct sum is of the form  $(\bar{\alpha}, \bar{\beta})$  where  $\alpha \in I^{r+1} \Lambda^{s-2}, \beta \in I^r J^{s-1}$ .

With the obvious meaning of the symbols its image is  $\overline{\alpha + \beta}$ .

Suppose now that  $\alpha + \beta \in I^r \Lambda^s$ ; then

$$\begin{aligned} \alpha &\in I^{r+1} \Lambda^{s-2} \cap (I^r J^{s-1} + I^r \Lambda^s) \\ &= I^{r+1} \Lambda^{s-2} \cap (I^r J^{s-1} + I^r J^s + I^{r+1} J^{s-1} + I^{r+2} J^{s-2} + \dots + I^{r+s}) \\ &= I^{r+1} \Lambda^{s-2} \cap (I^r J^{s-1} + I^{r+2} \Lambda^{s-2}) = I^{r+2} \Lambda^{s-2} \\ &\quad + (I^{r+1} \Lambda^{s-2} \cap I^r J^{s-1}) \subseteq I^{r+2} \Lambda^{s-2} + I^{r+1} \cap J^{s-1} \\ &= I^{r+2} \Lambda^{s-2} + I^{r+1} J^{s-1} = I^{r+1} \Lambda^{s-1} \text{ which means } \bar{\alpha} = 0. \end{aligned}$$

Now we have

$$\begin{aligned} \beta &\in I^r J^{s-1} \cap I^r \Lambda^s \\ &= I^r J^{s-1} \cap (I^{r+s} + \dots + I^{r+2} J^{s-2} + I^{r+1} J^{s-1} + I^r J^s) \\ &= I^{r+1} J^{s-1} + I^r J^s + I^r J^{s-1} \cap I^{r+2} \Lambda^{s-2} \\ &\subseteq I^r J^{s-1} \Lambda + J^{s-1} \cap I^{r+1} = I^r J^{s-1} \Lambda \text{ which means } \bar{\beta} = 0. \end{aligned}$$

PROPOSITION 2. Let  $n$  be a positive integer; let  $R$  be a ring,  $I, J, \Lambda = I + J$  ideals of  $R$ . Assume that  $I^r \cap J^s = I^r J^s$  for every pair of positive integers  $r, s$  such that  $r + s = n$ . Then the canonical epimorphism  $G_{n-1}(I; J) \longrightarrow G_{n-1}(\Lambda)$  is an isomorphism.

PROOF. It is an easy consequence of Lemma 1.

LEMMA 3. Let  $r, s$  be positive integers; with the usual notations assume that  $I^r \cap J = I^r J$  and  $G_s(J)$  is a free  $R/J$ -module. Then the canonical epimorphism  $G_r(I) \otimes_R G_s(J) \xrightarrow{\pi} G_{r,s}(I; J)$  is an isomorphism.

PROOF. Let  $\alpha = rk_{R/J}(G_s(J)) = rk_{R/\Lambda}(J^s/J^s\Lambda)$  and let us choose  $\theta_1, \dots, \theta_\alpha \in J^s$  so that  $\bar{\theta}_1, \dots, \bar{\theta}_\alpha$  is a free basis of  $G_s(J)$ . Then the homomorphism  $\rho: G_r(I) \otimes G_s(J) \longrightarrow (I'/I'\Lambda)^\alpha$  defined by  $\rho(\bar{a} \otimes \bar{\theta}_i) = (0, \dots, 0, \bar{a}, 0, \dots, 0)$  where  $\bar{a}$  is in the  $i$ th place, is clearly an isomorphism.

Now we can consider the homomorphism  $\tau: (I'/I'\Lambda)^\alpha \longrightarrow I'J^s/I'J^s\Lambda$  defined by  $\tau(\bar{a}_1, \dots, \bar{a}_\alpha) = \overline{\sum a_i \theta_i}$ .

It is clear that  $\pi = \tau \circ \rho$ , hence it is enough to prove that  $\tau$  is injective. Let  $\bar{0} = \tau(\bar{a}_1, \dots, \bar{a}_\alpha) = \overline{\sum a_i \theta_i}$ ; this means that  $\sum a_i \theta_i \in I'J^s\Lambda$  hence  $\sum a_i \theta_i = \sum b_i \theta_i \pmod{J^{s+1}}$  where  $b_i \in I'\Lambda$ . Therefore  $\sum (a_i - b_i) \theta_i = 0 \pmod{J^{s+1}}$  which implies  $a_i - b_i \in J$  for every  $i$ . We get  $a_i \in I' \cap (I'\Lambda + J) = I'\Lambda + I' \cap J = I'\Lambda + I'J = I'\Lambda$  for every  $i$ , and this concludes the proof.

LEMMA 4. *With the usual notations assume that  $I \cap J = IJ$  and  $G_r(J)$  is a free  $R/J$ -module for  $r = 1, \dots, n - 1$ . Then  $I \cap J^n = IJ^n$ .*

PROOF. It is an easy consequence of the fact that, given  $I_1, I_2$  ideals of a ring  $R$ ,  $I_1 \cap I_2 = I_1 \cdot I_2$  is equivalent to  $\text{Tor}_1^R(R/I_1, R/I_2) = 0$ .

THEOREM 5. *Let  $R$  be a ring,  $I, J, \Lambda = I + J$  ideals of  $R$  such that  $I \cap J = IJ$ . Let  $n$  be a positive integer and assume that  $G_i(I)$  is a free  $R/I$ -module for  $i = 1, \dots, n - 1$  and  $G_j(J)$  is a free  $R/J$ -module for  $j = 1, \dots, n - 1$ .*

*Then the canonical epimorphism  $\bigoplus_{\lambda+\mu=\nu} G_\lambda(I) \otimes_R G_\mu(J) \longrightarrow G_\nu(\Lambda)$  is an isomorphism, hence  $G_\nu(\Lambda)$  is a free  $R/\Lambda$ -module for  $\nu = 1, \dots, n - 1$ .*

PROOF. Using Lemma 4 we get  $I' \cap J^s = I'J^s$  for  $r = 1, \dots, n - 1$ ,  $s = 1, \dots, n - 1$ ; hence  $\bigoplus_{\lambda+\mu=\nu} G_{\lambda,\mu}(I; J) \simeq G_\nu(\Lambda)$  for  $\nu = 1, \dots, n - 1$  by Proposition 2.

Using Lemma 3 we are done.

REMARK. It is possible to use theorem 5 to prove the following (known) fact. Let  $R$  be a local ring, let  $a_1, \dots, a_n; b_1, \dots, b_m$  be  $R$ -sequences such that if we call  $I = (a_1, \dots, a_n)$ ,  $J = (b_1, \dots, b_m)$ ,  $I \cap J = IJ$ .

Then  $a_1, \dots, a_n, b_1, \dots, b_m$  is a regular  $R$ -sequence.

PROOF. (Hint). Use Theorem 5 to prove that  $G(I + J)$  is a polynomial ring in  $n + m$  indeterminates.

LEMMA 6. *With the usual notations denote by  $\bar{\phantom{x}}$  the reduction modulo  $I$ . Then the canonical sequence of  $R/\Lambda$ -modules*

$$0 \longrightarrow I \cap J^n / IJ^n + I \cap J^{n+1} \longrightarrow J^n / J^{n+1} \xrightarrow{\pi_n} \bar{J}^n / \bar{J}^{n+1} \longrightarrow 0$$

*is exact for every positive integer  $n$ .*

PROOF.  $\bar{J}^n / \bar{J}^{n+1} \simeq (J^n + I) / (J^{n+1} + I) \simeq J^n / (J^{n+1} + J^n \cap I)$ .

Hence  $\text{Ker } \pi_n \simeq (J^{n+1} + J^n \cap I) / (J^{n+1} + J^n I)$

$$\simeq (I \cap J^n) / (I \cap J^n \cap (IJ^n + J^{n+1})) \simeq (I \cap J^n) / (IJ^n + I \cap J^{n+1}).$$

Henceforth we assume that  $R$  is local.

COROLLARY 7. *With the notations of Lemma 6, assume that  $R$  is local and let  $n$  be a positive integer. Then the following conditions are equivalent*

- (i)  $\pi_r$  is an isomorphism for every  $r \geq n$ .
- (ii)  $I \cap J^r = IJ^r$  for every  $r \geq n$ .

PROOF. The only thing to be proved is that (i) implies  $I \cap J^n = IJ^n$ . Now, if  $\pi_r$  is an isomorphism for every  $r \geq n$ , then  $I \cap J^n = IJ^n + I \cap J^{n+1} = IJ^n + IJ^{n+1} + I \cap J^{n+2} = IJ^n + I \cap J^{n+2} = \dots = IJ^n + I \cap J^{n+k}$ . Hence  $I \cap J^n = \bigcap_k (IJ^n + I \cap J^{n+k}) = IJ^n$ .

COROLLARY 8. *With the usual notations assume that  $R$  is local  $I \cap J = IJ$  and  $G(J)$  is a free  $R/J$ -module. Then  $G(\bar{J}) \simeq G(J) \otimes_R R/\Lambda$ , hence is a free  $R/\Lambda$ -module.*

PROOF. Using Lemma 4 we get that  $I \cap J^n = IJ^n$  for every  $n$ , hence we can apply Corollary 7.

LEMMA 9. *Let  $R$  be a local ring,  $I, J, \Lambda$  as above,  $B = R/I$ ,  $A = B/\bar{\Lambda} \simeq R/\Lambda$  ( $\bar{\phantom{x}}$  denotes "modulo  $I$ ").*

*Let  $L$  be a finite free  $B$ -module and  $T$  a submodule of  $L$ . Suppose*

- (i)  $T_{\mathfrak{p}} = 0$  for every  $\mathfrak{p} \in \text{Ass}(A)$ .
- (ii)  $J^n / J^n \Lambda$  is a free  $A$ -module for every  $n$ .
- (iii)  $(I \cap J^n)_{\mathfrak{p}} = (IJ^n)_{\mathfrak{p}}$  for every  $\mathfrak{p} \in \text{Ass}(A)$ .

*Then  $T = 0$ .*

PROOF. If we make the identification of  $L$  with  $(R/I)^\alpha$  for a suitable  $\alpha$ , then  $T$  can be considered as the reduction modulo  $I$  of a sub-

module  $K$  of  $R^\alpha$ . Because of the type of the argument, we may assume that  $\alpha = 1$ , so it is enough to prove that  $K \subset I$ .

Using (i) we easily get that  $T \subset \bar{\Lambda}$ , hence  $K \subset \Lambda$ . Therefore if  $x \in K$  there exists an element  $a \in I$  such that  $x - a \in J$ . Let  $\vartheta_1, \dots, \vartheta_r$  be a minimal system of generators of  $J$ , such that  $\bar{\vartheta}_1, \dots, \bar{\vartheta}_r$  is a free basis of  $J/J\Lambda$ . Then  $x - a = \sum \lambda_i \vartheta_i$  and using (i) we get an element  $\lambda \notin \bigcup_{\mathfrak{p}} \mathfrak{p}$ ,  $\mathfrak{p} \in \text{Ass}(A)$  such that  $\lambda(x - a) = \sum \lambda \lambda_i \vartheta_i \in I$ . Hence  $\lambda(x - a) = \sum \lambda \lambda_i \vartheta_i \in I \cap J$ .

From (iii) we get an element  $\lambda' \notin \bigcup_{\mathfrak{p}} \mathfrak{p}$ ,  $\mathfrak{p} \in \text{Ass}(A)$  such that

$$\lambda \lambda'(x - a) = \sum (\lambda \lambda' \lambda_i) \vartheta_i \in IJ \subset J\Lambda.$$

From (ii) we deduce  $\lambda \lambda' \lambda_i \in \Lambda$  for every  $i$ , hence  $\lambda_i \in \Lambda$  for every  $i$ , hence  $x \in I + \Lambda J = I + J^2$ . Going on with the same type of argument we get  $x \in I + J^r$  for every  $r$ , hence  $x \in I$ .

Let us state a notation: if  $\alpha$  is an ideal of a ring  $R$ , we denote by  $\mathcal{Z}_\alpha(R)$  the set  $\{x \in R/\bar{x} \in \mathcal{Z}(R/\alpha)\}$  where  $\bar{\phantom{x}}$  denotes the reduction modulo  $\alpha$  and  $\mathcal{Z}(\cdot \cdot \cdot)$  means "zerodivisors of  $\dots$ ".

We are ready to prove the following

**THEOREM 10.** *Let  $R$  be a local ring,  $I, J, \Lambda = I + J$  ideals of  $R$  such that  $I \cap J = IJ$ . Then the following conditions are equivalent:*

- (1)  $G(I)$  and  $G(J)$  are free.
- (2)  $G(I) \otimes_R R/\Lambda$  and  $G(J) \otimes_R R/\Lambda$  are free and  $G(IR_\mathfrak{p})$  and  $G(JR_\mathfrak{p})$  are free for every  $\mathfrak{p} \in \text{Ass}(R/\Lambda)$ .
- (3)  $G(\Lambda)$  is free,  $G(IR_\mathfrak{p})$  and  $G(JR_\mathfrak{p})$  are free for every  $\mathfrak{p} \in \text{Ass}(R/\Lambda)$  and  $\mathcal{Z}_I(R)$ ,  $\mathcal{Z}_J(R)$  are contained in  $\mathcal{Z}_\Lambda(R)$ .

**PROOF.** (1) $\Rightarrow$ (3)  $G(\Lambda)$  free follows from Theorem 5.

Hence we have to prove that  $\mathcal{Z}_I(R) \subseteq \mathcal{Z}_\Lambda(R)$  (the same argument works for  $\mathcal{Z}_J(R)$ ).

If we call  $B = R/I$ ,  $\mathcal{Z}_I(R) \subseteq \mathcal{Z}_\Lambda(R)$  is equivalent to  $\mathcal{Z}_{(0)}(B) \subseteq \mathcal{Z}_{\bar{\Lambda}}(B)$  where  $\bar{\phantom{x}}$  denotes the reduction modulo  $I$ . Using Corollary 8 we get that  $G(\Lambda)$  is a free  $R/\Lambda \cong B/\bar{\Lambda}$ -module. Let now  $x, y$  be elements of  $B$  such that  $xy = 0$  and  $x \in \mathfrak{p} \in E_\mathfrak{p}$  where  $E = \text{Ass}(B/\bar{\Lambda})$ . Being  $G(\Lambda)$  free,  $\text{Ass}(B/\bar{\Lambda}^n) \subseteq \text{Ass}(B/\bar{\Lambda})$ , hence  $y \in \bigcap_n \bar{\Lambda}^n = (0)$ .

(2) $\Rightarrow$ (1) Let  $\vartheta_1, \dots, \vartheta_\alpha$  be a minimal system of generators of  $I^n$ , such that  $\bar{\vartheta}_1, \dots, \bar{\vartheta}_\alpha$  is a free basis of  $I^n/I^{n+1}$ .

Let  $0 \longrightarrow T \longrightarrow (R/I)^\alpha \xrightarrow{\pi} I^n/I^{n+1} \longrightarrow 0$  be the exact sequence of  $R/I$ -modules defined by  $\pi(e_i) = \bar{\vartheta}_i \in I^n/I^{n+1}$ .

Then we can apply Lemma 9, because (i) and (ii) are clearly satisfied and (iii) follows from Lemma 4.

In conclusion  $T = 0$  and  $G_n(I)$  is free. This works for every  $n$  and, in the same way, for  $J$ .

Since (1)  $\Rightarrow$  (2) is obvious, we only have to prove (3)  $\Rightarrow$  (1). Since  $I \cap J = IJ$ , by Proposition 2 we get

$$\Lambda/\Lambda^2 \simeq I/I\Lambda \oplus J/J\Lambda.$$

Let  $\vartheta_1, \dots, \vartheta_\alpha$  be a minimal system of generators of  $I$  such that  $\bar{\vartheta}_1, \dots, \bar{\vartheta}_\alpha$  is a free basis of  $I/I\Lambda$ .

Let  $0 \longrightarrow T \longrightarrow (R/I)^\alpha \longrightarrow I/I^2 \longrightarrow 0$  be the exact sequence of  $R/I$ -modules defined by  $\pi(e_i) = \bar{\vartheta}_i \in I/I^2$ .

By hypothesis  $T_{\mathfrak{p}} = 0$  for every  $\mathfrak{p} \in \text{Ass}(R/\Lambda)$ , hence, for every  $x \in T$  there exists  $\lambda \notin \mathcal{X}_\Lambda(R)$  such that  $\lambda x = 0$ . Since  $\mathcal{X}_I(R) \subseteq \mathcal{X}_\Lambda(R)$ , we get  $x = 0$ . Hence  $I/I^2$  and  $J/J^2$  (using the same argument) are free. Let us assume, by induction, that  $G_i(I), G_i(J)$  are free for  $i = 0, \dots, \nu - 1$ . Using Lemma 4 we get  $I^\lambda \cap J^\mu = I^\mu$  for  $\lambda + \mu \leq \nu + 1$  hence  $G_\nu(\Lambda) \simeq \bigoplus_{\lambda+\mu=\nu} G_{\lambda,\mu}(I; J)$  by Proposition 2.

In particular  $I^\nu/I^\nu\Lambda$  and  $J^\nu/J^\nu\Lambda$  are free  $R/\Lambda$ -modules.

Using the same argument as before we get that  $G_\nu(I)$  and  $G_\nu(J)$  are free.

#### REFERENCES

- [1] ACHILLES, SCHENZEL, VOGEL: Bemerkungen über normale Flachheit und normale Torsionsfreiheit und Anwendungen. (Preprint).
- [2] HERRMANN, SCHMIDT: Zur Transitivität der normalen Flachheit. *Invent. Math.* 28 (1975).
- [3] HERRMANN, SCHMIDT, VOGEL: *Theorie der normalen Flachheit*. Teuber-Texte zur Math. (1977).
- [4] HIRONAKA: Resolution of singularities of an algebraic variety over a field of characteristic zero. *Ann. of Math.* 79 (1964).
- [5] GROTHENDIECK: Éléments de géométrie algébrique I.H.E.S. Publ. Math. Paris 32 (1967).
- [6] ROBBIANO: An algebraic property of  $\mathbb{P}^1 \times \mathbb{P}^r$ . (In preparation).
- [7] ROBBIANO, VALLA: Primary powers of a prime ideal. *Pacific J. Math.* 63 (1976).
- [8] ROBBIANO, VALLA: On normal flatness and normal torsionfreeness. *J. of Algebra* 43 (1976).

(Oblatum 15-XI-1977)

Istituto Matematico  
dell'Università di Genova  
Via L.B. Alberti, n° 4 -  
16132 GENOVA (Italia)