

COMPOSITIO MATHEMATICA

J. H. RUBINSTEIN

C. GARDINER

A note on a 3-dimensional homogeneous space

Compositio Mathematica, tome 39, n° 3 (1979), p. 297-299

http://www.numdam.org/item?id=CM_1979__39_3_297_0

© Foundation Compositio Mathematica, 1979, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A NOTE ON A 3-DIMENSIONAL HOMOGENEOUS SPACE

J.H. Rubinstein* and C. Gardiner

We give a calculation of the homeomorphism type of the homogeneous space $SL(2, R)/SL(2, Z)$ using only elementary properties of Seifert manifolds (cf. p. 84 of Milnor's book, Introduction to algebraic K-theory, for a different proof). As a corollary, the well-known presentation of $SL(2, Z)$ follows.

THEOREM: *The (right) coset space of $SL(2, Z)$ in $SL(2, R)$ is homeomorphic to the complement of the trefoil knot in S^3 .*

PROOF: Since $SL(2, Z)$ is a discrete subgroup of $SL(2, R)$, the coset space which we denote by M is a 3-manifold without boundary. Right multiplication of $SO(2, R)$ on $SL(2, R)$ commutes with left multiplication by $SL(2, Z)$. Therefore there is an induced action of $SO(2, R)$ on M . We show that M is a Seifert manifold with two exceptional fibres of multiplicity 2 and 3 and with orbit surface equal to an open disk. This implies that M and the complement of the trefoil knot in S^3 have the same Seifert invariants and so are homeomorphic (see [2] or [3] for the properties of Seifert spaces).

Let $A \in SO(2, R)$. Then A is in the stabilizer of some point of M exactly when the equation $\Sigma S = SA$ can be solved for $\Sigma \in SL(2, Z)$ and $S \in SL(2, R)$. In this case $\Sigma = SAS^{-1}$ and $\text{tr } A = \text{tr } \Sigma$ is an integer. If λ, γ are the eigenvalues of A , then $|\lambda| = |\gamma| = 1$ since A is orthogonal, and $\lambda\gamma = 1$ as $\det A = 1$. Therefore $\lambda = \bar{\gamma}$ and because $\lambda + \gamma$ is an integer we see that λ is an n^{th} root of unity for $n = 1, 2, 3, 4$ or 6 . Consequently $A^n = I$. Also $-I$ acts trivially on M and so we conclude that the stabilizer of every point of M is Z_2, Z_4 or Z_6 .

Consider the case when A is of order 4, i.e. when $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

* The first author was supported by a Rothman's Research Fellowship.

Observe that $S^t = AS^{-1}A^{-1}$, so that $SS^t = SAS^{-1}A^{-1} = \Sigma A^{-1} \in SL(2, Z)$. Let $SS^t = \begin{pmatrix} m & p \\ p & n \end{pmatrix}$ and let $B = \begin{pmatrix} 0 & 1 \\ -1 & \pm 1 \end{pmatrix}$ or $\begin{pmatrix} \pm 1 & 1 \\ -1 & 0 \end{pmatrix}$. Then BSS^tB^t has its off-diagonal entry equal to $\pm n - p$ or $\pm m - p$ respectively. Now $mn - p^2 = 1$ and $m > 0, n > 0$. Consequently either $m \leq |p|$ or $n \leq |p|$, and by appropriate choice of B we get that the off-diagonal term of BSS^tB^t has absolute value strictly less than $|p|$. So by a sequence of such transformations, a matrix $\Sigma_1 \in SL(2, Z)$ can be found for which $\Sigma_1 SS^t \Sigma_1^t = I$. This implies that $\Sigma_1 S \in SO(2, R)$ and the coset of S is in the same orbit of $SO(2, R)$ as the coset of I . Therefore there is only one orbit of $SO(2, R)$ in M with stabilizer Z_4 .

For the case when the stabilizer is Z_6 , let A be the matrix of rotation by $\pi/3$. Then $SAS^{-1} \in SL(2, Z)$ if and only if $\sqrt{3}SS^t$ is a matrix with integer entries, determinant equal to 3, odd integers on the diagonal and even integers off the diagonal. By the same argument as above, we can find a matrix $\Sigma_1 \in SL(2, Z)$ so that $\Sigma_1 \sqrt{3}SS^t \Sigma_1^t = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}$. Therefore $\Sigma_1 SS^t \Sigma_1^t = S_1 S_1^t$ where $S_1 = \begin{pmatrix} \sqrt[3]{3} & 0 \\ 0 & -\sqrt[3]{3} \end{pmatrix}$. This implies $S_1^{-1} \Sigma_1 S \in SO(2, R)$ and the coset of S is in the same orbit as the coset of S_1 . Consequently, there is just one orbit with stabilizer Z_6 .

To complete the proof we have to determine the orbit surface F of M . Every matrix in $SL(2, R)$ can be written uniquely as the product of a real lower triangular matrix and a matrix in $SO(2, R)$. So there is a homeomorphism of $SL(2, R)$ onto $R^2 \times S^1$ and the inclusion of $SO(2, R)$ in $SL(2, R)$ induces an isomorphism of fundamental groups. Since $M = SL(2, R)/SL(2, Z)$, there is an exact sequence $1 \rightarrow Z \rightarrow \pi_1(M) \rightarrow SL(2, Z) \rightarrow 1$, and $SL(2, Z)$ is the quotient of $\pi_1(M)$ by a cyclic normal subgroup with generator α . Note that α is the image of a generator of $\pi_1(SO(2, R))$ under the mapping $SO(2, R) \rightarrow SL(2, R) \rightarrow M$. Let h be the homotopy class of an orbit in M of the $SO(2, R)$ action, with stabilizer Z_2 (i.e. an ordinary fibre in the terminology of [3]). Without loss of generality $\alpha = h^2$ and $SL(2, Z)$ is the quotient of $\pi_1(M)$ by the cyclic normal subgroup generated by h^2 .

Consequently there is an epimorphism from $SL(2, Z)$ to $\pi_1(F)$ and also from $H_1(SL(2, Z))$ to $H_1(F)$. Now it is well-known that $SL(2, Z)$ is generated by $X = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $Y = \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$. So since $X^4 = 1, Y^6 = 1$ and $X^2 = Y^3 = -I$, we see that $H_1(SL(2, Z))$ has order at most 12. Therefore F must be an open disk, a 2-sphere or a projective plane.

If $F = S^2$ then M is a lens space and so cannot be covered by $SL(2, R)$. Suppose next that $F = RP^2$. As M is orientable, the relation

matrix for $H_1(M)$ (see [2] or [3]) shows that h^2 is null homologous and $H_1(M)$ has order at least 24. But $h^2 \sim 0$ implies $H_1(M) = H_1(SL(2, Z))$, which gives a contradiction. This completes the proof.

COROLLARY: $SL(2, Z) = Z_4 * z_2 Z_6$.

PROOF: $\pi_1(M)$ has generators h, x, y and relations $[h, x] = [h, y] = 1, x^2 h = y^3 h = 1$ (see [2] or [3]). Then $SL(2, Z)$ is given by adding the relation $h^2 = 1$ to $\pi_1(M)$, and so has the presentation $\{x, y \mid x^4 = 1, x^2 = y^3\}$.

REFERENCES

- [1] J. MILNOR: Introduction to algebraic K-theory. *Annals of Math. Studies* #72, Princeton, 1971. MR50, #2304.
- [2] P. ORLIK: Seifert manifolds. *Lecture Notes in Math.* #291, Springer, 1972.
- [3] H. SEIFERT: Topologie dreidimensionaler gefaserner Räume. *Acta math.* 60 (1933) 147–238.

(Oblatum 3-VIII-1977 & 20-X-1978)

Department of Mathematics
University of Melbourne
Parkville, Victoria 3052
Australia