

COMPOSITIO MATHEMATICA

HANS KLEPPE

STEVEN L. KLEIMAN

Reducibility of the compactified jacobian

Compositio Mathematica, tome 43, n° 2 (1981), p. 277-280

http://www.numdam.org/item?id=CM_1981__43_2_277_0

© Foundation Compositio Mathematica, 1981, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

REDUCIBILITY OF THE COMPACTIFIED JACOBIAN

Hans Kleppe* and Steven L. Kleiman**

Let X be an integral, projective curve of arithmetic genus p over an algebraically closed ground field k . Denote by P the compactified jacobian of X , defined as the moduli space of torsion-free sheaves on X with rank 1 and Euler characteristic $1 - p$. Altman, Iarrobino and Kleiman proved an irreducibility theorem [1, Theorem (9)]: P is irreducible if X lies on a smooth surface, or equivalently, if the embedding dimension at each point of X is at most two [3, Corollary (9)]. They also constructed an example [1, Example (13)] of an X which is a complete intersection in \mathbb{P}^3 and for which P is reducible. The example suggests that the converse of the theorem holds. In the present article, we prove the converse in the following form.

THEOREM (1): *If X does not lie on a smooth surface, then the compactified jacobian P is reducible.*

Rego [5] asserted Theorem (1) and offered a sketchy proof, which runs as follows. First he showed that $\text{Hilb}^2(X)$ is reducible if X does not lie on a smooth surface. Then, if X is also Gorenstein, he concluded that P is reducible from the fact that the Abel map, $\text{Hilb}^n(X) \rightarrow P$, is smooth for large n . This map is no longer smooth if X is not Gorenstein, and so Rego devised other methods to obtain reducibility in general.

However, Altman and Kleiman [2] developed a theory in which $\text{Quot}^n(\omega/X)$, where ω is the dualizing sheaf on X , replaces $\text{Hilb}^n(X)$ as the source of an Abel map,

$$A_\omega^n : \text{Quot}^n(\omega/X) \rightarrow P.$$

* Supported by the Norwegian Research Council for Science and the Humanities.

** J.S. Guggenheim Fellow.

Whether or not X is Gorenstein, A_ω^n is smooth and its fibers are projective spaces for all $n \geq 2p - 1$ [2, Theorem (8.4) (v), Lemma (5.17) (ii) and Theorem (4.2)]. Hence, P will be reducible if $\text{Quot}^n(\omega/X)$ is reducible for large n .

This reducibility is proved below in two steps. First, we show that, if $\text{Quot}^m(\omega/X)$ is reducible, then $\text{Quot}^n(\omega/X)$ is reducible for $n \geq m$ (Proposition (3)). Secondly, we show that, if X does not lie on a smooth surface, then $\text{Quot}^d(\omega/X)$ is reducible, for small d , in fact, for $d = 2$ if X is Gorenstein, and for $d = 1$ if X is not Gorenstein (Proposition (4)). Thus, by a natural adaptation of part of Rego's method, Theorem (1) is proved.

Fix a torsion-free, rank-1 sheaf \mathcal{F} on X . Denote by U the open subscheme of X consisting of nonsingular points.

The open subscheme $Q^n U$ of $\text{Quot}^n(\mathcal{F}/X)$ parameterizing quotients of \mathcal{F} with support contained in U is isomorphic to $\text{Hilb}^n(U)$, because \mathcal{F} restricted to U is invertible; so $Q^n U$ is irreducible of dimension n [1, Lemma (1)]. Hence, $\text{Quot}^n(\mathcal{F}/X)$ is irreducible if and only if $Q^n U$ is dense in $\text{Quot}^n(\mathcal{F}/X)$. Using the valuative criterion [4, Ch. II, Prop. 7.1.4 (i)], we therefore get Lemma (2) below.

LEMMA (2): *$\text{Quot}^n(\mathcal{F}/X)$ is irreducible if and only if, for all quotients F of \mathcal{F} of length n , there exists a scheme $T = \text{Spec}(A)$, where A is a complete, discrete valuation ring, and a T -flat quotient \bar{F} of \mathcal{F}_T such that $\bar{F}(t) \simeq F$ and $\text{Supp } \bar{F}(g) \subseteq U_T(g)$, where t and g denote the closed and generic points of T .*

PROPOSITION (3): *If $\text{Quot}^n(\mathcal{F}/X)$ is irreducible, then $\text{Quot}^m(\mathcal{F}/X)$ is irreducible for all $m < n$.*

PROOF: Let F be a quotient of \mathcal{F} of length m . Let I denote the kernel of the natural map $\mathcal{F} \rightarrow F$ and let x_1, \dots, x_{n-m} be different nonsingular points on X such that $x_i \notin \text{Supp } F$ for $i = 1, \dots, n - m$. Then

$$F' = \mathcal{F}/M_1 \dots M_{n-m}I,$$

where M_i denotes the ideal of x_i , is a quotient of \mathcal{F} of length n . By Lemma (2) there exists a complete, discrete valuation ring A and a quotient \bar{F}' of \mathcal{F}_T , $T = \text{Spec}(A)$, with all the properties listed in that lemma and such that $\bar{F}'(t) \simeq F'$.

Let W be the closed subscheme of X_T defined by the annihilator of \bar{F}' . It is easy to see that we have an inclusion

$$\{x_1\} \cup \dots \cup \{x_{n-m}\} \cup V \subseteq W(t),$$

where V is the closed subscheme of X defined by the annihilator of F . Hence, since A is a henselian ring [4, Ch. IV, Prop. 18.5.14], W may be written in the form,

$$W = W_1 \oplus \cdots \oplus W_{n-m} \oplus W',$$

where $\{x_i\} \subseteq W_i(t)$ and $V \subseteq W'(t)$ [4, Ch. IV, Théorème 18.5.11(c)].

Denote by i the inclusion $W' \subseteq X_T$ and put

$$\bar{F} = i_* i^* \bar{F}'.$$

Then \bar{F} is a flat quotient of \bar{F}' and $\bar{F}(t) \simeq F$. Hence, by Lemma (2) the proposition is proved.

PROPOSITION (4): *Let x be a point of X and denote by M the ideal defining x .*

(a) *If $\dim_k(\omega/M\omega) \geq 2$, then $\text{Quot}^1(\omega/X)$ is reducible.*

(b) *If $\dim_k(\omega/M\omega) = 1$ and if $\dim_k(M/M^2) \geq 3$, then $\text{Quot}^2(\omega/X)$ is reducible.*

PROOF: (a). Set $\omega_1 = \omega/M\omega$. Obviously the functors $\underline{\text{Quot}}^1(\omega_1/X)$ and $\underline{\text{Grass}}_1(\omega_1/k)$ are isomorphic. Since $\dim_k(\omega_1) \geq 2$, $\underline{\text{Grass}}_1(\omega_1/k)$ has dimension at least 1. Hence, since $\text{Quot}^1(\omega_1/X)$ is a closed subscheme of $\text{Quot}^1(\omega/X)$, we therefore get

$$\dim \text{Quot}^1(\omega/X) \geq 1.$$

If equality holds, $\text{Quot}^1(\omega/X)$ is reducible since $\text{Quot}^1(\omega_1/X)$ is a closed 1-dimensional subscheme which is obviously different from $\text{Quot}^1(\omega/X)$. If equality fails, then the closure of Q^1U is a component of dimension 1, and so $\text{Quot}^1(\omega/X)$ is reducible.

(b) ω is torsion-free [2, 6.5], so ω is invertible at x because $\dim_k(\omega/M\omega) = 1$. Since $\dim_k(M/M^2) \geq 3$, we get that

$$\dim_k(M\omega/M^2\omega) \geq 3.$$

Set $\omega_2 = \omega/M^2\omega$. A vector subspace of $M\omega/M^2\omega$ of codimension 1 corresponds to a quotient of ω_2 of length 2. It is not hard to see that this correspondence extends to families of quotients and vector subspaces, so that $\underline{\text{Grass}}_1(M\omega/M^2\omega)$ can be considered as a subfunctor of $\underline{\text{Quot}}^2(\omega_2/X)$. Hence, since a proper monomorphism is a closed embedding [4, Ch. IV, Prop. 8.11.5], $\text{Quot}^2(\omega_2/X)$ contains

$\text{Grass}_1(M\omega/M^2\omega)$. Since the latter has dimension at least 2, reasoning as in the proof of (a) we conclude that $\text{Quot}^2(\omega/X)$ is reducible.

REFERENCES

- [1] ALTMAN, IARROBINO and KLEIMAN: Irreducibility of the Compactified Jacobian. Nordic Summer School NAVF, Oslo 1976, Noordhoff (1977).
- [2] ALTMAN and KLEIMAN: Compactifying the Picard Scheme. *Advances in Mathematics*, 35 (1980) 50–112.
- [3] ALTMAN and KLEIMAN: Bertini Theorems for Hypersurface Sections Containing a Subscheme. *Communications in Algebra* 7(8) (1979) 775–790.
- [4] GROTHENDIECK and DIEUDONNÉ: Elements de Géométrie Algébrique. *Publ. Math. I.H.E.S.*, Nos. 8, 28, 32.
- [5] REGO: *The Compactified Jacobian*, revised and expanded version, Tata Institute, Preprint, Feb. 1978.

(Oblatum 15-VII-1980 & 12-X-1980)

M.I.T. 2-278
Cambridge, MA 02139
USA