

# COMPOSITIO MATHEMATICA

PAOLO DE BARTOLOMEIS

GIUSEPPE TOMASSINI

## **Traces of pluriharmonic functions**

*Compositio Mathematica*, tome 44, n° 1-3 (1981), p. 29-39

[http://www.numdam.org/item?id=CM\\_1981\\_\\_44\\_1-3\\_29\\_0](http://www.numdam.org/item?id=CM_1981__44_1-3_29_0)

© Foundation Compositio Mathematica, 1981, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## TRACES OF PLURIHARMONIC FUNCTIONS

Paolo de Bartolomeis and Giuseppe Tomassini

### Introduction

Let  $M$  be a real oriented hypersurface in a complex manifold  $X$ , which divides  $X$  into two open sets  $X^+$  and  $X^-$ .

In this paper we characterize in terms of tangential linear differential operators on  $M$  the distributions  $T$  on  $M$  which are "jumps" or traces (in the sense of currents) of pluriharmonic functions in  $X^+$  and  $X^-$ .

The starting point of our investigation is the non-tangential characterizing equation  $\bar{\partial}_b \partial T = 0$ , which can be deduced from the theory of boundary values of holomorphic forms. If  $M$  is not Levi-flat, we construct a second order tangential local linear differential operator  $\omega_M$  such that if  $T$  is the trace on  $M$  of a pluriharmonic function  $h$ , then  $\omega_M(T) = \partial h$ . This enables us to prove that the tangential equation  $\bar{\partial}_b \omega_M(T) = 0$  characterizes locally the traces on  $M$  of pluriharmonic functions on  $X^+$  or  $X^-$  (local Cauchy-Dirichlet problem). From this local result we deduce directly the global solvability of the Cauchy-Dirichlet problem in the case  $M$  is either compact or its Levi form has everywhere at least one positive eigenvalue.

Finally, using standard cohomological arguments the global Riemann-Hilbert problem ("jumps" of pluriharmonic functions on  $X \setminus M$ ) is solved if  $H^2(X, \mathbb{C}) = 0$  or  $H^1(M, \mathbb{R}) = 0$ . Particular cases of our problem have been investigated in [1], [3] (cf. also [2], [4]).

The present paper contains an improved version of the results announced in [5].

### 1. Preliminaries and notations

In the present paper  $X$  will be a complex manifold of dimension  $n \geq 2$ , and  $M \subset X$  a real oriented connected  $C^\infty$  hypersurface.

We assume  $M$  is defined by  $\rho = 0$  where  $\rho: X \rightarrow \mathbb{R}$  is a  $C^\infty$  function such that  $d\rho \neq 0$  on  $M$ .

We say that such a  $\rho$  is a *defining function* for  $M$ ;  $M$  divides  $X$  into two open sets  $X^+$  and  $X^-$ , defined respectively by  $\rho > 0$  and  $\rho < 0$ ; if  $U$  is an open subset of  $X$ , we will set  $U^\pm = U \cap X^\pm$ ; we can also assume that there exists an  $\epsilon_0 > 0$  such that if  $|\epsilon| < \epsilon_0$  and  $M_\epsilon$  is the level hypersurface defined by  $\rho = \epsilon$ , there exists a diffeomorphism  $\pi_\epsilon: M_\epsilon \rightarrow M$ .

We will use the standard notations for currents and distributions spaces (cf. e.g. [9]); in particular we fix the orientation on  $M$  in such a way that  $d[X^+] = [M]$ .

Furthermore we list the following definitions:

- i) Let  $L: \mathcal{G}^{(r)}(X) \rightarrow \mathcal{G}^{(r)}(M)$  be the restriction operator; we set:  $\mathcal{G}^{(p,q)}(M) = L(\mathcal{G}^{(p,q)}(X))$ .
- ii) Let  $K \in \mathcal{D}'^{(r)}(M)$ :  $K \wedge [M]$  will be the  $(r+1)$ -current on  $X$  defined by:  $\langle K \wedge [M], \varphi \rangle = \langle K, L(\varphi) \rangle$ ,  $\varphi \in \mathcal{D}^{(2n-r-1)}(X)$ .
- iii) We say that  $K \in \mathcal{D}'^{(r)}(M)$  is a  $(r, 0)$ -current (resp.  $(0, r)$ -current) if  $(K \wedge [M])^{p,q} = 0$  for  $p \leq r$  (resp.  $(K \wedge [M])^{q,p} = 0$  for  $p \leq r$ ); we denote by  $\mathcal{D}'^{(r,0)}(M)$  (resp.  $\mathcal{D}'^{(0,r)}(M)$ ) the space of  $(r, 0)$ -currents (resp.  $(0, r)$ -currents).
- iv) If  $K \in \mathcal{D}'^{(r,0)}(M)$ , then  $K \wedge [M]^{1,0}$  is the  $(r+1, 0)$ -current defined by:  $\langle K \wedge [M]^{1,0}, \varphi \rangle = \langle K, L(\varphi) \rangle$ ,  $\varphi \in \mathcal{D}^{(n-r-1,n)}(X)$  and  $K \wedge [M]^{0,1}$  is the  $(r, 1)$ -current defined by:  $\langle K \wedge [M]^{0,1}, \varphi \rangle = \langle K, L(\varphi) \rangle$ ,  $\varphi \in \mathcal{D}^{(n-r, n-1)}(X)$ .
- v) Let  $\alpha \in \mathcal{G}^{(2)}(X^+)$ ; we say that  $\alpha$  *admits trace*  $K \in \mathcal{D}'^{(r)}(M)$  on  $M$  in the sense of currents if for every  $\varphi \in \mathcal{D}^{(2n-r-1)}(M)$  we have:

$$\begin{aligned} \lim_{\epsilon \rightarrow 0^+} \int_{M_\epsilon} \alpha \wedge \pi_\epsilon^*(\varphi) &= \lim_{\epsilon \rightarrow 0^+} \int_M \pi_{\epsilon*}(\alpha) \wedge \varphi = \\ &= \lim_{\epsilon \rightarrow 0^+} \langle \pi_{\epsilon*}(\alpha), \varphi \rangle = \langle K, \varphi \rangle \end{aligned}$$

(cf. [8]); we set  $K = \gamma_+(\alpha)$ ; in the same manner we define  $\gamma_-(\alpha)$  if  $\alpha \in \mathcal{G}^{(r)}(X^-)$ .

Let  $a \in \mathcal{G}^{(r)}(X \setminus M)$  such that  $\gamma_+(\alpha)$  and  $\gamma_-(\alpha)$  exist: we refer to  $\gamma_+(\alpha) - \gamma_-(\alpha)$  as the *jump* of  $\alpha$  on  $M$ .

We denote by  $\mathcal{G}_*^{(r,s)}(X^\pm)$  the space of forms  $\alpha \in \mathcal{G}^{(r,s)}(X^\pm)$  such that  $\gamma_\pm(\alpha)$  and  $\gamma_\pm(d\alpha)$  exist.

Observe that if  $\alpha \in \mathcal{G}_*^{(r,0)}(X^+)$ , in particular we have

$$\partial(\gamma_+(\alpha) \wedge [M]^{1,0}) = (-1)^{r+1} \gamma_+(\partial\alpha) \wedge [M]^{1,0}.$$

## 2. Tangential operators on $M$

### 2a) The smooth case

A local linear operator  $\Phi: \mathcal{E}^{(p,q)}(X) \rightarrow \mathcal{E}^{(r,s)}(M)$  is said to be *tangential on  $M$*  if from  $L(f) = 0$  it follows  $\Phi(f) = 0$ ; a tangential operator induces a new operator:  $\mathcal{E}^{(p,q)}(M) \rightarrow \mathcal{E}^{(r,s)}(M)$  which will be denoted again by  $\Phi$ .

Let now  $(\cdot, \cdot)$  be a Hermitian structure on  $X$ ; without loss of generality we can assume  $(\partial\rho, \partial\rho) \equiv \frac{1}{2}$  on  $M$ . Define:

$$\begin{aligned} \mathcal{N}^{(p,q)}(M) &= \{\alpha \in \mathcal{E}^{(p,q)}(M) \mid \alpha = \varphi \wedge L(\partial\rho)\} \quad p \geq 1 \\ \mathfrak{E}^{(p,q)}(M) &= \{\alpha \in \mathcal{E}^{(p,q)}(M) \mid (\alpha, \beta) = 0 \quad \forall \beta \in \mathcal{N}^{(p,q)}(M)\} \\ \bar{\mathcal{N}}^{(p,q)}(M) &= \{\alpha \in \mathcal{E}^{(p,q)}(M) \mid \alpha = \varphi \wedge L(\bar{\partial}\rho)\} \quad q \geq 1 \\ \bar{\mathfrak{E}}^{(p,q)}(M) &= \{\alpha \in \mathcal{E}^{(p,q)}(M) \mid (\alpha, \beta) = 0 \quad \forall \beta \in \bar{\mathcal{N}}^{(p,q)}(M)\} \end{aligned}$$

We have the decompositions:

$$\begin{aligned} \mathcal{E}^{(p,q)}(M) &= \mathcal{N}^{(p,q)}(M) \oplus \mathfrak{E}^{(p,q)}(M) \\ \mathcal{E}^{(p,q)}(M) &= \bar{\mathcal{N}}^{(p,q)}(M) \oplus \bar{\mathfrak{E}}^{(p,q)}(M) \end{aligned}$$

and we denote by:

$$\begin{aligned} \tau: \mathcal{E}^{(p,q)}(M) &\rightarrow \mathfrak{E}^{(p,q)}(M) \\ \bar{\tau}: \mathcal{E}^{(p,q)}(M) &\rightarrow \bar{\mathfrak{E}}^{(p,q)}(M) \end{aligned}$$

the natural projections; observe that  $\tau(\alpha) = \tau(\beta)$  is equivalent to  $\alpha \wedge [M]^{1,0} = \beta \wedge [M]^{1,0}$ ; we set  $\partial_b = \tau \circ L \circ \partial$  and  $\bar{\partial}_b = \bar{\tau} \circ L \circ \bar{\partial}$ ; by definition  $\partial_b$  and  $\bar{\partial}_b$  are tangential operators on  $M$  (cf. [6]).

The induced operators

$$\begin{aligned} \partial_b: \mathcal{E}^{(p,q)}(M) &\rightarrow \mathfrak{E}^{(p+1,q)}(M) \\ \bar{\partial}_b: \mathcal{E}^{(p,q)}(M) &\rightarrow \bar{\mathfrak{E}}^{(p,q+1)}(M) \end{aligned}$$

are described by the formulas

$$\begin{aligned} \partial_b \alpha &= \tau \circ L \circ \partial(\alpha \wedge [M]^{1,0}) \\ \bar{\partial}_b \alpha &= \bar{\tau} \circ L \circ \bar{\partial}(\alpha \wedge [M]^{0,1}). \end{aligned}$$

Thus

$$\begin{aligned}\partial(\alpha \wedge [M]^{1,0}) &= \partial_b \alpha \wedge [M]^{1,0} \\ \bar{\partial}(\alpha \wedge [M]^{0,1}) &= \bar{\partial}_b \alpha \wedge [M]^{0,1}.\end{aligned}$$

We consider in particular the following cases:  $f \in \mathcal{E}^{(0,0)}(X)$  and  $\beta \in \mathcal{E}^{(1,0)}(X)$ .

a) if  $f \in \mathcal{E}^{(0,0)}(X)$  set:

$$N(f) = L[(\partial f, \partial \rho)] \quad \bar{N}(f) = L[(\bar{\partial} f, \bar{\partial} \rho)];$$

so we obtain on  $M$ :

$$\begin{aligned}L(\partial f) &= \partial_b f + 2N(f)L(\partial \rho) \\ L(\bar{\partial} f) &= \bar{\partial}_b f + 2\bar{N}(f)L(\bar{\partial} \rho);\end{aligned}$$

furthermore it is easy to check that:

- i) at every point of  $M$ ,  $N(f)$  represents the complex normal derivative of  $f$  and  $N(f) + \bar{N}(f)$  is the real normal derivative of  $f$ .
- ii)  $f \mapsto N(f) - \bar{N}(f)$  is a tangential operator on  $M$ ; thus we have also e.g.:

$$L(\partial f) = \partial_b f + [N(f) - \bar{N}(f)]L(\partial \rho) + [N(f) + \bar{N}(f)]L(\partial \rho)$$

b) if  $\beta \in \mathcal{E}^{(1,0)}(X)$  define  $N_1(\beta)$  and  $\bar{N}_1(\beta)$  by the relations:

$$\begin{aligned}L(\partial \beta) &= \partial_b \beta + 2N_1(\beta) \wedge L(\partial \rho) \\ L(\partial \bar{\beta} + \bar{\partial} \beta) - \bar{\tau}(\partial \bar{\beta} + \bar{\partial} \beta) &= \bar{N}_1(\beta) \wedge L(\partial \rho)\end{aligned}$$

we observe that  $\beta \mapsto N_1(\beta) - \bar{N}_1(\beta)$  is a tangential operator, and so we obtain on  $M$  the decomposition:

$$L(\partial \beta) = \partial_b \beta + [N_1(\beta) - \bar{N}_1(\beta)] \wedge L(\partial \rho) + [N_1(\beta) + \bar{N}_1(\beta)] \wedge L(\partial \rho)$$

which again enables us to isolate the genuine non-tangential component of  $L(\partial \beta)$ , namely  $(N_1(\beta) + \bar{N}_1(\beta)) \wedge L(\partial \rho)$ .

Let  $U \subset X$  be an open set and let  $\mathcal{P}(U)$  be the space of real

pluriharmonic functions on  $U$  (i.e.  $f \in \mathcal{P}(U)$  iff  $f$  is real valued and  $\partial\bar{\partial}f = 0$ ).

We have the following:

**PROPOSITION 2.1:** *Assume  $L(\partial\rho \wedge \bar{\partial}\rho \wedge \partial\bar{\partial}\rho) \neq 0$ ; then there exists a local linear differential operator  $R: \mathcal{E}^{(0,0)}(M) \rightarrow \mathcal{E}^{(0,0)}(M)$  such that if  $h \in \mathcal{P}(X)$  then  $R(L(h)) = L(N(h) + \bar{N}(h))$ .*

**PROOF:** Let  $h \in \mathcal{P}(X)$ : define  $\delta_b = \partial_b + \bar{\partial}_b$ ,  $^*\delta_b = \bar{\tau}\partial_b + \tau\bar{\partial}_b$ ,  $\delta_b^\natural = i(\bar{\partial}_b - \partial_b)$ ; then omitting  $L$  to simplify our notations:

$$\begin{aligned} 0 &= \tau\bar{\tau}(\bar{\partial}\partial h) = \tau(\bar{\partial}_b\partial h) = \tau(\bar{\partial}_b[\partial_b h + 2N(h)\partial\rho]) = \\ &= \tau(\bar{\partial}_b\partial_b h + 2\bar{\partial}_b N(h) \wedge \partial\rho + 2N(h)\bar{\partial}_b\partial\rho) = \tau(\partial_b\partial_b h) + 2N(h)\tau(\bar{\partial}_b\partial\rho) \end{aligned}$$

and also

$$0 = \bar{\tau}\tau(\partial\bar{\partial}h) = \tau(\partial_b\bar{\partial}_b h) + 2N(h)\bar{\tau}(\partial_b\bar{\partial}\rho) = \bar{\tau}(\partial_b\bar{\partial}_b h) - 2N(h)\tau(\bar{\partial}_b\partial\rho)$$

thus we obtain the relation:

$$(\neq \neq) \quad ^*\delta_b\delta_b^\natural h = 2i[N(h) + \bar{N}(h)]\tau(\bar{\partial}_b\partial\rho);$$

taking the Hermitian product, we have:

$$(^*\delta_b\delta_b^\natural h, i\tau(\bar{\partial}_b\partial\rho)) = 2[N(h) + \bar{N}(h)](\|i\tau(\bar{\partial}_b\partial\rho)\|^2).$$

Since  $i\tau(\bar{\partial}_b\partial\rho)$ , which is a real operator, represents the restriction to  $M$  of the Levi form of  $\rho$ , in our assumption  $\|i\tau(\bar{\partial}_b\partial\rho)\|^2 > 0$  everywhere and so the  $R$  we are looking for is given by:

$$R(f) = \frac{1}{2}[(^*\delta_b\delta_b^\natural f, i\tau(\bar{\partial}_b\partial\rho))\|i\tau(\bar{\partial}_b\partial\rho)\|^{-2}].$$

We observe that in [10] a similar formula is proved in a more laborious way.

For example in the case  $X = B^2$ , the unit ball in  $C^2$ , and  $M = bB^2$ , we obtain the formula:

$$\begin{aligned} R(f) &= \frac{1}{\sqrt{2}} \left[ \frac{\partial f}{\partial z_1} z_1 + \frac{\partial f}{\partial z_2} z_2 + \frac{\partial f}{\partial \bar{z}_1} \bar{z}_1 + \frac{\partial f}{\partial \bar{z}_2} \bar{z}_2 \right. \\ &\quad \left. + 2 \left( \frac{\partial^2 f}{\partial z_1 \partial \bar{z}_2} z_1 \bar{z}_2 + \frac{\partial^2 f}{\partial \bar{z}_1 \partial z_2} z_1 \bar{z}_2 - \frac{\partial^2 f}{\partial z_1 \partial \bar{z}_1} z_2 \bar{z}_2 - \frac{\partial^2 f}{\partial z_2 \partial \bar{z}_2} z_1 \bar{z}_1 \right) \right] \end{aligned}$$

(cf. also [1]).

Proposition 2.1 shows that if  $h$  is a pluriharmonic function on  $X$  and  $M$  is not Levi-flat anywhere, then the real normal derivative of  $h$  on  $M$  can be expressed by means of a real tangential operator  $R$ .

Thus we have on  $M$ :

$$(*) \quad L(\partial h) = \partial_b h + [N(h) - \bar{N}(h)]L(\partial\rho) + R(h)L(\partial\rho)$$

let  $\omega_M: \mathcal{E}^{(0,0)}(M) \rightarrow \mathcal{E}^{(1,0)}(M)$  be defined by the right member of (\*) we have the following:

REMARKS 2.2: a) if  $f \in \mathcal{E}^{(0,0)}(M)$ , then  $\partial_b \omega_M(f) = 0$ .

(b) if  $f$  is the restriction of a pluriharmonic function  $F$  then  $\bar{\partial}_b \omega_M(f) = 0$ .

PROOF: a) Let  $\tilde{f} \in \mathcal{E}^{(0,0)}(X^+)$  be an extension of  $f$ ; by Stokes' theorem,  $L(\partial\rho) \wedge [M]^{1,0} = 0$ , so  $\omega_M(f) \wedge [M]^{1,0} = \partial_b f \wedge [M]^{1,0} = \partial(\tilde{f}[M]^{1,0})$  and therefore  $\partial\omega_M(f) \wedge [M]^{1,0} = 0$ , so  $\partial_b \omega_M(f) = 0$ .

b) By (\*),  $\omega_M(f) = L(\partial F)$  so  $\omega_M(f) \wedge [M]^{0,1} = \partial F \wedge [M]^{0,1}$ . Since  $F$  is pluriharmonic,  $\bar{\partial}(\omega_M(f) \wedge [M]^{0,1}) = 0$  and therefore  $\bar{\partial}_b \omega_M(f) = 0$ .

2b) *Extension to the general case.*

Consider the dual decompositions:

$$\mathcal{D}^{(p,q)}(M) = \mathcal{N}'^{(p,q)}(M) \oplus \mathfrak{F}'^{(p,q)}(M)$$

$$\mathcal{D}'^{(p,q)}(M) = \bar{\mathcal{N}}'^{(p,q)}(M) \oplus \bar{\mathfrak{F}}'^{(p,q)}(M)$$

where the projections  $\tau$  and  $\bar{\tau}$  are defined in an obvious way, so the operators  $\partial_b$  and  $\bar{\partial}_b$  extend naturally to currents on  $M$ . Furthermore it turns out that  $N_1$  and  $\bar{N}_1$  are defined as  $N_1, \bar{N}_1: \mathcal{E}_*^{1,0}(X^\pm) \rightarrow \mathcal{D}'^{(1,0)}(M)$  and they induce a continuous operator  $N_1 - \bar{N}_1: \mathcal{D}'^{(1,0)}(M) \rightarrow \mathcal{D}'^{(1,0)}(M)$ ; so if e.g.  $\beta \in \mathcal{E}_*^{(1,0)}(X^+)$  we have:

$$\gamma_+(\partial\beta) = \partial_b \beta + (N_1 - \bar{N}_1)(\gamma_+(\beta)) \wedge L(\partial\rho) + (N_1 + \bar{N}_1)(\beta) \wedge L(\partial\rho).$$

Of course  $\partial\beta = 0$  implies  $N_1(\beta) = 0$ ; we note also that if  $\beta$  is holomorphic or  $\beta = \partial f$  for a real valued  $f$ , then  $\bar{N}_1(\beta) = 0$ ; in particular if  $\gamma_+(\beta) = \gamma_+(f)\partial\rho$  for a real valued function  $f$ , then:

$$(\circ) \quad (N_1 - \bar{N}_1)(\beta) = (N_1 - \bar{N}_1)(\partial f\rho - \rho\partial f) = (N_1 - \bar{N}_1)(\partial f\rho) = 0.$$

Finally we have that  $\omega_M$  can be extended as an operator  $\omega_M: \mathcal{D}'^{(0,0)}(M) \rightarrow \mathcal{D}'^{(1,0)}(M)$  and remarks 2.2 a), b) hold.

### 3. Traces of pluriharmonic functions

We are able now to give the following local solution to the trace problem (Cauchy–Dirichlet problem) for the  $\bar{\partial}\partial$  operator.

**THEOREM 3.1:** *Let  $p \in M$  and  $U$  be a neighbourhood of  $p$ ; assume  $L(\partial\rho \wedge \bar{\partial}\rho \wedge \bar{\partial}\partial\rho) \neq 0$  on  $M \cap U$  and let  $T$  be a real distribution defined on  $U \cap M$ ; then the following statements are equivalent:*

- i)  $\bar{\partial}_b\omega_M(T) = 0$  on  $U \cap M$
- ii) *there exists a neighbourhood  $V$  of  $p$  and there exists  $F \in \mathcal{P}(V \setminus M)$  such that  $\gamma_+(F) - \gamma_-(F) = T$  in  $U \cap V \cap M$ ; more precisely if the Levi form of  $\rho$  has a positive (resp. negative) eigenvalue at  $p$  we can choose  $F|_{V^-} = 0$  (resp.  $F|_{V^+} = 0$ ) and so  $T$  is actually the trace of a pluriharmonic function.*

**PROOF:** We have already observed that ii) implies i); conversely assume  $\bar{\partial}_b\omega_M(T) = 0$  on  $U \cap M$ ; let  $W \subset U$  be a Stein neighbourhood of  $p$ : by assumption we have:  $\bar{\partial}[\omega_M(T) \wedge [M \cap W]^{0,1}] = 0$  in  $W$  and so there exists  $\tilde{K} \in \mathcal{D}'^{(1,0)}(W)$  such that  $\bar{\partial}\tilde{K} = \omega_M(T) \wedge [M \cap W]^{0,1}$ .

$\tilde{K}|_{W^+}$  and  $\tilde{K}|_{W^-}$  are holomorphic  $(1, 0)$ -forms in  $W^+$  and  $W^-$  respectively,  $\tilde{K}|_{W^\pm} \in \mathcal{E}_*^{(1,0)}(W^\pm)$  and  $\gamma_+(\tilde{K}|_{W^+}) - \gamma_-(\tilde{K}|_{W^-}) = \omega_M(T)$  on  $M \cap W$  (cf. [8] th.II 1.3). Now assume the Levi form of  $\rho$  has e.g. a positive eigenvalue at  $p$ ; then  $\tilde{K}|_{W^-}$  extends across  $M$  as a holomorphic  $(1,0)$ -form  $\beta$  to a Stein neighbourhood  $V$  of  $p$ : if  $K_+ = \tilde{K}|_{V^+}$  of course we have:  $\gamma_+(K_+ - \beta) = \omega_M(T)$  on  $M \cap W$ .

Furthermore we have:

$$(**) \quad \gamma_+[\partial(K_+ - \beta)] = 0$$

in fact  $\bar{N}_1(K_+ - \beta) = 0$  and so:

$$\begin{aligned} \gamma_+[\partial(K_+ - \beta)] &= \partial_b[\gamma_+(K_+ - \beta)] + 2(N_1 - \bar{N}_1)(\gamma_+(K_+ - \beta)) \\ &= \partial_b\omega_M(T) + 2(N_1 - \bar{N}_1)(\omega_M(T)) = 2(N_1 - \bar{N}_1)(\partial_b T) \end{aligned}$$

Let now  $(f_n)_{n \in \mathbb{N}}$  be a sequence of elements of  $\mathcal{E}^{(0,0)}(V)$  such that  $L(f_n) \rightarrow T$  in  $\mathcal{D}'^{(0,0)}(M \cap V)$ ; from (\*) it follows

$$2(N_1 - \bar{N}_1)(\partial_b T) = \lim_n 2(N_1 - \bar{N}_1)(\partial_b f_n) = \lim_n 2(N_1 - \bar{N}_1)(\partial f_n) = 0$$

which proves (\*\*).

Now in  $V$  we have:

$$\bar{\partial}[\partial(K_+ - \beta) \wedge [V^+]] = \gamma_+(\partial K_+ - \beta) \wedge [V \cap M]^{0,1} = 0$$

and thus  $\partial(K_+ - \beta) \wedge [V^+]$  is holomorphic in  $V$  and so  $\partial(K_+ - \beta) \equiv 0$  on  $V^+$ .

It follows that on  $V$ :  $\partial[(K_+ - \beta) \wedge [V^+] - T \wedge [M \cap V]^{1,0}] = 0$  and so it is possible to find  $\tilde{G} \in \mathcal{D}'^{(0,0)}(V)$  such that:

$$\partial \tilde{G} = (K_+ - \beta) \wedge [V^+] - T \wedge [M \cap V]^{1,0}.$$

Then if we set  $G = \tilde{G}|_{V \setminus M}$  we obtain  $\bar{\partial} \partial G = 0$ .

Thus we have the following:

- a)  $G$  is a pluriharmonic function in  $V \setminus M$
- b) Since  $G$  can be extended as a distribution across  $M$  and satisfies  $\bar{\partial} \partial G = 0$ , then (cf. again [8] corollaire I, 2.6.)  $\gamma_+(G)$  and  $\gamma_-(G)$  exist.
- c) On  $V^+$  one has  $\partial G = K_+ - \beta$ .

We have also that  $\gamma_+(G) + T$  is  $\partial_b$ -closed: in fact:

$$\begin{aligned} \partial[(\gamma_+(G) + T) \wedge [M \cap V]^{1,0}] &= \omega_M(T) \wedge [M \cap V]^{1,0} + \\ &- \gamma_+(\partial G) \wedge [M]^{1,0} = \omega_M(T) \wedge [M \cap V]^{1,0} - \gamma_+(K_+ - \beta) \wedge [M \cap V]^{1,0}. \end{aligned}$$

Thus there exists an antiholomorphic function  $H$  on  $V^+$  such that  $\gamma_+(H) = \gamma_+(G) + T$ .

It follows that  $F = H - G$  is a pluriharmonic function on  $V^+$  such that  $\gamma_+(F) = T$  and since  $T$  is real we actually have  $\gamma_+(\operatorname{Re} F) = T$  and the proof of Theorem 3.1. is complete.

From Theorem 3.1. we can deduce first the following global solutions of the Cauchy-Dirichlet problem:

**PROPOSITION 3.2:** *Suppose the Levi form of  $\rho$  has least one positive eigenvalue at every point  $p \in M$ . Then there exists a neighbourhood  $U$  of  $M$  such that the equation  $\bar{\partial}_b \omega_M(T) = 0$  characterizes those distributions on  $M$  which are traces of pluriharmonic functions in  $U^+$ .*

**PROOF:** Theorem 3.1. assures that there exists a covering  $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$  of  $M$  by open set of  $X$  such that for every  $n \in \mathbb{N}$  there exists  $f_n \in \mathcal{P}(U_n^+)$  for which  $\gamma_+(f_n) = T$  on  $M \cap U_n$ ; furthermore, if

$U_n \cap U_m \cap M \neq \emptyset$ , one has  $\gamma_+(f_m) = \gamma_+(f_n)$  on  $U_m \cap U_n \cap M$  and thus, since the trace on  $M$  characterizes a pluriharmonic function, we have  $f_m = f_n$  on  $U_n^+ \cap U_m^+$  etc ...

**PROPOSITION 3.3:** *Suppose  $X$  is a Stein manifold,  $X^+$  is relatively compact and  $L(\partial\rho \wedge \bar{\partial}\rho \wedge \bar{\partial}\partial\rho) \neq 0$ . If  $T$  is a real distribution on  $M$ , then the following statements are equivalent:*

- i)  $\bar{\partial}_b \omega_M(T) = 0$
- ii) *there exists  $F \in \mathcal{P}(X^+)$  such that  $\gamma_+(F) = T$*

**PROOF:** i) follows immediately from ii); assume now i) holds: in order to prove ii), we argue in the same way as in Theorem 3.1, setting  $W = X$  and using Hartog's theorem to extend  $\bar{K}|_{X^-}$  to the whole  $X$ .

Using standard cohomological arguments we can investigate the global Riemann–Hilbert problem.

Let  $\mathcal{S}$  be the sheaf of germs of real distributions  $T$  on  $M$  such that  $\bar{\partial}_b \omega_M(T) = 0$  and let  $\hat{\mathcal{S}}$  be its trivial extension to  $X$ . Let  $\mathcal{A}$  be the sheaf of germs of distributions  $T$  on  $M$  such that  $\bar{\partial}_b T = 0$ . Furthermore, let  $\mathcal{P}_X$  be the sheaf of germs of real pluriharmonic functions on  $X$  and  $*\mathcal{P}_M$  the sheaf on  $X$  associated to the canonical presheaf:

$$*\mathcal{P}(U) = \begin{cases} \mathcal{P}(U) & \text{if } U \cap M = \emptyset \\ \{f \in \mathcal{P}(U) \mid \gamma_+(f), \gamma_-(f) \text{ exist on } M\} & \text{if } U \cap M \neq \emptyset. \end{cases}$$

Assume  $L(\partial\rho \wedge \bar{\partial}\rho \wedge \bar{\partial}\partial\rho) \neq 0$  on  $M$ .

Let  $\text{Re}: \mathcal{A} \rightarrow \mathcal{S}$  be the sheaf homomorphism defined by  $T \mapsto$  real part of  $T$ , and let  $\alpha$  be the sheaf homomorphism defined by:  $\alpha: *\mathcal{P}_M \rightarrow \hat{\mathcal{S}}$

$$\alpha_x(f) = \begin{cases} [\gamma_+(f) - \gamma_-(f)]_x & \text{if } x \in M \\ 0 & \text{if } x \notin M \end{cases}.$$

**COROLLARY 3.4:** *The sequences*

$$(1) \quad 0 \rightarrow \mathcal{P}_X \rightarrow *\mathcal{P}_M \xrightarrow{\alpha} \hat{\mathcal{S}} \rightarrow 0$$

$$(2) \quad 0 \rightarrow \mathbb{R} \xrightarrow{i} \mathcal{A} \xrightarrow{\text{Re}} \mathcal{S} \rightarrow 0$$

*are exact.*

PROOF: We need the following

LEMMA 3.5: *Suppose  $M$  is not Levi-flat at  $p$  and let  $U$  be a neighbourhood of  $p$ ; let  $f_{\pm} \in \mathcal{P}(U^{\pm})$  admitting traces  $\gamma_+(f_+)$  and  $\gamma_-(f_-)$  on  $U \cap M$ . If furthermore  $\gamma_+(f_+) = \gamma_-(f_-)$ , then there exists  $f \in \mathcal{P}(U)$  such that  $f|_{U^{\pm}} = f_{\pm}$ .*

PROOF OF LEMMA 3.5: Since the problem is local, we can assume  $U$  is a domain of  $\mathbb{C}^n$ : then  $\frac{\partial f_{\pm}}{\partial z_j}$ ,  $1 \leq j \leq n$ , are holomorphic in  $U^{\pm}$ ,  $\gamma_+(\frac{\partial f_+}{\partial z_j})$  and  $\gamma_-(\frac{\partial f_-}{\partial z_j})$  exist and Proposition 2.1. assures that  $\gamma_+(\frac{\partial f_+}{\partial z_j}) = \gamma_-(\frac{\partial f_-}{\partial z_j})$ ,  $1 \leq j \leq n$ ; hence, we are essentially reduced to the case  $f_{\pm}$  holomorphic which follows from [8], Corollaire II 1.2.

PROOF OF THE COROLLARY 3.4: (1) In virtue of Theorem 3.1.  $\alpha$  is surjective and the previous lemma concludes the proof of the exactness of (1).

(2) As a consequence of Lemma 3.5, we deduce easily that  $\ker \operatorname{Re} = \operatorname{Im}(i)$  and so we have to check that if  $p \in M$  and  $T \in \mathcal{S}_p$  then there exists  $\tilde{T} \in \mathcal{A}_p$  such that  $\operatorname{Re} \tilde{T} = T$ .

Now in virtue of Theorem 3.1, there exist a neighbourhood  $U$  of  $p$  in  $X$  and  $F \in \mathcal{P}(U \setminus M)$  such that:  $T = \gamma_+(F) - \gamma_-(F)$ ; from [8] we deduce that there exists  $G \in \mathcal{O}(U \setminus M)$  such that  $\operatorname{Re} G = F$  and  $\gamma_+(G)$ ,  $\gamma_-(G)$  exist; (more in detail the argument runs as follows: holomorphic and pluriharmonic functions with traces in the sense of currents are characterized by finite order of growth with respect to  $\rho$  ([8] Corollaire I 2.6.) so  $F$  has finite order of growth with respect to  $\rho$  and so does  $G$ , which can be expressed locally as  $G = F + iH$ , where  $H$  satisfies  $dH = d^c F$  etc . . .); it follows that  $\operatorname{Re}: \mathcal{A} \rightarrow \mathcal{S}$  is surjective and (2) is exact: so the proof of Corollary 3.4 is complete.

THEOREM 3.6: *(Global solution of the Riemann–Hilbert problem for  $\bar{\partial}\bar{\partial}$ ) Suppose  $X$  is a Stein manifold and  $L(\partial\rho \wedge \bar{\partial}\rho \wedge \bar{\partial}\bar{\partial}\rho) \neq 0$ ; assume furthermore  $H^2(X, \mathbb{C}) = 0$  or  $H^1(M, \mathbb{R}) = 0$ ; then if  $T$  is a real distribution on  $M$ , the following statements are equivalent:*

- i)  $\bar{\partial}_b \omega_M(T) = 0$
- ii) *there exists  $F \in \mathcal{P}(X \setminus M)$  such that  $\gamma_+(F) - \gamma_-(F) = T$ .*

PROOF: We observe that for a Stein manifold  $X$  we have the isomorphism  $H^r(X, \mathcal{P}_X) \approx H^{r+1}(X, \mathbb{C})$  for  $r \geq 1$ ; moreover we have:  $H^0(X, \hat{\mathcal{P}}) \approx H^0(M, \mathcal{S})$ .

If  $H^2(X, \mathbb{C}) = 0$  we obtain the exact sequence:

$$0 \rightarrow H^0(X, \mathcal{P}_X) \rightarrow H^0(X, *_\mathcal{P}_M) \rightarrow H^0(M, \mathcal{S}) \rightarrow 0$$

If  $H^1(M, \mathbb{R}) = 0$  we obtain the exact sequence:

$$0 \rightarrow \mathbb{R} \rightarrow H^0(M, \mathcal{A}) \rightarrow H^0(M, \mathcal{S}) \rightarrow 0$$

This concludes the proof.

## REFERENCES

- [1] T. AUDIBERT: *Caractérisation locale par des opérateurs différentiels des restrictions à la sphère de  $\mathbb{C}^n$  des fonctions pluriharmoniques* C.R.A.S. (1977) A284 1029–1031.
- [2] E. BEDFORD:  $(\partial\bar{\partial})_b$  and the real part of CR functions *Indiana Univ. Math. J.* (1980) 29 3 333–340.
- [3] E. BEDFORD, P. FEDERBUSH: *Pluriharmonic boundary values* *Tohoku Math.* (1974) 26 505–511.
- [4] V. BELOSAPKA: *Functions pluriharmonic on manifolds* *Math. U.S.S.R. Izvestija* (1978) 12 439–447.
- [5] P. de BARTOLOMEIS, G. TOMASSINI: *Traces of pluriharmonic functions* *Analytic Functions Kozubnik 1979 Lecture Notes in Math.* 798 Springer (1980) 10–17.
- [6] J.J. KOHN, U. ROSSI: *On the extension of holomorphic functions from the boundary of a complex manifold* *Ann. of Math.* (1965) 81 451–473.
- [7] G. LAVILLE: *Fonctions pluriharmoniques et solution fondamentale d'un opérateur du 4<sup>e</sup> ordre* *Bull. Sc. Math.* 2s. (1977) 101 305–317.
- [8] S. LOJASIEWICZ, G. TOMASSINI: *Valeurs au bord des formes holomorphes* *Several Complex Variables, Proceedings of International Conferences, Cortona Italy 1976–1977* 222–245 *Scuola Normale Superiore, Pisa* (1978).
- [9] J. POLKING, R. O. WELLS: *Boundary values of Dolbeault Cohomology classes and a generalized Bochner–Hartogs Theorem* *Abh. Math. Sem. Univ. Hamburg* (1978) 47 3–24.
- [10] G.B. RIZZA: *Dirichlet problem for n-harmonic functions and related geometrical properties* *Math. Ann. Bd.* (1955) 130 s.202–218.

(Oblatum 6-III-1981 & 5-VI-1981)

Giuseppe Tomassini  
 Università delqe Studi  
 Istituto Matematico Ulisse  
 Dine  
 Viale Morgagni 67-A  
 Firenze 50134  
 Italy