

COMPOSITIO MATHEMATICA

J. DODZIUK

MIN-OO

**An L_2 -isolation theorem for Yang-Mills fields
over complete manifolds**

Compositio Mathematica, tome 47, n° 2 (1982), p. 165-169

http://www.numdam.org/item?id=CM_1982__47_2_165_0

© Foundation Compositio Mathematica, 1982, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

AN L_2 -ISOLATION THEOREM FOR YANG-MILLS FIELDS OVER COMPLETE MANIFOLDS

J. Dodziuk and Min-Oo

1. Introduction

In this note we extend the results of the preceding paper [4] to the case of non compact complete manifolds. Beyond the method of [4] we only make use of appropriate cut-off functions as in [2]. This cut-off trick is due to Andreotti and Vesentini (see [5, Th. 26]). It is the point of view of [2] that every vanishing theorem based on a Weitzenböck identity generalizes from the compact to the complete case for L_2 -forms. On the other hand the results in [4] are proved by applying a Sobolev inequality to a Weitzenböck formula for certain bundle valued harmonic forms. Thus it is not surprising that the L_2 -isolation theorem of the preceding paper extends to complete manifolds.

We shall use freely the notation and formulae of [4]. However the isoperimetric constant c_1 will have to be replaced by another isoperimetric constant $c_0 = c_0(M)$ defined as follows:

$$c_0 = \inf_D \frac{(\text{vol}(\partial D))^4}{(\text{vol}(D))^3},$$

where D ranges over all open, relatively compact subsets of M with smooth boundary. M is assumed from now on to be a noncompact, complete, oriented, 4-dimensional, Riemannian manifold.

We begin by stating the results. First, our method yields a simple proof of the following result of C.-L. Shen [6].

This work was supported in part by the Sonderforschungsbereich "Theoretische Mathematik" (SFB40) at the University of Bonn (both authors), and Sloan Fellowship and NSF Grant No. MCS8024276 (first author).

THEOREM 1: *Assume that*

$$(i) \quad k_- - \frac{1}{\sqrt{3}}|\Omega_-| \geq 0$$

with strict inequality holding at some point of M .

Suppose further that β is a harmonic section of the bundle $\Lambda^2 \otimes E$ satisfying the decay condition:

$$(ii) \quad \lim_{R \rightarrow \infty} \frac{1}{R^2} \int_{B_{2R}(x_0) \setminus B_R(x_0)} |\beta|^2 = 0$$

for some point $x_0 \in M$. Then $\beta \equiv 0$.

In particular, if ω is a sourceless Yang-Mills field such that Ω_- satisfies (i) and (ii), then $\Omega_- \equiv 0$.

As a generalization of Theorem 2 of [4] we obtain the following

THEOREM 2: *Assume that the curvature of M satisfies $k_- \geq 0$. If $\|\Omega_-\|_2^2 < c_0/108$, then every square integrable harmonic section of $\Lambda^2 \otimes E$ vanishes identically. In particular, if ω is a sourceless Yang-Mills field with $\|\Omega_-\|_2^2 < c_0/108$ then $\Omega_- \equiv 0$.*

Of course, this theorem is of interest only if $c_0(M) > 0$. This is the case for \mathbb{R}^4 with the flat metric and we obtain the following:

COROLLARY: *Let ω be a sourceless Yang-Mills field over \mathbb{R}^4 equipped with a complete conformally flat metric. If $\int_{\mathbb{R}^4} |\Omega_-|^2 < \pi^2 2^5/3^3$, then $\Omega_- \equiv 0$.*

This corollary yields an improvement of the constant in Theorem 3 of [4].

THEOREM 3: *Let ω be a sourceless Yang-Mills field over S^4 with a conformally flat metric. If*

$$\frac{1}{2} \int_{S^4} |\Omega|^2 < 2\pi^2 \left(|p_1(E)| + \frac{16}{27} \right)$$

then ω is either self-dual or anti-self-dual.

Finally the following result gives a lower bound of the spectrum of the Laplacian Δ_-^ω .

THEOREM 4: *Suppose $2k_- \geq \mu > 0$. If*

$$\|\Omega_-\|_2^2 \leq \frac{c_0}{108}, \text{ then } \text{Spec}(\Delta^\omega) \subset [\mu, \infty).$$

We now prove the theorems stated above. For a given $x_0 \in M$ we can construct (cf. [5]) a family $\{\lambda_R\}_{R>0}$ of Lipschitz continuous function $\lambda_R : M \rightarrow \mathbb{R}$ with the following properties

- (i) $\text{supp } \lambda_R \subset B_{2R}(x_0)$
- (ii) $0 \leq \lambda_R \leq 1$
- (2) (iii) $\lambda_R|_{B_R(x_0)} \equiv 1$
- (iv) $\lim_{R \rightarrow \infty} \lambda_R = 1$
- (v) $|d\lambda_R| < \frac{C}{R}$ a.e.,

where $d\lambda_R$ exists almost everywhere since λ_R is Lipschitz and the constant C is independent of R . In what follows we shall write λ for λ_R . Set $\beta_- = \beta$ in the Weitzenböck identity (3.3) of [4] and take the inner product with $\lambda^2\beta$. Integration by parts, which is permitted since $\text{supp } \lambda^2\beta \subset B_{2R}(x_0)$ is compact, now yields

$$(3) \quad (\Delta^\omega\beta, \lambda^2\beta) = (\nabla\beta, \nabla(\lambda^2\beta)) + \left(\frac{\kappa}{6}\beta, \lambda^2\beta\right) - (\beta \circ W_-, \lambda^2\beta) - (\Omega_-, [\beta, \lambda^2\beta]).$$

Leibnitz rule shows that

$$(4) \quad (\nabla\beta, \nabla(\lambda^2\beta)) = \|\nabla(\lambda\beta)\|_2^2 - \|d\lambda \otimes \beta\|_2^2.$$

Hence if $\Delta^\omega\beta = 0$, estimating the last three terms on the right hand side of (3) as in [4], we obtain

$$(5) \quad \|d\lambda \otimes \beta\|_2^2 \geq \int_M \left(\frac{\kappa}{6} - \mu_-\right) |\lambda\beta|^2 - \frac{2}{\sqrt{3}} \int_M |\Omega_-| |\lambda\beta|^2 \geq 2 \int_M \left(k_- - \frac{1}{\sqrt{3}} |\Omega_-|\right) |\lambda\beta|^2.$$

Observe that for $\lambda = \lambda_R$, (2) implies

$$(6) \quad \|d\lambda \otimes \beta\|_2^2 \leq \frac{c^2}{R^2} \int_{B_{2R}(x_0)/B_R(x_0)} |\beta|^2.$$

Passing to the limit as $R \rightarrow \infty$ in (5) we see that under the assumptions of Theorem 1

$$\int_M \left(k_- - \frac{1}{\sqrt{3}} \|\Omega_-\| \right) |\beta|^2 = 0.$$

Hence $\beta = 0$ on an open set. By the unique continuation theorem of Aronszajn, Krzywicki and Szarski (cf. [5]) $\beta \equiv 0$ and Theorem 1 is proved.

To prove Theorem 2 we use the Sobolev inequality of P. Li [3, Lemma 6]

$$\|\nabla f\|_2^2 \geq \frac{1}{9} \sqrt{c_0} \|f\|_4^2$$

for compactly supported functions, which implies in our case that

$$(7) \quad \|\nabla(\lambda\beta)\|_2^2 \leq \frac{1}{9} \sqrt{c_0} \|\lambda\beta\|_4^2.$$

Now assuming $\Delta^\omega \beta = 0$, substituting (4) into (3), using (7) together with the definition of k_- and the pointwise estimate (3.8) of [4], we obtain

$$\begin{aligned} \|d\lambda \otimes \beta\|_2^2 &\geq \left(\frac{1}{9} \sqrt{c_0} - \frac{2}{\sqrt{3}} \|\Omega_-\|_2 \right) \|\lambda\beta\|_4^2 \\ &\quad + 2 \int k_- |\lambda\beta|^2. \end{aligned}$$

Theorem 2 now follows by passing to the limit as $R \rightarrow \infty$, since by (6) $\lim_{R \rightarrow \infty} \|d\lambda \otimes \beta\|_2^2 = 0$ if β is square integrable.

The corollary follows from Theorem 2 by substituting the value $c_0(\mathbb{R}^4) = 2^7 \pi^2$. Theorem 3 follows from the corollary since \mathbb{R}^4 with the flat metric is conformally equivalent to $S^4 \setminus \{\text{pt.}\}$ with the standard metric and because the Yang-Mills functional is conformally invariant.

We now turn to the proof of Theorem 4. The Laplacian Δ^ω is essentially self-adjoint on $C_0^\infty(\Lambda^2 \otimes E)$. This is a consequence of completeness (cf. [1]). Thus it suffices to estimate $(\Delta^\omega \beta, \beta)$ for compactly supported β . From the Weitzenböck identity (3.3) of [4], the Sobolev inequality (7), the definition of k_- and the estimate (3.8) of [4] we obtain through integration by parts the following estimate:

$$\begin{aligned} (\Delta^\omega \beta, \beta) &\geq \left(\frac{1}{9} \sqrt{c_0} - \frac{2}{\sqrt{3}} \|\Omega_-\|_2 \right) \|\beta\|_4^2 + 2 \int k_- |\beta|^2 \\ &\geq \mu \|\beta\|_2^2 \end{aligned}$$

provided the assumptions of Theorem 4 are satisfied. This proves the theorem.

REFERENCES

- [1] P.R. CHERNOFF: Essential self-adjointness of powers of generators of hyperbolic equations. *Journal of Functional Analysis* 12 (1973) 401–414.
- [2] J. DODZIUK: Vanishing theorems for square integrable harmonic forms, *Geometry and analysis*, papers dedicated to the memory of V.K. Patodi, Indian Academy of Sciences and Tata Institute of Fundamental Research, Bombay, 1981, pp. 21–27.
- [3] P. LI: On the Sobolev constant and the p -Spectrum of a compact Riemannian manifold, *Ann. Scient. Éc. Norm. Sup. 4^e serie*, t. 13 (1980) 451–469.
- [4] MIN-OO: An L_2 -isolation theorem for Yang-Mills fields, this journal.
- [5] G. DERHAM: *Variétés différentiables*, Hermann, Paris 1973.
- [6] C.-L. SHEN: *Gap-phenomena of Yang-Mills fields over complete manifolds*, preprint.

(Oblatum 7-X-1981)

J. Dodziuk
Department of Mathematics
Queens College, CUNY
Flushing, N.Y. 11367
U.S.A.

Min-Oo
Mathematisches Institut der
Universität Bonn
Wegelerstr. 10
D5300-Bonn 1
W. Germany

Added in proof

In “Best Constant in Sobolev Inequality”, *Ann. Mat. Pure Appl.* 110 (1976) 353–372, G. Talenti shows that the best constant in the Sobolev inequality $\|\nabla f\|^2 \geq c\|f\|_2^2$ for functions on R^4 is $c = (8\pi/\sqrt{6})$. Using this we can improve the Corollary of Theorem 2 and Theorem 3. In the corollary the constant $\pi^2 2^5/3^3$ can be replaced by $8\pi^2$ and in Theorem 2 $16/27$ may be replaced by 2. The statements obtained this way are *optimal*. In fact, Bourguignon and Lawson, in “Stability and Isolation Phenomena for Yang-Mills Fields”, *Commun. Math. Phys.* 79 (1981) 189–230, exhibit Yang-Mills fields on S^4 with its canonical metric for which the pointwise norm $|\Omega_-| \equiv \sqrt{3}$, and hence $\|\Omega_-\|_2^2 = 8\pi^2$. Analyzing the case of equality carefully, we can show that if $\|\Omega_-\|_2^2 = 8\pi^2$, then $|\Omega_-| \equiv \sqrt{3}$. Such fields have been classified by Bourguignon and Lawson.