

# COMPOSITIO MATHEMATICA

S. GLASNER

D. MAON

## **On absolutely extremal points**

*Compositio Mathematica*, tome 59, n° 1 (1986), p. 51-56

<[http://www.numdam.org/item?id=CM\\_1986\\_\\_59\\_1\\_51\\_0](http://www.numdam.org/item?id=CM_1986__59_1_51_0)>

© Foundation Compositio Mathematica, 1986, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## ON ABSOLUTELY EXTREMAL POINTS

S. Glasner and D. Maon

### Abstract

Given three doubly asymptotic points  $x, y, z$  in a minimal flow  $X$ , we construct an affine embedding  $\varphi: X \rightarrow Q$  such that  $\varphi(x) = \frac{1}{2}(\varphi(y) + \varphi(z))$ . Thus  $x$  is not absolutely extremal. We produce an example of a metric minimal flow  $X$  with the property that for every  $x \in X$  a triple  $x, y, z$  as above exists, thereby showing that no point of  $X$  is absolutely extremal.

### Introduction

We recall the definitions of affine embedding and absolute extremality for flows, introduced in [1]. If  $(T, X)$  is a flow ( $T$  is a self homeomorphism of the compact space  $X$ ) and  $(T, Q)$  an affine flow (i.e.,  $Q$  is a compact convex set and  $T$  an affine homeomorphism) then an equivariant continuous map  $\varphi: X \rightarrow Q$  is called an *affine embedding* if  $\overline{\text{co}} \varphi(X) = Q$ . A point  $x \in X$  is called *absolutely extremal* if for every affine embedding  $\varphi: X \rightarrow Q$   $\varphi(x)$  is in  $\partial Q$ , the set of extreme points of  $Q$ .

Suppose  $(T, X)$  is metric and minimal (i.e., every orbit is dense) then for every affine embedding  $\varphi: X \rightarrow Q$  the set  $\{x \in X: \varphi(x) \in \partial Q\}$  is a dense  $G_\delta$ . It was shown in [1] that if  $(T, X)$  is metric and minimal then every distal point of  $X$  is absolutely extremal. Again under our assumptions on  $(T, X)$  the set of distal points is either empty or a dense  $G_\delta$ . These facts led the first author to ask in [1] whether every minimal metric flow must have absolutely extremal points.

The easiest examples where non-absolutely extremal points exist are given by certain almost automorphic flows where the flow  $X$  is presented as a set of sequences in  $l^\infty(Z)$  and the identity map of  $X$  into  $Q = \overline{\text{co}}(X) \subset l^\infty(Z)$  gives a natural affine embedding [1]. Some doubly asymptotic points of  $X$  turns out to be non-extreme in  $Q$ . In this note, we show that in any minimal flow a point with two doubly asymptotic points is not absolutely extremal. ( $x, y$  are doubly asymptotic if  $\lim_{|n| \rightarrow \infty} d(T^n x, T^n y) = 0$ ). We construct a minimal metric flow, every point of which has a

continuum of doubly asymptotic points; thus providing an example of a metric minimal flow no point of which is absolutely extremal.

The principle of construction is due to Grillenberger (see e.g. [4]), who first showed how to define a minimal set with some desired property as an intersection of a family of subshifts of finite type. A continuous version of Grillenberger's construction and applications of this method are described in [3] and [2]. The present paper can be considered as a sequel to [1] and we refer the reader to [1] for further motivation.

### Section 1. An affine embedding associated with doubly asymptotic points

**1.1 PROPOSITION:** *Let  $(X, T)$  be an infinite metric minimal flow,  $x_0, y_0, z_0 \in X$ , doubly asymptotic points. Then there exists an affine embedding  $\varphi: X \rightarrow Q$  such that  $\varphi(x_0) = \frac{1}{2}(\varphi(y_0) + \varphi(z_0))$ .*

**PROOF:** In  $C^*(X)$  we let  $V$  be the weak \* closed linear space spanned by the set

$$V_0 = \left\{ \delta_{T^n x_0} - \frac{1}{2}(\delta_{T^n y_0} + \delta_{T^n z_0}) : n \in \mathbb{Z} \right\}.$$

We let  $\pi: C^*(X) \rightarrow E = C^*(X)/V$  be the quotient map and define  $\varphi: X \rightarrow E$  by  $\varphi(x) = \pi(\delta_x) = \delta_x + V$ . Put  $Q = \overline{\text{co}}(\varphi(X))$  and let

$$W = \left\{ \eta \in C^*(X) : \eta = \sum_{n \in \mathbb{Z}} a_n \left( \delta_{T^n x_0} - \frac{1}{2}(\delta_{T^n y_0} + \delta_{T^n z_0}) \right), \right. \\ \left. \times \sum_{n \in \mathbb{Z}} |a_n| < \infty \right\}.$$

We claim that  $V = W$ ; to see this let  $\sum_{n \in \mathbb{Z}} |a_n| < \infty$  be given. Put

$$\eta_N = \sum_{|n| \leq N} a_n \left( \delta_{T^n x_0} - \frac{1}{2}(\delta_{T^n y_0} + \delta_{T^n z_0}) \right)$$

and let  $\eta$  be the infinite sum. For every  $f \in C(X)$  we have

$$|f(\eta_N) - f(\eta)| \leq \|f\| 2 \sum_{|n| > N} |a_n| \rightarrow 0.$$

Hence  $\eta_N \rightarrow \eta$  and  $\eta \in V$ . Thus  $W \subset V$ ; since  $V_0 \subset W$  it is enough to show that  $W$  is weak \* closed. By Krein-Šmulyan's theorem it suffices to

show that  $W \cap B_r$  is weak \* closed, where  $B_r = \{\nu \in C^*(X) : \|\nu\| \leq r\}$ . Since  $B_r$  is metrizable we can deal with sequences. So let

$$\eta^k = \sum_{n \in \mathbb{Z}} a_n^k \left( \delta_{T^{n}x_0} - \frac{1}{2}(\delta_{T^{n}y_0} + \delta_{T^{n}z_0}) \right)$$

be a sequence in  $W \cap B_r$  with  $\eta^k \rightarrow \eta \in V$ . We have

$$\text{Sup}_k \|\eta^k\| = \text{Sup}_k \text{Sup} \{ |f(\eta^k)| : \|f\| = 1 \} = \text{Sup}_k 2 \sum_{n \in \mathbb{Z}} |a_n^k| < r.$$

Using a diagonal process we can choose a subsequence  $\eta^{k_i}$  such that for each  $n$   $a_n^{k_i} \rightarrow b_n$ . For convenience we denote this subsequence also by  $\eta^k$ , thus we now assume  $a_n^k \rightarrow b_n$  for every  $n$ . Using Fatou's lemma (in  $l_1(\mathbb{Z})$ ) we have

$$\sum_{n \in \mathbb{Z}} |b_n| \leq \underline{\lim} \sum_{n \in \mathbb{Z}} |a_n^k| < r.$$

Put

$$\tilde{\eta} = \sum_{n \in \mathbb{Z}} b_n \left( \delta_{T^{n}x_0} - \frac{1}{2}(\delta_{T^{n}y_0} + \delta_{T^{n}z_0}) \right)$$

and let  $f \in C(X)$  and  $\epsilon > 0$  be given. Choose  $\delta > 0$  such that  $d(x, y) < \delta \Rightarrow |f(x) - f(y)| < \epsilon$  and  $N$  with

$$|n| > N \Rightarrow d(T^n y_0, T^n z_0), d(T^n z_0, T^n x_0), d(T^n y_0, T^n x_0) < \delta.$$

Then

$$\begin{aligned} |f(\eta^k - \tilde{\eta})| &= \left| \sum_{|n| \leq N} (a_n^k - b_n) [f(T^n x_0) - \frac{1}{2}(f(T^n y_0) + f(T^n z_0))] \right. \\ &\quad \left. + \sum_{|n| > N} (a_n^k - b_n) [f(T^n x_0) - \frac{1}{2}(f(T^n y_0) + f(T^n z_0))] \right| \\ &\leq 2\|f\| \sum_{|n| \leq N} |a_n^k - b_n| + \epsilon \sum_{|n| > N} (|a_n^k| + |b_n|) \\ &\leq 2\|f\| \sum_{|n| \leq N} |a_n^k - b_n| + \epsilon(\|\eta^k\| + \|\tilde{\eta}\|). \end{aligned}$$

It follows that  $\eta^k \rightarrow \tilde{\eta}$  so that  $\tilde{\eta} = \eta$  is in  $W$  and  $V = W$ .

Clearly  $\varphi$  is continuous and equivariant from  $X$  into the affine flow  $\mathcal{Q}$ . If  $\varphi(x) = \varphi(y)$  then  $\delta_x - \delta_y \in V$ . But as  $V = W$ , every non-zero

measure in  $V$  is supported by at least three points. Thus  $x = y$  and  $\varphi$  is one to one. Finally

$$\begin{aligned}\varphi(x_0) &= \delta_{x_0} + V = \delta_{x_0} - \left( \delta_{x_0} - \frac{1}{2}(\delta_{y_0} + \delta_{z_0}) \right) + V \\ &= \frac{1}{2}(\delta_{y_0} + \delta_{z_0}) + V = \frac{1}{2}(\varphi(y_0) + \varphi(z_0)).\end{aligned}$$

This completes the proof.  $\square$

## Section 2. A metric minimal flow every point of which has a continuum of asymptotic points

Let  $\Omega = [0, 1]^Z$  denote the compact metric space of two sided  $[0, 1]$  valued sequences with the metric  $d(x, y) = \sup_{n \in Z} 2^{-|n|} |x_n - y_n|$ . For a closed  $W \subset [0, 1]^n$  and  $i \in Z$  we let

$$C_i(W) = \{x \in \Omega : \forall j \in Z, x[i + jn, i + (j + 1)n - 1] \in W\}$$

and  $C(W) = \bigcup_{i=1}^n C_i(W)$ . We define inductively a sequence  $n_k$  and closed sets  $W_k \subset W_{k-1}^{n_k}$  as follows. Let  $W_0 = [0, 1]$ . Given  $W_{k-1}$  we choose an arbitrary but fixed  $2^{-k}$ -net  $\{u_1, u_2, \dots, u_{l_k}\}$  of  $W_{k-1}^2$ , where the metric on a finite dimensional cube  $[0, 1]^n$  is  $d(w, v) = \sup_{1 \leq i \leq n} |w_i - v_i|$ . Let  $n_k = 100l_k$  and define

$$\begin{aligned}W_k &= \{w \in W_{k-1}^{n_k} : \text{there exist odd indices } 1 \leq i_1, i_2, \dots, i_{l_k} \leq n_k \\ &\quad \text{such that } w_{i_j} w_{i_j+1} = u_j \text{ for } j = 1, 2, \dots, l_k \\ &\quad \text{where } w = w_1 w_2 \dots w_{n_k}, w_{i_j} \in W_{k-1}\}.\end{aligned}$$

We call the set  $\{i_1, i_2, \dots, i_{l_k}\}$  a  $u$ -set for  $w$ . Put  $X = \bigcap_{k=1}^{\infty} C(W_k)$ .

**2.1 PROPOSITION:** *Let  $T$  be the shift on  $X$ , then  $(X, T)$  is a minimal flow.*

**PROOF:** Follows directly from the way  $X$  was defined.

**2.2 PROPOSITION:** *For every  $k$ ,  $W_k$  is pathwise connected.*

**PROOF:** Assume  $W_{k-1}$  is connected. Let  $w, w' \in W_k$ ,  $w = w_1 w_2 \dots w_{n_k}$ ,  $w' = w'_1 w'_2 \dots w'_{n_k}$ ,  $w_i, w'_i \in W_{k-1}$ . Assume first that there exists a  $u$ -set  $A = \{i_1, i_2, \dots, i_{l_k}\}$  common to  $w$  and  $w'$ . We let  $w(t) = w_1(t) w_2(t) \dots w_{n_k}(t)$  be defined by  $w_{i_j}(t) w_{i_j+1}(t) \equiv u_j$  if  $i_j \in A$  and where for all other  $i$ 's  $w_i(t)$  is a path in  $W_{k-1}$  connecting  $w_i$  and  $w'_i$ . Clearly  $w(t) \in W_k$  for every  $t \in [0, 1]$ . For the general case let  $A = \{i_1, i_2, \dots, i_{l_k}\}$

and  $A' = \{i'_1, i'_2, \dots, i'_k\}$  be  $u$ -sets for  $w$  and  $w'$  respectively. Choose  $A'' = \{i''_1, i''_2, \dots, i''_k\} \subset \{1, 2, \dots, n_k\}$  a set of odd indices disjoint from  $A \cup A'$  and define  $v = v_1 v_2 \dots v_{n_k}$ ,  $v' = v'_1 v'_2 \dots v'_{n_k}$  as follows

$$v_i v_{i+1} = \begin{cases} u_j & \text{if } i = i_j \text{ or } i = i'_j \\ w_i w_{i+1} & \text{otherwise} \end{cases}$$

$$v'_i v'_{i+1} = \begin{cases} u_j & \text{if } i = i'_j \text{ or } i = i''_j \\ w'_i w'_{i+1} & \text{otherwise.} \end{cases}$$

Clearly  $v, v' \in W_k$ . Now as  $A$  is a common  $u$ -set for  $w$  and  $v$ ,  $A'$  a common  $u$ -set for  $v$  and  $v'$  and  $A''$  a common  $u$ -set for  $v'$  and  $w''$ , we conclude by the first part of the proof, that there exists a path in  $W_k$  connecting  $w$  and  $w'$ .  $\square$

**DEFINITION:** Let  $K$  be a natural number  $1 \leq r, s \leq l_k$ ,  $r \neq s$ . A *chain* from  $u_r$  to  $u_s$  is a set  $\{j_0, j_1, \dots, j_l\}$  of indices such that  $j_0 = r$ ,  $j_l = s$  and  $d(u_{j_n}, u_{j_{n+1}}) < 2^{-k}$ ,  $0 \leq n < l$ .

For every  $r$  and  $s$  as above, the existence of a chain from  $u_r$  to  $u_s$  follows from the fact that  $W_k$  is pathwise connected.

**DEFINITION:** For  $x \in X$  there exists by definition a sequence of integers  $\{t_k\}$  such that  $t_k \leq 0 < t_k + m_k$  (where  $m_k$  is the length of sequences in  $W_k$ ) and such that for every  $k$   $x \in C_{t_k}(W_k)$ . It is easy to see that one can choose  $\{t_k\}$  so that  $\forall k$   $t_{k-1} \equiv t_k \pmod{m_{k-1}}$ . Such a sequence  $\{t_k\}$  will be called a *block partition* for  $x$ .

**2.3 PROPOSITION:** Let  $x \in X$ ,  $\{t_k\}$  a block partition for  $x$  and  $w_0 \in W_{k_0}$  for some  $k_0$ . Then there exists  $y \in X$  such that

- (1)  $y[t_{k_0}, t_{k_0} + m_{k_0} - 1] = w_0$
- (2)  $y$  is doubly asymptotic to  $x$ .

**PROOF:** We define  $y[t_k, t_k + m_k - 1]$  by induction on  $k$ . Put  $y[t_{k_0}, t_{k_0} + m_{k_0} - 1] = w_0$ . Let  $x[t_k, t_k + m_k - 1] = w_1 w_2 \dots w_{n_k} = w$ ,  $w_i \in W_{k-1}$ ,  $i = 1, \dots, n_k$ , and suppose  $x[t_{k-1}, t_{k-1} + m_{k-1} - 1]$  is  $w_n$ . Let  $A$  be a  $u$ -set for  $w$ . If  $n, n-1 \notin A$  define  $y[i] = x[i]$  for  $t_k \leq i \leq t_k + m_k - 1$ ,  $i \notin [t_{k-1}, t_{k-1} + m_{k-1} - 1]$  and then clearly  $y[t_k, t_k + m_k - 1] \in W_k$ .

If  $n = i_r \in A$ , let  $m$ ,  $1 \leq m \leq n_k - 1$  be an odd integer such that  $m \notin A$ . There exists an  $s$ ,  $1 \leq s \leq l_k$  such that  $d(u_s, w_m w_{m+1}) < 2^{-k}$ . Let  $j_0, j_1, \dots, j_l$  be a chain from  $u_r$  to  $u_s$ ,  $j_0 = r$ ,  $j_l = s$ ; thus for  $i_{j_t} \in A$   $w_{i_{j_t}} w_{i_{j_t}+1} = u_{j_t}$ ,  $t = 1, \dots, l$ . Put  $w' = w'_1 w'_2 \dots w'_{n_k}$  where for an odd  $i$

$$w'_i w'_{i+1} = \begin{cases} u_{j_{t-1}} & \text{if } i = i_{j_t}, 0 < t \leq l \\ u_s & \text{if } i = m \\ w_i w_{i+1} & \text{otherwise} \end{cases}$$

Since for every  $t$ ,  $d(u_{j_{t-1}}, u_{j_t}) < 2^{-k}$  and also  $d(u_s, w_m w_{m+1}) < 2^{-k}$  we have  $d(w, w') < 2^{-k}$ . Let  $y[t_k, t_k + m_k - 1] = w'_1 w'_2 \dots w'_{n-1} y[t_{k-1}, t_{k-1} + m_{k-1} - 1] w'_{n+1} \dots w'_{n_k}$ , then clearly  $y[t_k, t_k + m_k - 1] \in W_k$ . If  $n - 1 = i_r \in A$  the construction is similar.

There are now three possibilities:

- (1)  $t_k \rightarrow \infty$ ,  $t_k + m_k \rightarrow \infty$ , in which case  $y$  is now fully defined.
- (2) There exists  $k_0$  such that for  $k \geq k_0$ ,  $t_k = t_{k_0}$ . In this case define  $y[-\infty, t_{k_0} - 1] = x[-\infty, t_{k_0} - 1]$ .
- (3) There exists  $k_0$  such that for  $k \geq k_0$ ,  $t_k + m_k = t_{k_0} + m_{k_0}$ .

In this case define  $y[t_{k_0} + m_{k_0} + 1, \infty] = x[t_{k_0} + m_{k_0} + 1, \infty]$ . By definition of  $y$  we have for  $i < t_k$  and  $i > t_k + m_k$ ,  $|y[i] - x[i]| < 2^{-k}$ . Thus  $y$  and  $x$  are asymptotic.  $\square$

## References

- [1] S. GLASNER: Absolutely extremal points in minimal flows. *Compositio Math.* 55 (1985) 263–268.
- [2] S. GLASNER and D. MAON: An inverted tower of almost 1-1 extensions. *Journal d'Analyse Math.* Vol 44 (1984/85) 67–75.
- [3] D. MAON: On the relation  $N$ , the obstruction to uniform rigidity. Tel Aviv University.
- [4] M. DENKER, C. GRILLENBERGER and K. SIGMUND: Ergodic theory on compact spaces, *Lecture Notes in Mathematics* 527. Springer Verlag (1976).

(Oblatum 14-XI-1984 & 6-II-1985)

Department of Mathematics  
University of Washington  
Seattle, WA 98195  
USA

and

Department of Mathematics  
Tel Aviv University  
Tel Aviv  
Israel