

# COMPOSITIO MATHEMATICA

TAKASHI ICHIKAWA

## **The universal periods of curves and the Schottky problem**

*Compositio Mathematica*, tome 85, n° 1 (1993), p. 1-8

[http://www.numdam.org/item?id=CM\\_1993\\_\\_85\\_1\\_1\\_0](http://www.numdam.org/item?id=CM_1993__85_1_1_0)

© Foundation Compositio Mathematica, 1993, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## The universal periods of curves and the Schottky problem

TAKASHI ICHIKAWA

*Department of Mathematics, Faculty of Science and Engineering, Saga University, Saga 840, Japan*

Received 13 February 1991; accepted 8 October 1991

### Introduction

The aim of this paper is to give a new approach to the Schottky problem which means, in this paper, to characterize Siegel modular forms vanishing on the jacobian locus. Our characterization uses the formal power series expressing periods of algebraic curves and deduces new relations between the Fourier coefficients.

In [10] and [7], Schottky and Manin-Drinfeld obtained the infinite product expression of the multiplicative periods of Schottky uniformized Riemann surfaces and Mumford curves respectively. First, we show the infinite products can be expressed as certain formal power series, which we call the *universal power series for multiplicative periods of algebraic curves*, in terms of the *Koebe coordinates* which are the fixed points and the ratios of generators of the corresponding Schottky groups. The existence of the universal periods is suggested by the result of Gerritzen [4] which determined a fundamental domain in the Schottky space over a complete valuation field. Our result also gives an algorithm to compute the universal periods. By using our result, one can deduce Mumford's result on the periods of degenerate Riemann surfaces ([9], IIIb, §5) and the asymptotic behavior of the periods of Riemann surfaces ([3], Corollary 3.8).

Next, we study the Schottky problem by using the universal periods. Let  $f$  be a Siegel modular form of degree  $g$  over  $\mathbf{C}$  given by  $(Z = (z_{ij}) \in$  the Siegel upper half space of degree  $g$ )

$$f(Z) = \sum_T a(T) \exp(2\pi\sqrt{-1} \cdot \text{Tr}(TZ)).$$

Since  $\exp(2\pi\sqrt{-1} \cdot z_{ij})$  are the multiplicative periods of the Riemann surface with period matrix  $Z$ ,  $f|_{q_{ij}=p_{ij}}$  ( $q_{ij} := \exp(2\pi\sqrt{-1} \cdot z_{ij})$ ,  $p_{ij}$ : the universal periods) is the expansion of the form (on the moduli space of curves of genus  $g$ ) induced from  $f$  in terms of the Koebe coordinates. Then as an analogy of a part of the  $q$ -

expansion principle, we have

$$f \text{ vanishes on the jacobian locus} \Leftrightarrow f|_{q_{ij}=p_{ij}} = 0.$$

This follows from the retrosection theorem ([6]) which says that every Riemann surface can be Schottky uniformized. As an immediate consequence of this result, we have

**THEOREM** (cf. Corollary 3.3). *Let*

$$f(Z) = \sum_T a(T) \exp(2\pi\sqrt{-1} \cdot \text{Tr}(TZ))$$

*be any Siegel modular form of degree  $g$  over  $\mathbf{C}$  vanishing on the jacobian locus, and put  $S = \min\{\text{Tr}(T) \mid a(T) \neq 0\}$ . Then for any set  $\{s_1, \dots, s_g\}$  of non-negative  $g$ -integers satisfying  $\sum_{i=1}^g s_i = S$ ,*

$$\sum_{t_i=s_i} a(T) \prod_{i < j} \left( \frac{(x_i - x_j)(x_{-i} - x_{-j})}{(x_i - x_{-j})(x_{-i} - x_j)} \right)^{2t_{ij}} = 0 \quad (T = (t_{ij})),$$

*where  $x_{\pm 1}, \dots, x_{\pm g}$  are variables.*

It is shown by Faltings [2] that Siegel modular forms over any field can be defined and have Fourier expansions in that field. In this paper, we prove our results for Siegel modular forms over any field by using the theory of Mumford curves ([8]) and the irreducibility of the moduli space of curves of given genus ([1]).

## 1. Mumford curves

In this section, we review some results in [4], [7], and [8].

1.1. Let  $K$  be a complete discrete valuation field with valuation  $\|\cdot\|$ . Let  $PGL_2(K)$  act on  $\mathbf{P}^1(K)$  by the Möbius transformations, i.e.,

$$g(z) = \frac{az+b}{cz+d} \left( g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \bmod K^\times \in PGL_2(K), z \in \mathbf{P}^1(K) \right).$$

Let  $\Gamma$  be a Schottky group over  $K$ , i.e., a finitely generated discrete subgroup of  $PGL_2(K)$  consisting of hyperbolic elements. Then it is known that  $\Gamma$  is a free group (cf. [5]). Let  $g$  be the rank of  $\Gamma$ , and  $\{\gamma_i \mid i = 1, \dots, g\}$  be a set of free generators of  $\Gamma$ . Since  $\gamma_i$  is hyperbolic, this can be expressed as

$$\gamma_i = \begin{pmatrix} \alpha_i & \alpha_{-i} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \beta_i \end{pmatrix} \begin{pmatrix} \alpha_i & \alpha_{-i} \\ 1 & 1 \end{pmatrix}^{-1} \bmod K^\times,$$

where  $\alpha_i, \alpha_{-i} \in \mathbf{P}^1(K)$  and  $\beta_i \in K^\times$  with  $|\beta_i| < 1$ . Then  $\alpha_{\pm i}$  are the fixed points of  $\gamma_i$  and  $\beta_i$  is one of the ratios of the eigenvalues of  $\gamma_i$ . Let

$$[a, b; c, d] = \frac{a-c}{a-d} \cdot \frac{b-d}{b-c}$$

denote the cross-ratio of four points. In [4], Gerritzen showed that if  $(\alpha_i, \alpha_{-i}, \beta_i)_{1 \leq i \leq g} \in (\mathbf{P}^1(K) \times \mathbf{P}^1(K) \times K^\times)^g$  satisfies  $\alpha_j \neq \alpha_k$  ( $j \neq k$ ) and  $|\beta_i| < \min\{|[\alpha_j, \alpha_k, \alpha_i, \alpha_{-i}]|\}$  for any  $i, j, k \in \{\pm 1, \dots, \pm g\}$  with  $j, k \neq \pm i$ , then the subgroup of  $PGL_2(K)$  generated by  $\gamma_i$  is a Schottky group with free generators  $\gamma_i$ . For each Schottky group  $\Gamma$  of rank  $g$  over  $K$ , let  $D_\Gamma$  be the set of points which are not limits of fixed points of elements of  $\Gamma - \{1\}$  in  $\mathbf{P}^1(K)$ . Then in [8], Mumford proved that the quotient  $D_\Gamma/\Gamma$  is the  $K$ -analytic space associated with a unique proper and smooth curve of genus  $g$  over  $K$  with multiplicative reduction, and that any proper and smooth curve of genus  $g$  over  $K$  with multiplicative reduction can be obtained in this way. Then  $D_\Gamma/\Gamma$  is called the Mumford curve associated with  $\Gamma$ , and we denote it by  $C_\Gamma$ .

1.2. Let  $x_i, x_{-i}$ , and  $y_i$  ( $i = 1, \dots, g$ ) be variables, and put  $\Omega = \mathbf{Q}(x_{\pm i}, y_i)$ . Let  $f_i$  be the element of  $PGL_2(\Omega)$  given by

$$(1.2.1) \quad f_i = \begin{pmatrix} x_i & x_{-i} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & y_i \end{pmatrix} \begin{pmatrix} x_i & x_{-i} \\ 1 & 1 \end{pmatrix}^{-1} \pmod{\Omega^\times},$$

and  $F$  the subgroup of  $PGL_2(\Omega)$  generated by  $f_i$  ( $i = 1, \dots, g$ ). Then  $F$  is a free group with generators  $f_i$ . For  $1 \leq i, j \leq g$ , let  $\psi_{ij}: F \rightarrow \Omega^\times$  be the map given by

$$\psi_{ij}(f) = \begin{cases} y_i & (\text{if } i = j \text{ and } f \in \langle f_i \rangle) \\ [x_i, x_{-i}; f(x_j), f(x_{-j})] & (\text{otherwise}). \end{cases}$$

Then it is easy to see that  $\psi_{ij}$  depends only on double coset classes  $\langle f_i \rangle f \langle f_j \rangle$  ( $f \in F$ ). Then in [7], Manin and Drinfeld showed that for any Schottky group  $\Gamma = \langle \gamma_i \rangle$  over  $K$ ,

$$q_{ij} = \prod_f \psi_{ij}(f)|_{x_{\pm i} = \alpha_{\pm i}, y_i = \beta_i}$$

( $f$  runs through all representatives of  $\langle f_i \rangle \backslash F / \langle f_j \rangle$ ) converges in  $K^\times$  and  $q_{ij}$  ( $1 \leq i, j \leq g$ ) are the multiplicative periods of  $C_\Gamma$ , i.e., the  $K$ -analytic space associated with the jacobian variety of  $C_\Gamma$  is given by the quotient

$$(K^\times)^g / \left( \left( \prod_j q_{ij}^{n_j} \right)_{1 \leq i \leq g} \mid n_j \in \mathbf{Z} \right).$$

## 2. Periods as power series

2.1. Let the notation be as above. For each integer  $k = -1, \dots, -g$ , put  $y_k = y_{-k}$  and  $f_k = (f_{-k})^{-1}$ . Then  $f_k$  satisfies (1.2.1) also. If the reduced expression of  $f \in F$  is  $\prod_{p=1}^n f_{\sigma(p)}$  ( $\sigma(p) \in \{\pm 1, \dots, \pm g\}$ ), then we put  $n(f) = n$ . Put

$$R = \mathbf{Z} \left[ x_i, \prod_{j < k} 1/(x_j - x_k) \right] \quad (i, j, k \in \{\pm 1, \dots, \pm g\}).$$

Let  $A$  be the ring of formal power series over  $R$  with variables  $y_1, \dots, y_g$ , i.e.,

$$A = R[[y_1, \dots, y_g]],$$

and  $I$  the ideal of  $A$  generated by  $y_i$  ( $i = 1, \dots, g$ ).

2.2. LEMMA. *For any  $f \in F$  and  $i \in \{\pm 1, \dots, \pm g\}$ ,  $f(x_i) \in A$ . Moreover, if the reduced expression of  $f \in F - \langle f_i \rangle$  is  $f_k \cdot f'$ , then  $f(x_i) \in x_k + I$ .*

*Proof.* We prove this by the induction on  $n(f)$ . Let  $f$  be an element of  $F - \langle f_i \rangle$  with reduced expression  $f_k \cdot f'$  such that  $f(x_i) = x_k + a$  for some  $a \in I$ . Then for any  $j \neq -k$ ,

$$(f(x_i) - x_{-j})^{-1} = \{(x_k - x_{-j})(1 + a/(x_k - x_{-j}))\}^{-1} \in A.$$

Hence

$$(f_j \cdot f)(x_i) = \left\{ x_j - \frac{(f(x_i) - x_j)x_{-j}y_j}{f(x_i) - x_{-j}} \right\} \cdot \left\{ 1 - \frac{(f(x_i) - x_j)y_j}{f(x_i) - x_{-j}} \right\}^{-1}$$

belongs to  $x_j + I$ . Assume that  $f \in \langle f_i \rangle$ . Then  $f(x_i) = x_i$ , and hence for any  $j \neq \pm i$ ,  $(f_j \cdot f)(x_i) \in x_j + I$ .

2.3. LEMMA. *For any  $f \in F$  having the reduced expression  $f' \cdot f_l$  with  $l \neq \pm j$ ,  $f(x_j) - f(x_{-j}) \in I^{n(f)}$ .*

*Proof.* We prove this by the induction on  $n(f)$ . If  $n(f) = 1$ , then  $f = f_l$  for some  $l \neq \pm j$ , and hence

$$f_l(x_j) - f_l(x_{-j}) = \frac{(x_l - x_{-l})^2(x_j - x_{-j})y_l}{\{x_j - x_{-l} - y_l(x_j - x_l)\}\{x_{-j} - x_{-l} - y_l(x_{-j} - x_l)\}}$$

belongs to  $I$ . Assume that the reduced expression of  $f \in F$  is  $f_k \cdot f' \cdot f_l$  ( $l \neq \pm j$ ) and  $f(x_j) - f(x_{-j}) \in I^{n(f)}$ . Then by Lemma 2.2, there exists  $b \in I$  such that

$f(x_j) = x_k + b$ . Therefore, for any  $m \neq -k$ ,

$$\{f(x_j) - x_{-m} - y_m(f(x_j) - x_m)\}^{-1} = \{(x_k - x_{-m}) + (b - y_m(f(x_j) - x_m))\}^{-1}$$

belongs to  $A$ . Similarly,  $\{f(x_{-j}) - x_{-m} - y_m(f(x_{-j}) - x_m)\}^{-1}$  belongs to  $A$ . Hence

$$\begin{aligned} & (f_m \cdot f)(x_j) - (f_m \cdot f)(x_{-j}) \\ &= \frac{(x_m - x_{-m})^2 (f(x_j) - f(x_{-j})) y_m}{\{f(x_j) - x_{-m} - y_m(f(x_j) - x_m)\} \{f(x_{-j}) - x_{-m} - y_m(f(x_{-j}) - x_m)\}} \end{aligned}$$

belongs to  $I^{n(f)+1}$ .

**2.4. PROPOSITION.** For any  $f \in F$  having the reduced expression  $f_k \cdot f' \cdot f_l$  with  $k \neq \pm i$  and  $l \neq \pm j$ ,

$$[x_i, x_{-i}; f(x_j), f(x_{-j})] \in 1 + I^{n(f)}.$$

*Proof.* By Lemma 2.2,  $f(x_j) - x_k$  and  $f(x_{-j}) - x_k$  belongs to  $I$ . Hence  $(f(x_j) - x_{-i})^{-1}$  and  $(f(x_{-j}) - x_i)^{-1}$  belongs to  $A$ . Therefore, by Lemma 2.3,

$$[x_i, x_{-i}; f(x_j), f(x_{-j})] = 1 + \frac{(x_i - x_{-i})(f(x_j) - f(x_{-j}))}{(f(x_j) - x_{-i})(f(x_{-j}) - x_i)}$$

belongs to  $1 + I^{n(f)}$ .

**2.5. COROLLARY.** For any  $1 \leq i, j \leq g$ , the infinite product  $\prod_f \psi_{ij}(f)$  ( $f$  runs through all representatives of  $\langle f_i \rangle \setminus F / \langle f_j \rangle$ ) is convergent in  $A$ .

2.6. We call  $\prod_f \psi_{ij}(f) \in A$  the universal power series for multiplicative periods of algebraic curves and denote them by  $p_{ij}$ . As seen in 1.2, when  $f_i \in PGL_2(\Omega)$  ( $i = 1, \dots, g$ ) are specialized to generators of any Schottky group  $\Gamma$  over a complete discrete valuation field,  $p_{ij}$  are specialized to the multiplicative periods  $q_{ij}$  of  $C_\Gamma$ .

2.7. By the above formulas, one can compute  $p_{ij}$  explicitly. For example,

$$\begin{aligned} p_{ij} &\equiv \left( 1 + \sum_{|k| \neq ij} \frac{(x_i - x_{-i})(x_j - x_{-j})(x_k - x_{-k})^2}{(x_i - x_k)(x_{-i} - x_k)(x_j - x_{-k})(x_{-j} - x_{-k})} y_{|k|} \right) \\ &\text{mod} \begin{cases} I^2 & (\text{if } i \neq j) \\ I^3 & (\text{if } i = j), \end{cases} \end{aligned}$$

where

$$c_{ij} = \begin{cases} [x_i, x_{-i}; x_j, x_{-j}] & (\text{if } i \neq j) \\ y_i & (\text{if } i = j). \end{cases}$$

### 3. The Schottky problem

3.1. Let  $k$  be a field and let  $f$  be a Siegel modular form of degree  $g \geq 2$  and weight  $h > 0$  defined over  $k$ , i.e., an element of

$$\Gamma(\mathcal{X}_g \otimes_{\mathbf{Z}} k, (\Lambda^g \pi_* (\Omega_{\mathcal{A}/\mathcal{X}_g}^1)^{\otimes h}).$$

Here  $\mathcal{X}_g$  denotes the moduli stack of principally polarized abelian varieties of dimension  $g$ , and  $\pi: \mathcal{A} \rightarrow \mathcal{X}_g$  denotes the universal abelian scheme. By a result of Faltings ([2], §6),  $f$  has the Fourier expansion rational over  $k$ :

$$F(f) = \sum_{T=(t_{ij})} a(T) \prod_{i,j=1}^g q_{ij}^{t_{ij}} \quad (a(T) \in k),$$

where  $T = (t_{ij})$  runs through all semi-integral (i.e.,  $2t_{ij} \in \mathbf{Z}$  and  $t_{ii} \in \mathbf{Z}$ ) and positive semi-definite symmetric matrices of degree  $g$  (if  $k = \mathbf{C}$ , then  $q_{ij} = \exp(2\pi\sqrt{-1} \cdot z_{ij})$ , where  $Z = (z_{ij})$  is in the Siegel upper half space of degree  $g$ ). Then by the symmetry  $q_{ij} = q_{ji}$  and the positive semi-definiteness of  $T$ ,  $F(f)$  belongs to

$$k \left[ q_{ij}, \prod_{i < j} 1/q_{ij} \right] [[q_{11}, \dots, q_{gg}]] \quad (1 \leq i, j \leq g).$$

Let  $\mathcal{M}_g$  denote the moduli stack of proper and smooth curves of genus  $g$ , and let  $\tau: \mathcal{M}_g \rightarrow \mathcal{X}_g$  denote the Torelli map.

3.2. THEOREM. *Let  $A$  be as in 2.1, and let*

$$\rho: k \left[ q_{ij}, \prod_{i < j} 1/q_{ij} \right] [[q_{11}, \dots, q_{gg}]] \rightarrow A \widehat{\otimes}_{\mathbf{Z}} k$$

*be the ring homomorphism over  $k$  defined by  $\rho(q_{ij}) = p_{ij}$  ( $1 \leq i, j \leq g$ ). Then for any Siegel modular form  $f$  of degree  $g$  defined over  $k$ , the pull back  $\tau^*(f)$  of  $f$  to  $\mathcal{M}_g \otimes_{\mathbf{Z}} k$  is equal to 0 if and only if  $\rho(F(f)) = 0$  in  $A \widehat{\otimes}_{\mathbf{Z}} k$ .*

*Proof.* We may assume that  $k$  is algebraically closed. First we assume that  $\tau^*(f) = 0$ . We regard  $k((z))$  as the complete discrete valuation field over  $k$  with prime element  $z$ . Then by the result of Gerritzen quoted in 1.1,  $\rho(F(f))$  vanishes

for any  $(x_i, x_{-i}, y_i)_{1 \leq i \leq g} \in (k \times k \times k[[z]])^g$  with  $x_j \neq x_k$  ( $j \neq k$ ) and  $y_i \in z \cdot k[[z]]$ . Since  $k$  is an infinite field,  $\rho(F(f)) = 0$ . Next we assume that  $\rho(F(f)) = 0$ . Let  $Z$  be the closed subset of  $\mathcal{M}_g \otimes_{\mathbb{Z}} k$  defined by  $\tau^*(f) = 0$ . Then  $Z$  can be identified with a closed subset of  $M_g \otimes_{\mathbb{Z}} k$ , where  $M_g$  denotes the coarse moduli scheme of proper and smooth curves of genus  $g$ . Since  $M_g \otimes_{\mathbb{Z}} k$  is irreducible ([1], 5.15), there exist a complete discrete valuation field  $K$  containing  $k$ , and a proper and smooth curve over  $K$  with multiplicative reduction which corresponds to the generic point  $\eta$  of  $M_g \otimes_{\mathbb{Z}} k$ . Hence by the assumption,  $Z$  contains  $\eta$ . Therefore, by the irreducibility of  $M_g \otimes_{\mathbb{Z}} k$ , we have  $Z = M_g \otimes_{\mathbb{Z}} k$ . This completes the proof.

3.3. COROLLARY. *Let  $f$  be a Siegel modular form of degree  $g$  defined over  $k$  whose Fourier expansion is  $\sum_{T=(t_i)} a(T) \prod_{i,j} q_i^{t_{ij}}$ , and put*

$$S = \min\{\text{Tr}(T) \mid a(T) \neq 0\}.$$

*If  $\tau^*(f) = 0$ , then for any set  $\{s_1, \dots, s_g\}$  of non-negative  $g$ -integers satisfying  $\sum_{i=1}^g s_i = S$ ,*

$$\sum_{t_i = s_i} a(T) \prod_{i < j} [x_i, x_{-i}; x_j, x_{-j}]^{2t_{ij}} = 0.$$

*Proof.* This follows from 2.7 and 3.2.

### Acknowledgements

The author wishes to express his sincere thanks to Professor D.B. Zagier who was interested in this subject and suggested some improvements in an earlier version of the paper.

### References

- [1] P. Deligne and D. Mumford: The irreducibility of the space of curves of given genus, *Publ. Math. IHES.* 36 (1969), 75–109.
- [2] G. Faltings: Arithmetische Kompaktifizierung des Modulraums der abelschen Varietäten, Proc. Arbeitstagung Bonn 1984, *Lecture Notes in Math.* 1111 (1985), 321–383, Springer-Verlag.
- [3] J.D. Fay: Theta functions on Riemann surfaces, *Lecture Notes in Math.* 352 (1973), Springer-Verlag.
- [4] L. Gerritzen: Zur analytischen Beschreibung des Raumes der Schottky-Mumford-Kurven, *Math. Ann.* 255 (1981), 259–271.
- [5] Y. Ihara: On discrete subgroups of the  $2 \times 2$  projective linear group over  $p$ -adic fields, *J. Math. Soc. Japan* 18 (1966), 219–235.

- [6] P. Koebe: Über die Uniformisierung der algebraischen Kurven IV, *Math. Ann.* 75 (1914), 42–129.
- [7] Yu. I. Manin and V. Drinfeld: Periods of  $p$ -adic Schottky groups, *J. Reine Angew. Math.* 262/263 (1972), 239–247.
- [8] D. Mumford: An analytic construction of degenerating curves over complete local fields, *Comp. Math.* 24 (1972), 129–174.
- [9] D. Mumford: Tata lectures on theta II, *Progress in Math.* 43 (1984), Birkhäuser.
- [10] F. Schottky: Über eine spezielle Function, welche bei einer bestimmten linearen Transformation ihres Arguments unverändert bleibt, *J. Reine Angew. Math.* 101 (1887), 227–272.