

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

K. A. HARDIE

K. H. KAMPS

Homotopy over B and under A

Cahiers de topologie et géométrie différentielle catégoriques, tome 28, n° 3 (1987), p. 183-196

http://www.numdam.org/item?id=CTGDC_1987__28_3_183_0

© Andrée C. Ehresmann et les auteurs, 1987, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

HOMOTOPY OVER B AND UNDER A
 by K.A. HARDIE and K.H. KAMPS

RÉSUMÉ, Dans cet article, on décrit une certaine catégorie d'homotopie cohérente \mathbb{H}_B^A d'espaces au-dessus d'un espace B et sous un espace A . Le problème d'isomorphisme et le problème de classification sont résolus. On indique aussi les liens avec les classes de composition secondaires.

0. INTRODUCTION.

An object X of \mathbb{H}_B^A is a diagram

$$(0.1) \quad A \xrightarrow{\sigma} X \xrightarrow{\rho} B$$

where X is a space and A and B are fixed spaces. An arrow from X to X' in \mathbb{H}_B^A will be a certain equivalence class of homotopy commutative diagrams of the form

$$\begin{array}{ccc}
 A & \xlongequal{\quad} & A \\
 \sigma \downarrow & \nearrow \sigma_t & \downarrow \sigma' \\
 X & \xrightarrow{\quad} & X' \\
 \rho \downarrow & \nearrow \rho_t & \downarrow \rho' \\
 B & \xlongequal{\quad} & B
 \end{array}$$

where u is a continuous map and σ_t and ρ_t are homotopies. Our first result (Theorem 1.2) is to the effect that such an arrow is an isomorphism in \mathbb{H}_B^A whenever the map u is (an ordinary) homotopy equivalence. As an immediate corollary we obtain (Corollary 1.8) that allowing σ and ρ to vary up to homotopy does not change the isomorphism class of X .

We shall denote the set of morphisms in \mathbb{H}_B^A from X to X' by $\pi(A/X, X'/B)$. If $u: X \rightarrow X'$ is a continuous map, let $\pi_1^X(X'; u)$ denote the u -based track group. (See [3, 2].) Then $\pi_1^X(X'; u)$ depends up to isomorphism only on the homotopy class $\{u\}$ of u . In Section 2 we classify the elements of $\pi(A/X, X'/B)$ in the sense that we exhibit a bijection between $\pi(A/X, X'/B)$ and the union over certain classes $\{u\}$ of sets of double cosets in $\pi_1^X(X'; u)$. The result generalizes classifications obtained in [2] for $\pi(X, X'/B)$ (the case $A = \emptyset$) and for $\pi(A/X, X')$ (the case B is a singleton).

In view of Corollary 1.6 the set $\pi(A/X, X'/B)$ is determined by the spaces A, X, X', B and by the homotopy classes $\{\sigma\}, \{\sigma'\}, \{\rho\}, \{\rho'\}$. Thus it is an invariant that is defined whenever we are given two factorizations of a homotopy class. In Section 3 we show, in a special case, that it is related to, and in a sense measures, the set of possible nontrivial secondary homotopy compositions of a certain type.

1. THE EQUIVALENCE RELATION.

If $h_0: Y \rightarrow Z$ and $h_1: Y \rightarrow Z$ are homotopies from Y to Z with the property that $h_0 = k_0$ and $h_1 = k_1$ then h_0 and h_1 are *relatively homotopic*, denoted $h_0 \equiv k_1$, if there exists a homotopy of homotopies $H_{t,s}: Y \rightarrow Z$ such that

$$h_0 = H_{t,0}, \quad k_1 = H_{t,1}, \quad H_{0,s} = h_0 = k_0 \quad (s \in I), \quad \text{and} \quad H_{1,s} = h_1 = k_1 \quad (s \in I).$$

The *track* $\{h_0\}$ of a homotopy h_0 is its relative homotopy class. The track of the constant homotopy $h_0 \equiv h_0$ is denoted by $\{h_0\}$.


For each homotopy commutative diagram 0.2 there is a triple $(\{\rho\}, u, \{\sigma\})$, where $\{\sigma\}$ and $\{\rho\}$ are tracks. We obtain an equivalence relation in the set of such triples if, whenever $u': X \rightarrow X'$ is a homotopy with $u_0 = u$, we define

$$(1.1) \quad (\{\rho' + \rho'u\}, u', \{u'\sigma + \sigma\}) \sim (\{\rho\}, u, \{\sigma\}).$$

In 1.1 the $+$ refers to the usual track addition of homotopies. Denoting the set of equivalence classes by $\pi(A/X, X'/B)$ we obtain a category with composition induced by juxtaposition of diagrams. Formally we set

$$(\{\rho'u', u'\sigma\}) \circ (\{\rho\}, u, \{\sigma\}) = (\{\rho' + \rho'u\}, u'u, \{u'\sigma + \sigma\}).$$

To check that composition respects the equivalence relation may at first glance present a problem, but we may bear in mind that a diagram



represents equally well the tracks $(\psi_0\beta_\epsilon + \psi_\epsilon\beta_1)$ and $(\psi_\epsilon\beta_0 + \psi_1\beta_\epsilon)$, for we have

$$\psi_0\beta_\epsilon + \psi_\epsilon\beta_1 \equiv \psi_\epsilon\beta_0 + \psi_1\beta_\epsilon.$$

The required verifications are now easily done. The identity morphism $X \rightarrow X$ in \mathbb{H}_B^A is the equivalence class of the triple $(\{\rho\}, 1_X, \{\sigma\})$. The equivalence class of a triple $(\{\rho_\epsilon\}, u, \{\sigma_\epsilon\})$ will be denoted simply by $(\rho_\epsilon, u, \sigma_\epsilon)$.

THEOREM 1.2. *The arrow $(\rho_\epsilon, u, \sigma_\epsilon)$ is an isomorphism in \mathbb{H}_B^A if and only if the map $u: X \rightarrow X'$ is a homotopy equivalence.*

PROOF. If $(\rho_\epsilon, u, \sigma_\epsilon)$ is an isomorphism then it is obvious that u is a homotopy equivalence. Suppose conversely that u is a homotopy equivalence. Then by [6] there exists a homotopy inverse u' of u and homotopies

$$\beta_\epsilon: uu' \simeq 1_X, \quad \psi_\epsilon: u'u \simeq 1_{X'}$$

$$(1.3) \quad \beta_\epsilon u \equiv u \psi_\epsilon.$$


It is claimed that

$$(\rho'\beta_{1-\epsilon} + \rho_{1-\epsilon}u'u', u\psi_{1-\epsilon} + \psi_\epsilon\sigma)$$


is inverse to $(\rho_\epsilon, u, \sigma_\epsilon)$ in \mathbb{H}_B^A , for firstly

$$(\rho_\epsilon, u, \sigma_\epsilon) \circ (\rho'\beta_{1-\epsilon} + \rho_{1-\epsilon}u'u', u\psi_{1-\epsilon} + \psi_\epsilon\sigma)$$

is represented by the following composite diagram




whose associated triple is equal to that of




(using 1.3) which is equal to $(\{\rho'\}, 1_X, \{\sigma'\})$. Secondly the composite

$$\{\rho'\beta_{1-t} + \rho_{1-t}U', U', U\sigma'_{1-t} + \psi\sigma\} \circ \{\rho_t, U, \sigma_t\}$$

is represented by the composite diagram



whose associated triple is equal to



(using 1.3) which is equal to $(\{\rho\}, 1_X, \{\sigma\})$, completing the proof.

COROLLARY 1.4. *Each object X in H_B^A is isomorphic to an object X' such that σ' is a closed cofibration and ρ' a Hurewicz fibration.*

PROOF. Let

$$A \xrightarrow{\sigma} X \xrightarrow{\rho} B$$

be the given object. By the mapping cylinder construction we can factor σ as qj where j is a closed cofibration and q is a homotopy equivalence. By [5], Proposition 2 (see also [4], Remark (d)) we can factor ρq as pi with p a fibration and i a closed cofibration and a homotopy equivalence. Now apply Theorem 1.2 twice to the diagram

$$(1.5) \quad \begin{array}{ccccc} A & \xlongequal{\quad} & A & \xlongequal{\quad} & A \\ ij \downarrow & & \downarrow j & & \downarrow \sigma \\ \bullet & \xleftarrow{\quad i \quad} & \bullet & \xrightarrow{\quad q \quad} & X \\ P \downarrow & & \downarrow \rho q & & \downarrow \rho \\ B & \xlongequal{\quad} & B & \xlongequal{\quad} & B. \end{array}$$

Now suppose that $\alpha_i: A \rightarrow X$ and $\rho_i: X \rightarrow B$ are homotopies. Applying Theorem 1.2 to the diagram

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ \downarrow \alpha_i & \nearrow & \downarrow \\ X & \xlongequal{\quad} & X \\ \downarrow \rho_i & \nearrow & \downarrow \\ B & \xlongequal{\quad} & B \end{array}$$

we obtain the following corollary.

COROLLARY 1.6. *The isomorphism type of an object X of H_B^A depends only on the homotopy classes of α and ρ .*

Let $f: A \rightarrow B$ be a fixed map and consider the class of all objects X of H_B^A such that $\rho\sigma = f$. Let HR_B^A be the category with these objects whose arrows are equivalence classes under the relation 1.1 of triples $(\{\rho_i\}, u, \{\sigma_i\})$ for which the tracks $\{\rho_i\}$ and $\{\sigma_i\}$ satisfy the additional condition

$$(1.7) \quad \{\rho_i\sigma_i\} + \{\rho_i\sigma_i\} = \{f\},$$

where $\{f\}$ denotes the track of the constant homotopy f . Note that the relation 1.1 still makes sense for such triples and that the inclusion $HR_B^A \rightsquigarrow H_B^A$ is an embedding.

COROLLARY 1.8. *An arrow $(\rho_\epsilon, u, \sigma_\epsilon)$ of \underline{HR}_B^A is an isomorphism if and only if u is a homotopy equivalence.*

PROOF. If $(\{\rho_\epsilon\}, u, \{\sigma_\epsilon\})$ satisfies (1.7) then it can be checked that the triple

$$(\{\rho'_\epsilon\}_{1-\epsilon} + \rho_{1-\epsilon}u, u', \{u'\sigma'_{1-\epsilon} + \psi_\epsilon\sigma\})$$

as constructed in the proof of Theorem 1.2 also satisfies condition 1.7. •

REMARK 1.9. Since the diagram 1.5 is strictly commutative, a result corresponding to Corollary 1.4 also holds in \underline{HR}_B^A .

2. THE KERVAIRE DIAGRAM.

Let

$$(2.1) \quad \begin{array}{ccc} A & \xlongequal{\quad} & A \\ \sigma \downarrow & \nearrow \chi_\epsilon & \downarrow \sigma' \\ X & \xrightarrow{\quad u \quad} & X' \\ \rho \downarrow & \nearrow \mu_\epsilon & \downarrow \rho' \\ B & \xlongequal{\quad} & B \end{array}$$

be a fixed homotopy commutative diagram over B and under A. Let

$$U = (\mu_\epsilon, u) = \begin{array}{ccc} X & \xrightarrow{\quad u \quad} & X' \\ \rho \downarrow & \nearrow \mu_\epsilon & \downarrow \rho' \\ B & \xlongequal{\quad} & B \end{array}$$

and

$$V = (u, \chi_\epsilon) = \begin{array}{ccc} A & \xlongequal{\quad} & A \\ \sigma \downarrow & \nearrow \chi_\epsilon & \downarrow \sigma' \\ X & \xrightarrow{\quad u \quad} & X' \end{array}$$

be the induced homotopy commutative diagrams over B and under A whose equivalence classes are

$$U \sim \epsilon \in \pi(X, X'/B) \quad \text{and} \quad V \sim \epsilon \in \pi(A/X, X')$$

respectively. Let us also denote by $U \sim V$ the element of $\pi(A/X, X'/B)$ represented by 2.1.

Consider the following interlocking (Kervaire) diagram of groups and pointed sets with base points as indicated.

$$\begin{array}{ccccc}
 \lambda(A/X, X'; V) & \xrightarrow{(\rho') \cdot i} & \pi_1^x(B; \rho' u) & \xrightarrow{m_u = r_B n_{uv}} & \pi(X, X'/B)_u & \xrightarrow{(\sigma) \cdot d} & \pi(A, X')_\sigma \\
 \downarrow i & & \downarrow \rho' & \searrow n_{uv} & \downarrow r_B & \downarrow d & \downarrow \sigma \\
 (2.2) \quad \lambda(X, X'/B; U) & \xrightarrow{j} & \pi_1^x(X'; u) & \xrightarrow{n^{vu}} & \pi(A/X, X'/B)_{u-v} & \xrightarrow{c} & \pi(X, X')_u \\
 \downarrow j & & \downarrow \sigma & \searrow r^A & \downarrow r^A & \downarrow c & \downarrow \rho' \\
 \lambda(X, X'/B; U) & \xrightarrow{(\sigma) \cdot j} & \pi_1^A(X'; u\sigma) & \xrightarrow{m^v = r^A n^{vu}} & \pi(A/X, X')_v & \xrightarrow{(\rho') \cdot c} & \pi(X, B)_\rho
 \end{array}$$

Here, $\lambda(A/X, X'; V)$ and $\lambda(X, X'/B; U)$ are subgroups of $\pi_1^x(X'; u)$ defined as the kernel of the induced group homomorphism

$$\sigma: \pi_1^x(X'; u) \rightarrow \pi_1^A(X'; u\sigma), \text{ respectively } \rho': \pi_1^x(X'; u) \rightarrow \pi_1^x(B; \rho' u);$$

the maps i and j are inclusions, r_B, r^A, d and c are the obvious restriction operators and n_{uv}, n^{vu} are defined by the rules


$$\begin{array}{ccc}
 (2.3) \quad \begin{array}{ccc} X & \xrightarrow{\rho' u} & B \\ \parallel & \nearrow h_t & \parallel \\ X & \xrightarrow{\rho' u} & B \end{array} & \xrightarrow{n_{uv}} & \begin{array}{ccc} A & \xrightarrow{\quad} & A \\ \sigma \downarrow & \nearrow \chi_t & \downarrow \sigma' \\ X & \xrightarrow{u} & X' \\ \rho \downarrow & \nearrow h_t & \downarrow \rho' \\ B & \xrightarrow{\quad} & B \end{array} \\
 \\
 (2.4) \quad \begin{array}{ccc} A & \xrightarrow{u \sigma} & X' \\ \parallel & \nearrow k_t & \parallel \\ A & \xrightarrow{u \sigma} & X' \end{array} & \xrightarrow{n^{vu}} & \begin{array}{ccc} A & \xrightarrow{\quad} & A \\ \sigma \downarrow & \nearrow \chi_t & \downarrow \sigma' \\ X & \xrightarrow{u} & X' \\ \rho \downarrow & \nearrow \mu_t & \downarrow \rho' \\ B & \xrightarrow{\quad} & B \end{array}
 \end{array}$$

THEOREM 2.5. *The Kervaire diagram 2.2 is commutative and its four interlocking sequences of homomorphisms and pointed maps are exact. Moreover*

$$\bullet \xrightarrow{(\cdot, \sigma)j} \bullet \xrightarrow{n^{\nu u}} \bullet$$

is exact of type E3 (see [2], 1.A) at $\pi_1^A(X'; u)$.

PROOF. We prove commutativity from $\pi_1^X(X'; u)$. We have that



The two sequences passing through $\pi_1^X(X'; u)$ are exact by definition of $\lambda(A/X, X'; V)$, $\lambda(X, X'/B; U)$ and by [2], Theorem 4.3 and its dual. Thus the following proposition remains to be proved.


PROPOSITION 2.6. *The sequence*

$$\lambda(X, X'/B; U) \xrightarrow{(\cdot, \sigma)j} \pi_1^A(X'; u) \xrightarrow{n^{\nu u}} \pi(A/X, X'/B)_{u \cdot \nu} \xrightarrow{r_B} \pi(X, X'/B)_{u \cdot \nu} \xrightarrow{(\cdot, \phi)d} \pi(A, X')_{\sigma}$$


is exact. At $\pi_1^A(X'; u)$ it is exact of type E3.

PROOF. The exactness at $\pi(X, X'/B)_{u \cdot \nu}$ is obvious.

Exactness at $\pi(A/X, X'/B)_{\sigma, \rho}$: n^{\cup} is given by 2.4. Clearly r_B applied to the composite square on the right of 2.4 yields U^{\sim} . Now suppose that




Then there exists $h_\epsilon: X \rightarrow X'$ with $h_0 = h$ such that $h_\epsilon = u$ and $\rho_\epsilon + \rho'h_\epsilon \equiv \mu_\epsilon$ and it can be checked that




as required.

Exactness at $\pi_1^*(X'; u)$: Let $\{u_\epsilon\} \in \lambda(X, X'/B; U)$, i.e., $\{u_\epsilon\}$ is in $\pi_1^*(X'; u)$ such that $\rho'u_\epsilon \equiv \rho'u$, whence
 (2.7) $\mu_\epsilon + \rho'u_\epsilon \equiv \mu_\epsilon$.

Now $n^{\cup}(\sigma)\{u_\epsilon\}$ is represented by the diagram



and in view of the relation 2.7, we have that $n^{\nu}(\sigma)(u_\epsilon) = U \sim V$. Conversely, if $\{k_\epsilon\} \in \pi_1^A(X'; u)$ is such that



then there exists a homotopy $u_\epsilon: X \rightarrow X'$ with $u_0 = u_1 = u$ such that

$$(2.8) \quad \mu_\epsilon + \rho' u_\epsilon \equiv \mu_\epsilon$$

and

$$(2.9) \quad u_{\epsilon^{-1}\sigma} + k_\epsilon + \chi_\epsilon \equiv \chi_\epsilon.$$

By 2.8 $\{u_\epsilon\}$ belongs to $\lambda(X, X'/B; U)$ and by 2.9 we have that

$$(\sigma)(u_\epsilon) = \{k_\epsilon\}.$$

Exactness of type E3 is proved by checking that

$$n^{\nu}(\{k_\epsilon\}) = n^{\nu}(\{k'_\epsilon\}) \quad \text{iff} \quad \{k'_\epsilon\}(\{k_\epsilon\})^{-1} \in \text{Im}(\sigma)$$

and applying ordinary exactness at $\pi_1^A(X'; u)$.

Applying [2], Theorem 2.13 and Lemma 2.11, we obtain

COROLLARY 2.10. *The sequence*

$$\lambda(A/X, X'; V) \times \lambda(X, X'/B; U) \xrightarrow{\theta} \pi_1^A(X'; u) \xrightarrow{\Delta} \pi(A/X, X'/B)_{\sigma^{-1}\sigma} \longrightarrow \pi(A/X, X')_{\sigma^{-1}\sigma} \times \pi(X, X'/B)_{\sigma^{-1}\sigma}$$

is exact, where $\Delta = n^{\nu}(\sigma)$ and $\theta(\alpha, \beta) = \beta^{-1}\alpha$.

Moreover the images of two elements under Δ coincide iff they belong to the same double coset of the subgroups $\lambda(X, X'/B; U)$ and $\lambda(A/X, X'; V)$.

Let $K(U, V)$ denote the set of double cosets in $\pi_1^A(X'; u)$ of the subgroups $\lambda(X, X'/B; U)$ and $\lambda(A/X, X'; V)$.

COROLLARY 2.11. *There is a bijection*

$$\pi(A/X, X'/B) \longleftrightarrow \bigcup_{\mathcal{S}} K(u, U, V),$$

where

$$S = \{(u, U, V) \mid U \sim \in d^{-1}\{u\}, V \sim \in c^{-1}\{u\}, \{u\} \in (\sigma)^{-1}\{\sigma'\} \cap (\rho')^{-1}\{\rho\}\}.$$

REMARK 2.12. The homomorphisms

$$\rho' : \pi_1^*(X'; u) \longrightarrow \pi_1^*(B; \rho'u) \quad \text{and} \quad \sigma : \pi_1^*(X'; u) \longrightarrow \pi_1^*(X'; u)$$

can, in certain special cases, be computed as discussed by Rutter [3].

3. SECONDARY COMPOSITION CLASSES.

In this final section we examine in a special case some interactions between elements of the set $\pi(A/X, X'/B)$ and secondary composition classes.

We consider pointed topological spaces A, B, X, X' and the case in which $\rho: X \rightarrow B$ and $\sigma': A \rightarrow X'$ are the trivial maps (denoted by $*$). The following operators may be defined.

$$(3.1) \quad R: \pi(A/X, X'/B) \longrightarrow \pi(X, X').$$

$$\text{Set } R\{\rho_\varepsilon, u, \sigma_\varepsilon\} = \{u\}.$$

$$(3.2) \quad M: \pi(A/X, X'/B) \longrightarrow \pi(\sigma, \rho').$$

Here $\pi(\sigma, \rho')$ refers to the homotopy pair set, for details see [1]. Set

$$M\{\rho_\varepsilon, u, \sigma_\varepsilon\} = \{*, *, \rho_\varepsilon\sigma + \rho'\sigma_\varepsilon\}.$$

Let $\alpha = \{\rho_\varepsilon, u, \sigma_\varepsilon\}$ denote an arbitrary element of $\pi(A/X, X'/B)$. Let

$$\{(\rho'), \{R\alpha\}, \{\sigma\}\} \subset \pi(\Sigma A, B)$$

denote the Toda bracket coset of $\{\sigma\}$, $\{\rho'\}$ and $\{R\alpha\}$.

We have the following result.

THEOREM 3.3. *The following are equivalent.*

$$(i) \quad 0 \in \{(\rho'), \{R\alpha\}, \{\sigma\}\}.$$

(ii)

$$M\alpha = 0.$$

PROOF. Allowing diagrams to represent elements we have

$$M\alpha = \begin{array}{ccc} A & \xrightarrow{*} & X' \\ \sigma \downarrow & \nearrow \sigma_t & \downarrow \rho' \\ X & \xrightarrow{\rho_t} & B \end{array} \sim \begin{array}{ccc} A & \xrightarrow{u\sigma} & X' \\ \sigma \downarrow & \nearrow \sigma_{t-t} & \downarrow \rho' \\ X & \xrightarrow{\rho_t} & B \\ \downarrow \rho_{t-t} & \nearrow \rho'u & \downarrow \\ X & \xrightarrow{\rho'u} & B \end{array} = \begin{array}{ccc} A & \xrightarrow{u\sigma} & X' \\ \sigma \downarrow & & \downarrow \rho' \\ X & \xrightarrow{\rho'u} & B \end{array}$$

By [2], Proposition 3.14 it follows that (i) and (ii) are equivalent. •

REFERENCES .

1. K.A. HARDIE & A.V. JANSEN, Toda brackets and the category of homotopy pairs, *Questiones Math*, 6 (1983), 107-128.
2. K.A. HARDIE & K.H. KAMPS, Exact sequence interlocking and tree homotopy theory, *Cahiers Top, et Géom, Diff*, XXVI-1 (1985), 3-31.
3. J.W. RUTTER, A homotopy classification of maps into an induced fibre space, *Topology* 6 (1967), 379-403.
4. R. SCHÖN, The Brownian classification of fiber spaces, *Arch, Math*, 39 (1982), 359-365.
5. A. STRØM, The homotopy category is a homotopy category, *Arch, Math*, 23 (1972), 435-441.
6. R.M. VOGT, A note on homotopy equivalences, *Proc, A.M.S*, 32 (1972), 627-629.

The first author acknowledges grants to the Topology Research Group from the University of Cape Town and the South African Council for Scientific and Industrial Research.

K.A. HARDIE:
 Department of Mathematics
 University of Cape Town
 RONDEBOSCH 7700
 SOUTH AFRICA

K.H. KAMPS:
 Fachbereich Mathematik u. Informatik
 Fernuniversität
 Postfach 940
 D-5800 HAGEN, Fed. Rep. of GERMANY