

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

G. J. ELLIS

C. RODRIGUEZ-FERNANDEZ

An exterior product for the homology of groups with integral coefficients modulo p

Cahiers de topologie et géométrie différentielle catégoriques, tome
30, n° 4 (1989), p. 339-343

http://www.numdam.org/item?id=CTGDC_1989__30_4_339_0

© Andrée C. Ehresmann et les auteurs, 1989, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

**AN EXTERIOR PRODUCT FOR THE
 HOMOLOGY OF GROUPS
 WITH INTEGRAL COEFFICIENTS MODULO p**

by G. J. ELLIS¹ and C. RODRIGUEZ-FERNANDEZ

RÉSUMÉ. Les auteurs généralisent la suite exacte à 8 termes de groupes d'homologie entière obtenue par Brown et Loday en une suite exacte de groupes d'homologie à coefficients dans \mathbb{Z}_p , où p est un entier non-négatif.

In this article we generalize Brown and Loday's eight term exact sequence in integral group homology [2] to an exact sequence in group homology with coefficients in \mathbb{Z}_p , where p is any nonnegative integer.

Let G be a group with a normal subgroup N , and consider $\mathbb{Z}_p = \mathbb{Z}/p\mathbb{Z}$ as a trivial G -module. We prove

THEOREM 1. *There is a natural exact sequence*

$$\begin{aligned} H_3(G, \mathbb{Z}_p) &\longrightarrow H_3(G/N, \mathbb{Z}_p) \longrightarrow \text{Ker}(\partial: N\Delta^p G \rightarrow G) \longrightarrow H_2(G, \mathbb{Z}_p) \\ &\longrightarrow H_2(G/N, \mathbb{Z}_p) \longrightarrow N/N\#_p G \longrightarrow H_1(G, \mathbb{Z}_p) \longrightarrow H_1(G/N, \mathbb{Z}_p) \longrightarrow 0. \end{aligned}$$

Here $N\#_p G$ denotes the subgroup of N generated by the elements $[n, g]$ and n^p , for $g \in G$, $n \in N$. (When x, y are elements of a group, we write $[x, y] = xyx^{-1}y^{-1}$ and $x^y = xyx^{-1}$.)

The group $N\Delta^p G$ is a new construction. It is generated by the symbols $n \wedge g$ and $\{n\}$ for $n \in N$, $g \in G$, subject to the relations

- (1) $n \wedge gh = (n \wedge g)(g n \wedge g h),$
- (2) $nm \wedge g = ({}^n m \wedge {}^n g)(n \wedge g),$

¹ This author would like to thank the University of Santiago de Compostela for its hospitality during the preparation of this article.

- (3) $n \wedge n = 1,$
- (4) $\{n\}(m \wedge g)\{n\}^{-1} = n^P m \wedge n^P g,$
- (5) $\{nm\} = \{n\} \left(\prod_{i=1}^{P-1} (n^{-1} \wedge (n^{1-P+i} m)^i) \right) \{m\},$
- (6) $[\{n\}, \{m\}] = n^P \wedge m^P,$
- (7) $\{[n, g]\} = (n \wedge m)^P$

for $g, h \in G, m, n \in N$. Note that if $N = G$, then (2) and (7) are redundant.

Clearly $N\Delta^P G$ is functorial in N and G .

The homomorphism $\partial: N\Delta^P G \rightarrow G$ is defined by

$$\partial(n \wedge g) = [n, g] \text{ and } \partial\{n\} = n^P.$$

It is routine to check that ∂ is a well-defined homomorphism, and that its image is $N *_P G$.

As an immediate consequence of Theorem 1 we have

COROLLARY 2. *There is an isomorphism*

$$H_2(G, \mathbb{Z}_p) \approx \text{Ker}(\partial: G\Delta^P G \rightarrow G)$$

Also, for any presentation $R \triangleright F \twoheadrightarrow G$ of G , there is an isomorphism

$$H_3(G, \mathbb{Z}_p) \approx \text{Ker}(\partial: R\Delta^P F \rightarrow F).$$

In order to prove Theorem 1 we need the following

LEMMA 3. *If F is a free group, then ∂ induces an isomorphism $F\Delta^P F \approx F *_P F$.*

PROOF. Recall from [2] that $N \wedge G$ is the group generated by the symbols $n \wedge g$ for $g \in G, n \in N$, subject to relations (1), (2), (3). There is thus a homomorphism $\iota: N \wedge G \rightarrow N\Delta^P G, n \wedge g \mapsto n \wedge g$. By (4) the image of ι is normal in $N\Delta^P G$. On taking $G = N = F$ we thus have a commutative diagram

$$\begin{array}{ccccc}
 F \wedge F & \longrightarrow & F\Delta^P F & \longrightarrow & F\Delta^P F / \text{Im}(\iota) \\
 \partial' \downarrow & & \downarrow \partial & & \downarrow \partial'' \\
 [F, F] & \triangleright & F *_P F & \longrightarrow & \rho F^{\text{ab}}
 \end{array}$$

where ∂'' is induced by ∂ , and where ∂' is the isomorphism proved in [2] (see [3] for an algebraic proof of this isomorphism).

AN EXTERIOR PRODUCT FOR THE HOMOLOGY ...

The homomorphism ∂'' is clearly surjective, and hence has a splitting since ρF^{ab} is free abelian and $F\Delta^P F / \text{Im}(\iota)$ is abelian by (6). This splitting is surjective because of (5). Therefore ∂'' is an isomorphism. Since the rows of the diagram are both short exact, it follows that $\partial: F\Delta^P F \rightarrow F \#_P F$ is an isomorphism. ■

In [1] the following natural exact sequence

$$\begin{aligned} H_3(G, \mathbb{Z}_P) &\rightarrow H_3(G/N, \mathbb{Z}_P) \rightarrow \text{Ker}(L_0 V_1^P(\alpha) \rightarrow G) \rightarrow H_2(G, \mathbb{Z}_P) \\ &\rightarrow H_2(G/N, \mathbb{Z}_P) \rightarrow N/N \#_P G \rightarrow H_1(G, \mathbb{Z}_P) \rightarrow H_1(G/N, \mathbb{Z}_P) \rightarrow 0. \end{aligned}$$

is obtained. Thus to prove Theorem 1 it suffices to exhibit an isomorphism $\approx: N\Delta^P G \rightarrow L_0 V_1^P(\alpha)$ such that

$$(*) \quad \begin{array}{ccc} N\Delta^P G & \xrightarrow{\quad} & G \\ \approx \downarrow & & \parallel \\ L_0 V_1^P(\alpha) & \xrightarrow{\quad} & G \end{array}$$

commutes.

For any surjection $\varepsilon: F \rightarrow G$ with F a free group let S be the kernel of the composite homomorphism

$$F \xrightarrow{\varepsilon} G \xrightarrow{\alpha} G/N.$$

Let $i: S' \rightarrow S$ be an isomorphism. Let T be the kernel of

$$\begin{pmatrix} \varepsilon i \\ \varepsilon \end{pmatrix}: S' * F \rightarrow G.$$

Then it is shown in ([1], Propositions 6.4 and 8.1) that

$$L_0 V_1^P(\alpha) = \frac{(S' * S' * F) \#_P (S' * F)}{((S' * S' * F) \#_P T) ((T \cap S' * S' * F) \#_P (S' * F))}.$$

As in [1] let $\mu: G \rightarrow F$ be any set theoretic section of $\varepsilon: F \rightarrow G$. Then μ induces a section $N \rightarrow S \approx S'$; under this section we denote the image of $n \in N$ by $\mu(n)' \in S'$. Let

$$D = ((S' * S' * F) \#_P T) ((T \cap S' * S' * F) \#_P (S' * F)).$$

With this notation, we have

LEMMA 4. *There is a homomorphism $h: N\Delta^P G \rightarrow L_0 V_1^P(\alpha)$ defined by*

$$h(n \wedge g) = [\mu(n)', \mu(g)]D, \quad h(\{n\}) = (\mu(n)')^P D.$$

PROOF. We need to show that h preserves the relations (1)-(7). By [1] clearly h preserves (1)-(3). Relation (4) is preserved because

$$\begin{aligned} \mu(nm)' \in S^*S^*F, \quad ((\mu(n))' \mu(g))^{-1} \mu(n^P g) \in T \\ ((\mu(n))' \mu(m))^{-1} \mu(n^P m)' \in T \cap S^*S^*F \end{aligned}$$

from which we see that

$$h(n^P m \wedge n^P g) = (\mu(n))' \mu(m), \mu(g) \mu(n)) \mu(n))' P D = h(\{n\}(g \wedge m)\{n\}^{-1}).$$

Relation (5) is preserved because

$$\mu(n^{-1})' \mu(n)' \in T \cap S^*S^*F$$

and

$$\mu(n)' \in S^*S^*F, \quad \mu(n^{1-P+i} m)^i (\mu(n))^{1-P+i} \mu(m)^{-1} \in T$$

which implies

$$h(\{n\} \left(\prod_{i=1}^{P-1} (n^{-1} \wedge (n^{1-P+i} m)^i) \right) \{m\}) = (\mu(n))' \mu(m))' P D.$$

Since

$$(\mu(nm))'^{-P} (\mu(n))' \mu(m))' P \in T \cap S^*S^*F$$

we have

$$h(\{n\} \left(\prod_{i=1}^{P-1} (n^{-1} \wedge (n^{1-P+i} m)^i) \right) \{m\}) = (\mu(nm))' P D = h(\{nm\}).$$

Relation (6) is preserved because we have

$$(\mu(n))' P (\mu(n^P))'^{-1} \in T \cap S^*S^*F \quad \text{and} \quad \mu(n^P)' \in S^*S^*F, \quad \mu(m^P)^{-1} (\mu(m))' P \in T.$$

Clearly relation (7) is preserved. ■

Consider the homomorphism

$$d = \begin{pmatrix} \varepsilon & i \\ \varepsilon & \end{pmatrix} : S^*F \longrightarrow G.$$

By Lemma 3 we have a homomorphism $\varphi: S^* \#_P S^* \approx S^* \Delta^P S^* \rightarrow N \Delta^P G$ defined by

$$\varphi([s'_1, s'_2]) = d(s'_1) \wedge d(s'_2), \quad \varphi((s')^P) = \{d(s')\}.$$

We therefore have a set theoretic map $g: S^*S^*F \#_P (S^*F) \rightarrow N \Delta^P G$ defined as follows (cf. [1], 8.12):

if $\prod_{i=1}^n s'_i f_i \in S^*S^*F \#_P (S^*F)$ then $\prod_{i=1}^n s'_i \in S^* \#_P S^*$ and we can define

$$g\left(\prod_{i=1}^n s'_i f_i\right) = \left(\prod_{i=1}^{n-1} (d(\prod_{j=1}^i s'_j) \wedge d(\prod_{j=1}^i f_j)) (d(\prod_{j=1}^i s'_j) \wedge d(\prod_{j=1}^{i+1} f_j))^{-1} \right) \varphi\left(\prod_{i=1}^n s'_i\right).$$

LEMMA 5. *The composite function*

$$S^*S^*F \#_P (S^*F) \xrightarrow{g} N \Delta^P G \longrightarrow N \Delta^P G / \iota(N \wedge G)$$

is a homomorphism, and induces a homomorphism

$$\psi: L_0V_1^P(\alpha) \longrightarrow N\Delta^PG/\iota(N\wedge G).$$

PROOF. The first homomorphism is clear, and certainly D is in the kernel of the first homomorphism. By Lemmas 4 and 5 we have a commutative diagram

$$\begin{array}{ccccc} N\wedge G & \xrightarrow{\iota} & N\Delta^PG & \longrightarrow & N\Delta^PG/\iota(N\wedge G) \\ h' \downarrow \approx & & h \downarrow & \nearrow \psi & \downarrow \bar{h} \\ L_0V_1^0(\alpha) & \twoheadrightarrow & L_0V_1^P(\alpha) & \twoheadrightarrow & L_0V_1^P(\alpha)/L_0V_1^0(\alpha) \end{array}$$

where h' is the restriction of $h\iota$, and is an isomorphism by Section 8 of [1]. Clearly $L_0V_1^0(\alpha)$ lies in the kernel of ψ , and so ψ induces a splitting of \bar{h} . But ψ is surjective and hence \bar{h} is an isomorphism. It follows that h is an isomorphism. It is readily seen that the above diagram (*) commutes. So Theorem 1 is proved.

REFERENCES.

1. J. BARJA & C. RODRIGUEZ, Homology groups $H_n^q(-)$ and eight-term exact sequences, *Cahiers Top. et Géom. Diff. Cat.* XXXI (1990).
2. R. BROWN & J.-L. LODAY, Van Kampen Theorems for diagrams of spaces, *Topology* 26 (3) (1987), 311-335.
3. G.J. ELLIS, Non-abelian exterior products of groups and exact sequences in the homology of groups, *Glasgow Math. J.* 29 (1987), 13-19.

Mathematics Department
University College
GALWAY
IRELAND

Departamento de Algebra
Universidad de
SANTIAGO DE COMPOSTELA
ESPAGNE