JOURNÉES ÉQUATIONS AUX DÉRIVÉES PARTIELLES

CHAOHAO GU

On the mixed partial differential equations in n independent variables

Journées Équations aux dérivées partielles (1980), p. 1-2

[<http://www.numdam.org/item?id=JEDP_1980____A7_0>](http://www.numdam.org/item?id=JEDP_1980____A7_0)

© Journées Équations aux dérivées partielles, 1980, tous droits réservés.

L'accès aux archives de la revue « Journées Équations aux dérivées partielles » (<http://www.math.sciences.univ-nantes.fr/edpa/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE MIXED PARTIAL DIFFERENTIAL EQUATIONS
IN n INDEPENDENT VARIABLES.

par GU Chaohao

The boundary value problems for mixed partial differential equations were thought to be very difficult. However, now we are able to point out a large class of well-posed boundary value problems [1]. Instead of talking about the general methods I prefer to presenting here some special cases [2], [3], [4].

Let

$$L\phi = (\delta^{ij} - x^i x^j) \partial_{ij} \phi + 2ax^i \partial_i \phi - a(a+1)\phi = f$$

$$(i, j = 1, 2, \dots, n)$$
(1)

and Ω be a boundary domain in $\mathbb{R}^n$. The operator L is elliptic inside the unit sphere and hyperbolic outside the unit sphere. Suppose that the $\partial\Omega$ is sufficiently smooth and the tangent planes of $\partial\Omega$ do not meet the unit sphere. We have

Theorem 1 : If $f \in H_s(\Omega)$ and $a > -\frac{n}{2} + s$ ($s > 1$), there exists uniquely a solution $\phi \in H_{s+1}(\Omega)$ of (1) such that the boundary conditions

$$\phi|_{\partial\Omega} = 0, \quad \frac{\partial\phi}{\partial n}|_{\partial\Omega} = 0$$
(2)

both be satisfied. In particular, if $s \geq [\frac{n}{2}] + 2$, the boundary value problem (1), (2) admits a classical solution.

Theorem 2 : If $f \in H_s(\Omega)$ and $a < -\frac{n}{2}$ ($s > 1$), there exists a solution $\phi \in H_{s+1}(\Omega)$ of (1) uniquely.

Theorem 3 : If $(a+1)$ is not a non-negative integer, then the map $L : \phi \rightarrow L\phi$ is an homeomorphism of $C^\infty(\bar{\Omega})$ to $C^\infty(\bar{\Omega})$. If $(a+1)$ is a non-negative integer, then the image of L is a subspace of codimension N_a and the dimension of $L^{-1}(0)$ is N_a , where $N_a = C_{a+1}^{n+a} + C_a^{n+a-1}$ (if $a \neq -1$) or 1 (if $a = -1$). Moreover, for an analytic f , $L^{-1}(f)$, if exists, is analytic too.

These theorems indicate that the properties of mixed equations essentially depend on the coefficients of lower order terms and the boundary conditions depend on the functional spaces of the unknown functions.

The similar results were obtained for the operator [3], [4]

$$L(a,b) = (\delta^{ij} - x^i x^j) \partial_{ij} \phi + 2ax^i \partial_i \phi - b\phi.$$
(3)

Moreover, for the quasilinear equation

$$\left(\left(1 + \frac{1}{a} \lambda(x, \phi, \partial \phi) \right) \delta^{ij} - x^i x^j \right) \partial_{ij} \phi + 2ax^i \partial_i \phi - a(a+1)\phi = f(x, \phi, \partial \phi) \quad (4)$$

with $\lambda(x, 0, 0) = 0$ we have

Theorem 4 : If a is sufficiently large the boundary value problem (2), (4) admits a classical solution ϕ .

We note that the equation (1) is hyperbolic near the boundary $\partial\Omega$ and elliptic near the origin. Hence theorem 4 is an existence theorem for an essentially quasilinear mixed equation whose elliptic region and hyperbolic region cannot be determined before the solution is obtained.

For the proof we use the theory of positive symmetric systems and some of its developments [7], [5].

In [8] some existence theorems of H_1 strong solutions were obtained.

Références

- [1] Gu Chaohao, Preprint ITP-SB-80 - 1
- [2] Gu Chaohao, Scientia Sinica 14 (1965), 1574-1580.
- [3] Gu Chaohao, Bull. Sciences 1978, 335-339.
- [4] Gu Chaohao, Preprint ITP-SB-80-20 (to appear in Commun. in P. D. E.).
- [5] Gu Chaohao, Acta Math. Sinica 21 (1978), 119-129.
- [6] Friedrichs, K. O. , Commun. Pure Appl. Math. 11 (1958), 333-418.
- [7] Gu Chaohao, Acta Math. Sinica 14 (1964), 503-516, Chinese Math. 5 (1964) 541-555.
- [8] Karatoprakliiev G. D. , Differential Equations 13 (1977), 64-75 (Russ.), 43-51 (Eng).

*
* *
*