

JOURNÉES ÉQUATIONS AUX DÉRIVÉES PARTIELLES

ARNE JENSEN

Stark hamiltonians with periodic potentials

Journées Équations aux dérivées partielles (1989), p. 1-6

http://www.numdam.org/item?id=JEDP_1989____A11_0

© Journées Équations aux dérivées partielles, 1989, tous droits réservés.

L'accès aux archives de la revue « Journées Équations aux dérivées partielles » (<http://www.math.sciences.univ-nantes.fr/edpa/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Stark Hamiltonians with Periodic Potentials

Arne Jensen

Aalborg University

Denmark

1. Introduction

Let $H_0 = -\Delta + Fx_1$ denote the free Stark Hamiltonian on $L^2(\mathbb{R}^n)$. It is essentially selfadjoint on the Schwartz space $\mathcal{S}(\mathbb{R}^n)$. Let V be a realvalued bounded function. Then $H = H_0 + V$ is selfadjoint with domain $\mathcal{D}(H) = \mathcal{D}(H_0)$. The time-dependent Schrödinger equation $i\frac{d\psi}{dt} = H\psi$, $\psi(0) = \psi_0$, has the solution $\psi(t) = e^{-itH}\psi_0$. The questions we want to consider here are the following:

- 1^o Describe the asymptotic behavior of $\psi(t) = e^{-itH}\psi_0$ as $t \rightarrow \pm\infty$. This is in a general form the basic question in scattering theory.
- 2^o Describe the spectrum $\sigma(H)$ of H in detail, i.e. classify it according to the usual categories: point spectrum, continuous spectrum, absolutely continuous and singular continuous spectrum.

For the one-dimensional case we obtain fairly complete results, see section 4. For the higher dimensional case we obtain some general results, see section 3, and for the case of a half-crystal we obtain some interesting new results, see section 5.

This presentation is a *preliminary* report on [J]₃. Concerning previous papers on Stark effect Hamiltonians with decaying potentials, we refer to the references given in [J]₂.

2. Periodic potentials and lattices

A discrete subset of \mathbb{R}^n is called a lattice, if it can be represented in the form

$$T = \{ k_1 \mathbf{a}_1 + k_2 \mathbf{a}_2 + \dots + k_n \mathbf{a}_n \mid k_1, \dots, k_n \in \mathbb{Z} \},$$

where $\mathbf{a}_1, \dots, \mathbf{a}_n$ are linearly independent vectors in \mathbb{R}^n . A function V on \mathbb{R}^n is said to be periodic with the period lattice T , if for all $\mathbf{x} \in \mathbb{R}^n$ and all $\boldsymbol{\tau} \in T$ we have $V(\mathbf{x} + \boldsymbol{\tau}) = V(\mathbf{x})$.

The position of the lattice T relative to the x_1 -axis plays an important role in our study. We introduce the following definitions. Let $\mathbf{e}_1 = (1, 0, \dots, 0) \in \mathbb{R}^n$. The inner product on \mathbb{R}^n is denoted $\langle \cdot, \cdot \rangle$.

Definition 2.1. (i) The lattice T is said to be irrational with respect to \mathbf{e}_1 , if the set $\{ \langle \mathbf{e}_1, \boldsymbol{\tau} \rangle \mid \boldsymbol{\tau} \in T \}$ is dense in \mathbb{R} .

(ii) The lattice T is said to be rational with respect to \mathbf{e}_1 , if the set $\{ \langle \mathbf{e}_1, \boldsymbol{\tau} \rangle \mid \boldsymbol{\tau} \in T \}$ is discrete in \mathbb{R} .

This is a classification, since it is easy to see that these are the only possibilities. The translation group associated to the lattice is given by $(U(\boldsymbol{\tau})f)(\mathbf{x}) = f(\mathbf{x} - \boldsymbol{\tau})$. Assume that the potential V above is periodic with period lattice T . Then we have the important relation

$$(2.1) \quad U(\boldsymbol{\tau})H U(\boldsymbol{\tau})^{-1} = H - F \langle \mathbf{e}_1, \boldsymbol{\tau} \rangle.$$

3. General spectral results

Throughout this section we assume that the potential V is a realvalued function with period lattice T .

Proposition 3.1. Assume that T is irrational with respect to \mathbf{e}_1 . Then $\sigma(H) = \mathbb{R}$.

Proof: By (2.1) $\sigma(H) = \sigma(H) - F \langle \mathbf{e}_1, \boldsymbol{\tau} \rangle$. Since $\sigma(H) \neq \emptyset$ and $\{ F \langle \mathbf{e}_1, \boldsymbol{\tau} \rangle \mid \boldsymbol{\tau} \in T \}$ is dense in \mathbb{R} , the result follows. \square

Proposition 3.2. Assume that T is rational with respect to e_1 . Assume that $\{\tau \in T \mid \langle e_1, \tau \rangle = 0\}$ is a sublattice of dimension $n-1$. Assume that $\sigma(-d^2/dx_1^2 + Fx_1 + V(x_1, \tilde{x})) = \mathbb{R}$ for a dense set of $\tilde{x} \in \mathbb{R}^{n-1}$. Then $\sigma(H) = \mathbb{R}$.

Remark 3.3. A sufficient condition for $\sigma(-d^2/dx_1^2 + Fx_1 + V(x_1, \tilde{x})) = \mathbb{R}$ is $V(x_1, \tilde{x}) = (\partial/\partial x_1)W(x_1, \tilde{x})$ for some bounded function W with two bounded derivatives, see [J]₁.

Proof: We use a direct integral decomposition with respect to the sublattice in the proposition and the the variable \tilde{x} . The proof is somewhat long, so the details are omitted. See also section 5. \square

Propositions 3.1 and 3.2 cover all cases for $n = 2$. For $n > 2$ not all cases are covered. We expect to find $\sigma(H) = \mathbb{R}$ in all cases. For a strong electric field it is easy to obtain a result on the type of spectrum.

Theorem 3.4. Assume V , $\partial V/\partial x_1$ and $\partial^2 V/\partial x_1^2$ continuous realvalued bounded functions on \mathbb{R}^n and $\alpha_0 = \inf\{F + (\partial V/\partial x_1)(\mathbf{x}) \mid \mathbf{x} \in \mathbb{R}^n\} > 0$. Assume $\sigma(H) = \mathbb{R}$. Then the spectrum is purely absolutely continuous.

Proof: This result is an immediate consequence of Mourre's commutator method [M]. We use the conjugate operator $A = i\partial/\partial x_1$. The assumption implies that we have the Mourre commutator estimate

$$i[H, A] = F + \partial V/\partial x_1(x) \geq \alpha_0 I.$$

Furthermore, the second commutator $i[i[H, A], A] = \partial^2 V/\partial x_1^2$ is a bounded operator on $L^2(\mathbb{R}^n)$ by our assumption. Thus all the essential conditions for applying Mourre result are verified. The remaining technical conditions are easily verified. \square

4. One-dimensional Stark Hamiltonians

In the one-dimensional case there are fairly complete answers to questions 1* and 2* in section 1. We shall briefly recall these results from [J]₁. Let us recall that the basic objects in the scattering theory for the pair of operators H and H_0 are the wave operators $W_{\pm}(H, H_0) = s\text{-}\lim_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0}$. One asks whether these

operators exist and are complete, i.e. $\text{Ran}(W_{\pm}) = \mathcal{H}_p(H)^{\perp}$, the orthogonal complement to the closed subspace $\mathcal{H}_p(H)$ spanned by the L^2 -eigenfunctions of H . The point spectrum of H is denoted $\sigma_p(H)$.

Theorem 4.1. ($n=1$) Assume $V \in C^2(\mathbb{R})$, V periodic with period a , and $\int_0^a V(x)dx = 0$. Then $W_{\pm}(H, H_0)$ exist and are unitary.

Theorem 4.2. ($n=1$) Assume $V = V_1 + V_2$, where V_1 satisfies the assumptions of the previous theorem and V_2 satisfies $V_2(x) = O(|x|^{1-\epsilon})$ as $x \rightarrow \infty$, $V_2(x) = O(|x|^{-1/2-\epsilon})$ as $x \rightarrow -\infty$ for some $\epsilon > 0$. Then $W_{\pm}(H, H_0)$ exist and are complete. Furthermore, $\sigma_p(H)$ is discrete in \mathbb{R} .

Theorem 4.3. ($n=1$) Assume $V = W''$, where W is a realvalued bounded function with four bounded derivatives. Then $W_{\pm}(H, H_0)$ exist and are unitary.

Theorem 4.1 is of the expected type. It shows that the crystal becomes "transparent" with respect to the time evolution, if one waits a long time. Theorem 4.2 shows that we can add "impurities" (in the form of V_2) and retain the same result, except the possible occurrence of a discrete set of embedded eigenvalues.

Theorem 4.3 shows that the same result holds, even for sums of periodic potentials and for a large class of almost-periodic functions. For example, one can take

$$V(x) = \int_{\mathbb{R}} e^{i\omega x} d\mu(\omega)$$

where μ is a Borel measure satisfying

$$\int_{\mathbb{R}} (\omega^{-2} + \omega^2) d|\mu|(\omega) < \infty.$$

As a special case we can take

$$V(x) = \sum_{k=1}^{\infty} a_k \sin(\omega_k x)$$

with

$$\sum_{k=1}^{\infty} |a_k|(\omega^{-2} + \omega^2) < \infty.$$

5. The half-crystal model

We now consider the case where the crystal fills up half the space.

Half-solids have been briefly considered in [S]. Here we add a constant electric field orthogonal to the surface directed into the empty part of space. The results below show that after a long time an electron will eventually move freely, irrespective of the initial position.

Let V_1 be a periodic function on \mathbb{R}^n with period lattice $T = \mathbb{Z} \times \tilde{T}$, where \tilde{T} is a lattice in \mathbb{R}^{n-1} . We assume $V_1 \in C^2(\mathbb{R}^n)$. Let χ be a cutoff function, i.e. $\chi \in C^\infty(\mathbb{R})$ realvalued, $0 \leq \chi(x_1) \leq 1$, $\chi(x_1) = 0$ for $x_1 < -\delta$, and $\chi(x_1) = 1$ for $x_1 > \delta$, where $\delta > 0$ is a fixed parameter. We take as our potential

$$V(\mathbf{x}) = \chi(x_1)V_1(\mathbf{x}).$$

The main result is the following

Theorem 5.1. ($n \geq 2$) Let V satisfy the assumptions above. Then $W_{\pm}(H, H_0)$ exist and are unitary. Consequently, $\sigma(H) = \sigma_{ac}(H) = \mathbb{R}$.

The proof of this theorem will only be sketched. Let $F_{\tilde{T}}$ denote a fundamental region for the lattice \tilde{T} , chosen diffeomorphic to the $n-1$ -dimensional torus \mathbb{T}^{n-1} . The dual lattice is denoted \tilde{T}^* and a fundamental region $F_{\tilde{T}^*}$, again chosen diffeomorphic to \mathbb{T}^{n-1} . We now use the Floquet-Bloch reduction, see for example [Sk] for details. There exists a unitary operator $W_{\tilde{T}}$ from $L^2(\mathbb{R}^n)$ to the direct integral space $\mathcal{H} = \int^{\oplus} \mathcal{H}(k) dk$, where k varies over $F_{\tilde{T}^*}$. The operator H is transformed into $W_{\tilde{T}} H W_{\tilde{T}}^{-1} = \int^{\oplus} H(k) dk$. In our case we do not reduce in x_1 , so we have $\mathcal{H}(k) = L^2(\mathbb{R}) \otimes L^2(F_{\tilde{T}})$ and $H(k) = \hbar_0 \otimes I_2 + I_1 \otimes Q(k) + V(x_1, \tilde{x})$ with $\hbar_0 = -(d^2/dx_1^2) + Fx_1$ on $L^2(\mathbb{R})$ and $Q(k) = (-i\nabla_{\tilde{x}} - k)^2$ on $L^2(F_{\tilde{T}})$ with periodic boundary conditions. Here $k \in F_{\tilde{T}^*}$. The main step is the following lemma.

Lemma 5.2. The wave operators $\mathfrak{W}_{\pm}(H(k), H_0(k))$ exist and are unitary on $\mathcal{H}(k)$, $k \in F_{\Gamma}^*$.

To prove this lemma, we verify the conditions in the abstract theorems in [J]₂. The proof of absence of embedded eigenvalues requires a separate argument. Details can be found in [J]₃.

References

- [J]₁ A. Jensen, *Asymptotic Completeness for a New Class of Stark Effect Hamiltonians*. Commun. Math. Phys. **107** (1986), 21–28.
- [J]₂ A. Jensen, *Scattering Theory for Hamiltonians with Stark Effect*. Ann. Inst. Henri Poincaré, Phys. Théor. **46** (1987), 383–395.
- [J]₃ A. Jensen, in preparation.
- [M] E. Mourre, *Absence of Singular Continuous Spectrum for Certain Self-Adjoint Operators*. Commun. Math. Phys. **78** (1981), 391–408.
- [S] B. Simon, *Phase Space Analysis of Simple Scattering Systems: Extensions of some Work of Enss*. Duke Math. J. **46** (1979), 119–168.
- [Sk] M. M. Skriganov, *Geometric and Arithmetic Methods in the Spectral Theory of Multidimensional Periodic Operators*. English translation: Proc. Steklov Inst. Math. vol. 171, 1987.