A. D. POLLINGTON

R. C. VAUGHAN

The k -dimensional Duffin and Schaeffer conjecture

Séminaire de Théorie des Nombres de Bordeaux, tome 1, n° 1 (1989),
p. 81-88

<http://www.numdam.org/item?id=JTNB_1989__1_1_81_0>

© Université Bordeaux 1, 1989, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

The k -dimensional Duffin and Schaeffer conjecture.

par A.D. POLLINGTON AND R.C. VAUGHAN

Résumé — Nous montrons que la conjecture de Duffin et Schaeffer est vraie en toute dimension supérieure à 1.

Abstract — We show that the Duffin and Schaeffer conjecture holds in all dimensions greater than one.

In 1941 Duffin and Schaeffer [1] made the following conjecture:

CONJECTURE. Let $\{\alpha_n\}$ denote a sequence of real numbers with

$$0 \leq \alpha_n < \frac{1}{2}$$

then the inequalities

$$(1) \quad |nx - a| < \alpha_n, \quad (a, n) = 1,$$

have infinitely many solutions for almost all x if and only if

$$(2) \quad \sum_{n=1}^{\infty} \frac{\alpha_n \varphi(n)}{n}$$

diverges.

If (2) converges then it easily follows from the Borel-Cantelli lemma that the set of x 's satisfying infinitely many of the inequalities (1) has Lebesgues measure zero. Duffin and Schaeffer gave conditions on the α_n for which the conjecture is true and showed that the condition $(a, n) = 1$ is necessary. In 1970 Erdős [2] showed that the conjecture holds if $\alpha_n = \frac{\epsilon}{n}$ or 0. This was later extended by Vaaler [6] who showed that $\alpha_n = O(\frac{1}{n})$ is sufficient. In his book on metric number theory [5] Sprindzuk considers a k -dimensional analogue of the conjecture in which (1) is replaced by

$$(3) \quad \max(|x_1 n - a_1|, \dots, |x_k n - a_k|) < \alpha_n \quad (a_i, n) = 1, \quad i = 1, \dots, k$$

and (2) by

$$(4) \quad \sum_{n=1}^{\infty} \left(\frac{\alpha_n \varphi(n)}{n} \right)^k$$

where the measure is now k -dimensional Lebesgues measure. He states that the study of such approximations subject to the conditions $(a_1, n) = \cdots = (a_k, n) = 1$ is probably a problem of the same degree of complexity as the case $n = 1$. This appears not to be the case. For we can now prove the k -dimensional analogue of the Duffin and Schaeffer conjecture.

We prove the following result:

THEOREM. *Let $k > 1$ and let $\{\alpha_n\}$ denote a sequence of real numbers with*

$$0 \leq \alpha_n < \frac{1}{2}$$

and suppose that

$$\sum_{n=1}^{\infty} \left(\frac{\alpha_n \varphi(n)}{n} \right)^k$$

diverges. Then the inequalities

$$\max(|x_1 n - a_1|, \dots, |x_k n - a_k|) < \alpha_n \quad (a_i, n) = 1, \quad i = 1, \dots, k$$

have infinitely many solutions for almost all $x \in R^k$.

Unfortunately our method does not readily extend to the case $k = 1$. We are able find some more sequences $\{\alpha_n\}$ for which the conjecture holds, for example if $\alpha_n = 0$ or $1 \ll \alpha_n$, but not to settle the dimension one case. In the k -dimensional case, $k \geq 2$, Vilchinski has previously shown that we may take $\alpha_n = O(n^{-\gamma})$ for any $\gamma > 0$.

Put

$$E_n = E_n^{(1)} \times \cdots \times E_n^{(k)}$$

where

$$(5) \quad E_n^{(i)} = \bigcup_{\substack{1 \leq a_i \leq n \\ (a_i, n) = 1}} \left(\frac{a_i - \alpha_n}{n}, \frac{a_i + \alpha_n}{n} \right).$$

Then E_n is the set counted in (3) and

$$(6) \quad \lambda_k(E_n) = \left(\frac{2\alpha_n \varphi(n)}{n}\right)^k.$$

Thus (4) becomes

$$(7) \quad \sum_{k=1}^{\infty} \lambda_k(E_n)$$

diverges.

We are interested in

$$(8) \quad \lambda_k \left(\bigcap_{m=1}^{\infty} \bigcup_{n=m}^{\infty} E_n \right).$$

By an ergodic theorem of Gallagher [3], see Sprindzuk [5], this is either zero or one. Gallagher proves his result for dimension one. The corresponding k -dimensional result is proved by Vilchinski [7]. In order to prove the theorem it therefore suffices to show that (8) is not zero.

Since (7) diverges, and $\lambda_k(E_n) \rightarrow 0$ as $n \rightarrow \infty$, given any $1 > \eta > 0$, for every N we can find a finite set Z so that if $z \in Z$ then $z > N$, and

$$(9) \quad \eta^2 < \Lambda(Z) = \sum_{n \in Z} \lambda_k(E_n) < \eta.$$

By the Cauchy-Schwarz inequality

$$(10) \quad \lambda_k \left(\bigcup_{n \in Z} E_n \right) \geq \frac{(\sum_n \lambda_k(E_n))^2}{(\sum_n \sum_m \lambda_k(E_n \cap E_m))},$$

this is Lemma 5 of Sprindzuk [5]. So provided there is some absolute constant c for which

$$(11) \quad \sum_{\substack{n \neq m \\ m, n \in Z}} \lambda_k(E_n \cap E_m) \leq c \sum_{n \in Z} \lambda_k(E_n)$$

the theorem is proved.

Thus we need to bound

$$\sum_{\substack{n \neq m \\ m, n \in \mathbb{Z}}} \lambda_k(E_n \cap E_m).$$

In the subsequent discussion constants in the Vinogradov $\ll$ symbol depend at most on the dimension k .

Note that

$$(12) \quad \lambda_k(E_n) = \prod_{i=1}^k \lambda(E_n^{(i)})$$

and

$$(13) \quad \lambda_k(E_n \cap E_m) = \prod_{i=1}^k \lambda(E_n^{(i)} \cap E_m^{(i)}).$$

We now concentrate on $k = 1$ and use (12) and (13) to obtain a bound for $k > 1$.

We wish to obtain an estimate for

$$\lambda(E_n \cap E_m) \quad \text{with } n > m.$$

Put

$$(14) \quad \delta = \min\left(\frac{\alpha_n}{n}, \frac{\alpha_m}{m}\right) \quad \text{and} \quad \Delta = \max\left(\frac{\alpha_n}{n}, \frac{\alpha_m}{m}\right),$$

and

$$d = (m, n) \quad m' = \frac{m}{d} \quad n' = \frac{n}{d}.$$

Then

$$(15) \quad \lambda(E_n \cap E_m) \leq 2\delta \sum_{\substack{|\frac{a}{n} - \frac{b}{m}| < 2\Delta \\ (a, n) = 1 \\ (b, m) = 1}} 1.$$

By estimating

$$(16) \quad \sum_{\substack{|\frac{a}{n} - \frac{b}{m}| < 2\Delta \\ (a, n) = 1 \\ (b, m) = 1}} 1$$

we obtain

LEMMA 1. With E_n as above and $d = (m, n)$ there is an absolute constant c so that

$$(17) \quad \lambda(E_n \cap E_m) \leq c\lambda(E_n)\lambda(E_m) \prod_{\substack{p|m'n' \\ p > \frac{2mn\Delta}{d}}} (1 - \frac{1}{p})^{-1}.$$

Let $g(n)$ be defined as the least number so that

$$\sum_{\substack{p|n \\ p > g(n)}} \frac{1}{p} < 2$$

Put

$$t = \max(g(n'), g(m')).$$

Then by Lemma 1

$$(18) \quad \lambda(E_n \cap E_m) \ll \lambda(E_n)\lambda(E_m) \prod_{\substack{p|m'n' \\ \frac{2mn\Delta}{d} < p < t}} (1 - \frac{1}{p})^{-1}.$$

In particular if $\frac{2mn\Delta}{d} \geq t$ then

$$(19) \quad \lambda(E_n \cap E_m) \ll \lambda(E_n)\lambda(E_m).$$

Now

$$(20) \quad \prod_{\substack{p|\frac{nm}{d^2} \\ p < t}} (1 - \frac{1}{p})^{-1} \leq \prod_{p < t} (1 - \frac{1}{p})^{-1} \ll \log t.$$

We now return to the k -dimensional case. By (12) and (13)

$$(21) \quad \sum_{\substack{n \neq m \\ \log t \leq \Lambda(Z)^{-1/k}}} \lambda_k(E_n \cap E_m) \ll \Lambda(Z)^{-1} \sum_{n \neq m} \lambda_k(E_n) \lambda_k(E_m) \\ \ll \Lambda(Z).$$

From now on we shall assume that

$$\log t > \Lambda(Z)^{-1/k}.$$

We distinguish four cases:

$$(a) \quad \Delta = \frac{\alpha_m}{m}, \quad \alpha_n < \alpha_m$$

$$(b) \quad \Delta = \frac{\alpha_m}{m}, \quad \alpha_n \geq \alpha_m$$

$$(c) \quad \Delta = \frac{\alpha_n}{n}, \quad \alpha_n < \alpha_m$$

$$(d) \quad \Delta = \frac{\alpha_n}{n}, \quad \alpha_n \geq \alpha_m.$$

Recall that $m < n$ so (c) is impossible. We will consider the other three cases separately. The first case corresponds to the situation considered by Erdős .

Case (a).

We have

$$\frac{mn\Delta}{d} = n'\alpha_m.$$

We need to consider pairs m, n with

$$(22) \quad 1 < n'\alpha_m < t$$

Let $A_{u,v}$ denote that part of the sum

$$\sum_{\substack{m \neq n \\ \text{case (a)}}} \lambda_k(E_n \cap E_m)$$

for which $g(m') = u$ and $g(n') = v$. Then, by Lemma 1 and (22)

$$A_{u,v} \ll \log^k t \sum_m \lambda_k(E_m) \sum_{\substack{(u) \\ m' | m \\ 1 \leq m' < t\alpha_m^{-1}}} \sum_{\substack{(v) \\ \alpha_m^{-1} \leq n' < t\alpha_m^{-1} \\ n = n'd}} \lambda_k(E_n).$$

Where $\sum_{(u)}$ means the sum of those m' with $g(m') = u$ and $m = m'd$. Since

$$\lambda_k(E_n) = \left(2\alpha_n \frac{\varphi(n)}{n} \right)^k \ll \alpha_m^k$$

then

$$A_{u,v} \ll \log^k t \sum_m \lambda_k(E_m) \alpha_m^k \sum_{\substack{m'|m \\ 1 \leq m' < t\alpha_m^{-1}}} \sum_{\substack{(u) \\ \alpha_m^{-1} \leq n' < t\alpha_m^{-1} \\ n=n'd}} 1.$$

To estimate the inner sum we will use the following result due to Erdős which also appears in Vaaler [6]. Let $N(\xi, v, x)$ be the number of integers $n \leq x$ which satisfy

$$\sum_{\substack{p|n \\ p \geq v}} \frac{1}{p} \geq \xi.$$

Then

LEMMA 2. *For any $\epsilon > 0$ and any $\xi > 0$ there exists a positive integer $v_0 = v_0(\xi, \epsilon)$ such that for all $x > 0$ and all $v > v_0$,*

$$N(\xi, v, x) \leq x \exp\{-v^{\beta(1-\epsilon)}\}$$

where $\log \beta = \xi$.

Applying this result with $\xi = 1$ and $\beta(1 - \epsilon) = \frac{3}{2}$ we have

COROLLARY. *Given $x > 0$ and $v \geq 1$*

$$(23) \quad N(1, v, x) \ll x \exp(-v^{\frac{3}{2}}).$$

Using (23) and partial summation

$$\sum_{\substack{(v) \\ 1 \leq n' \leq t\alpha_m^{-1}}} \ll t\alpha_m^{-1} \exp(-v^{\frac{3}{2}}).$$

Applying (23) again

$$A_{u,v} \ll t^2 \log^k t \exp(-u^{\frac{3}{2}}) \exp(-v^{\frac{3}{2}}) \sum_m \lambda_k(E_m),$$

since $k \geq 2$. Then summing over u and v we have

$$(24) \quad \sum_{\substack{m \neq n \\ \text{case}(a)}} \lambda_k(E_n \cap E_m) \ll \Lambda(Z)$$

if η is sufficiently small.

The other cases are similar and we have

$$(25) \quad \sum_{n \neq m} \lambda_k(E_n \cap E_m) \leq c \sum_n \lambda_k(E_n)$$

where c is a constant depending only on k . This completes the proof of the Theorem .

REFERENCES

1. R.J. Duffin and A.C. Schaeffer, *Khinchine's problem in metric Diophantine approximation*, Duke Math. J. **8** (1941), 243–255.
2. P. Erdős, *On the distribution of convergents of almost all real numbers*, J. Number Theory **2** (1970), 425–441.
3. P.X. Gallagher, *Approximation by reduced fractions*, J. Math. Soc. of Japan **13** (1961), 342–345.
4. Halberstam and Richert, "Sieve methods," Academic Press, London, 1974.
5. V.G. Sprindzuk, "Metric theory of Diophantine approximations," V.H. Winston and Sons, Washington D.C., 1979.
6. J.D. Vaaler, *On the metric theory of Diophantine approximation*, Pacific J. Math. **76** (1978), 527–539.
7. V.T. Vilchinski, *On simultaneous approximations*, Vesti Akad Navuk BSSR Ser Fiz.-Mat (1981), 41–47.
8. ———, *The Duffin and Schaeffer conjecture and simultaneous approximations*, Dokl. Akad. Nauk BSSR **25** (1981), 780–783.

Mots clefs: Diophantine approximation, k -dimensional, Lebesgues measure, Duffin and Schaeffer conjecture.

Imperial College
LONDON.