

JACEK POMYKAŁA

**Non-vanishing of  $n$ -th derivatives of twisted elliptic  
 $L$ -functions in the critical point**

*Journal de Théorie des Nombres de Bordeaux*, tome 9, n° 1 (1997), p. 1-10

[http://www.numdam.org/item?id=JTNB\\_1997\\_\\_9\\_1\\_1\\_0](http://www.numdam.org/item?id=JTNB_1997__9_1_1_0)

© Université Bordeaux 1, 1997, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

**Non-vanishing of  $n$ -th derivatives of twisted  
elliptic  $L$ -functions in the critical point**

par JACEK POMYKAŁA

RÉSUMÉ. On note  $L^{(n)}(s, E)$  la dérivée  $n$ -ième de la série  $L$  de Hasse-Weil associée à une courbe elliptique modulaire  $E$  définie sur  $\mathbb{Q}$ . On évalue dans cet article le nombre de tordues  $E_d$ ,  $d \leq D$ , de la courbe elliptique  $E$  telles que  $L^{(n)}(1, E_d) \neq 0$ .

ABSTRACT. Let  $E$  be a modular elliptic curve over  $\mathbb{Q}$  and  $L^{(n)}(s, E)$  denote the  $n$ -th derivative of its Hasse-Weil  $L$ -series. We estimate the number of twisted elliptic curves  $E_d$ ,  $d \leq D$  such that  $L^{(n)}(1, E_d) \neq 0$ .

## 1. Introduction

Let  $E = E/\mathbb{Q}$  be a modular elliptic curve over  $\mathbb{Q}$  with conductor  $N$  defined by the Weierstrass equation  $y^2 = f(x)$  and let  $\mathbb{D}$  be defined as follows:

$$\mathbb{D} = \{d - \text{square-free} : 0 < d \equiv -\nu^2 \pmod{4N} \text{ for some } \nu \text{ prime to } 4N\}.$$

For any  $d \in \mathbb{D}$  we consider the twisted elliptic curve  $E_d$  given by the equation  $-dy^2 = w(x)$ . We denote by  $L(s, E)$  and  $L(s, E_d)$  the Hasse-Weil  $L$ -functions associated to the curves  $E$  and  $E_d$ , respectively.

The celebrated Birch and Swinnerton-Dyer conjecture (see [B-S]) asserts that the rank of the group of rational points of  $E/\mathbb{Q}$  is equal to the vanishing order of the associated Hasse-Weil function  $L(s, E)$  at  $s = 1$ . Kolyvagin [Ko] has proved that

$E/\mathbb{Q}$  has rank equal to zero if:

- 1)  $L(E, 1) \neq 0$ ,
- 2) There exists  $d \in \mathbb{D}$  such that  $L(s, E_d)$  has a simple zero at  $s = 1$ .

$E/\mathbb{Q}$  has rank equal to one if:

- 1')  $L(s, E)$  has a simple zero at  $s = 1$ .
- 2') There exists  $d \in \mathbb{D}$  such that  $L(1, E_d) \neq 0$ .

The condition 2') is true according to Waldspurger's theorem (see [Wa]).

The condition 2) has been proved to hold for infinitely many  $d \in \mathbb{D}$  (see [B-F-H], [M-M]).

Iwaniec [Iw] has also proved a quantitative result on this condition. Let

$$N(D) = \#\{d \in \mathbb{D}, d \leq D : L'(1, E_d) \neq 0\}.$$

He has obtained the estimate

$$N(D) \gg D^{\frac{2}{3}-\varepsilon}$$

with arbitrary  $\varepsilon > 0$ .

The above exponent is improved to  $1-\varepsilon$  in [P-P]. Here we will generalize this result to the  $n$ -th derivative of  $L(s, E_d)$  where  $n$  is an arbitrary non-negative integer.

In this connection let  $w = w(E)$  be the sign in the functional equation (see (1)). We define

$$N_n(D) = \#\{d \in \mathbb{D}, d \leq D : L^{(n)}(1, E_d) \neq 0\}.$$

We will prove

**THEOREM 1.** *Let  $\varepsilon$  be an arbitrary positive real number and  $n$  be a fixed non-negative integer. Then we have as  $D$  tends to infinity*

$$N_n(D) \gg (n + |w - 1|)D^{1-\varepsilon}.$$

where the constant implied in the symbol  $\gg$  depends on  $n$  and  $\varepsilon$ .

Our result is based on a recent large sieve type estimates over fundamental discriminants obtained by Heath-Brown [HB] (see Theorem 3 of [P-P]) and the method applied by Iwaniec in [Iw].

## 2. Outline of the proofs

For  $\operatorname{Re} s > 1$  the corresponding  $L$ -functions are given by

$$L(s, E) = \sum_{k=1}^{\infty} a_k k^{-s}, \quad L(s, E_d) = \sum_{k=1}^{\infty} \chi_d(k) a_k k^{-s},$$

where  $\chi_d(\cdot) = \left(\frac{-d}{\cdot}\right)$  (the Kronecker symbol) is a real character to modulus  $d$  prime to  $4N$ . They have the analytic prolongation to the whole complex plane, where they satisfy the equations

$$\left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma(s) L(s, E) = w \left(\frac{\sqrt{N}}{2\pi}\right)^{2-s} \Gamma(2-s) L(E, 2-s), \quad (1)$$

$$\left(\frac{d\sqrt{N}}{2\pi}\right)^s \Gamma(s) L(s, E_d) = w_d \left(\frac{d\sqrt{N}}{2\pi}\right)^{2-s} \Gamma(2-s) L(2-s, E_d), \quad (2)$$

where  $w$  and  $w_d = \pm 1$  are suitable constants depending on the reduction of  $E$  at the primes dividing  $N$ , which satisfy the equality

$$w_d = w \chi_d(-N). \quad (3)$$

From (1) and (2) it follows that

$$w = (-1)^{\operatorname{ord}_{s=1} L(s, E)}, \quad w_d = (-1)^{\operatorname{ord}_{s=1} L(s, E_d)}$$

Theorem 1 is a consequence of the following two theorems.

**THEOREM 2.** *For any integer  $n \geq 0$  we have the asymptotical equality*

$$\sum_{d \leq D}^* L^{(n)}(1, E_d) F\left(\frac{d}{Y}\right) \sim c_n L(1) \int F(t) dt D(\ln D)^n, \quad \text{as } D \rightarrow \infty,$$

where  $F$  is a smooth function compactly supported in  $\mathbb{R}^+$  with positive mean-value and the star (\*) above means that the summation is restricted to  $d \in \mathbb{D}$ . The (nonzero) constant  $L(1)$  is described in [Iw], while

$$c_n = \begin{cases} \frac{1}{2}(1-w)c & \text{if } n = 0 \\ w(-2)^{n-1}c & \text{if } n \geq 1 \end{cases} \quad (4)$$

$$c = \frac{3}{\pi^2 N} \prod_{p|4N} \left(1 - \frac{1}{p^2}\right)^{-1} \times \quad (5)$$

$$\times \#\{d \pmod{4N} : d \equiv -\nu^2 \pmod{4N}, (\nu, 4N) = 1\}.$$

The proof follows the idea exploited in [Iw] (cf. also [M-M] and [P-S]). We postpone it to sect.3.

Along the same lines as Theorem 3 of [P-P] we obtain

**THEOREM 3.** *Let  $\varepsilon > 0$  and  $n$  be a fixed non-negative integer. Then we have*

$$\sum_{d \leq D}^* |L^{(n)}(1, E_d)|^2 \ll_{\varepsilon} D^{1+\varepsilon}$$

Now from Theorem 2 and Theorem 3 we obtain by an application of the Cauchy-Schwarz inequality that

$$(n + |w-1|)D(\ln D)^n \ll \sum_{d \leq D}^* |L^{(n)}(1, E_d)| \leq \left( \sum_{d \leq D}^* |L^{(n)}(1, E_d)|^2 \right)^{\frac{1}{2}} N_n(D)^{\frac{1}{2}}$$

hence

$$N_n(D) \gg (n + |w-1|)D^{1-2\varepsilon}$$

and Theorem 1 follows.

### 3. Proof of Theorem 2

For  $n \geq 0$  we introduce the approximate functions

$$A_n(X, \chi_d) = \sum_{m=1}^{\infty} a_m \frac{\chi_d(m)}{m} V_n\left(\frac{2\pi m}{X}\right) \quad (6)$$

with

$$V_n(X) = \frac{1}{2\pi i} \int_{(3/4)} \frac{\Gamma(s)}{s^n} X^{-s} ds, \quad (7)$$

where we integrate over the line  $\operatorname{Re} s = \frac{3}{4}$ .

By the functional equation (2) we have

$$L(1+s, E_d) = w_d \left( \frac{d\sqrt{N}}{2\pi} \right)^{-2s} L(1-s, E_d) \frac{\Gamma(1-s)}{\Gamma(1+s)}.$$

Therefore

$$\begin{aligned} \frac{1}{2\pi i} \int_{(-3/4)} L(1+s, E_d) \frac{\Gamma(s)}{s^n} \left( \frac{X}{2\pi} \right)^s ds \\ &= \frac{w_d}{2\pi i} \int_{(-3/4)} L(1-s, E_d) \frac{\Gamma(1-s)}{\Gamma(1+s)} \frac{\Gamma(s)}{s^n} \left( \frac{2\pi X}{d^2 N} \right)^s ds \\ &= -\frac{w_d}{2\pi i} \int_{(-3/4)} L(1-s, E_d) \frac{\Gamma(-s)}{s^n} \left( \frac{2\pi X}{d^2 N} \right)^s ds \\ &= (-1)^{n+1} \frac{w_d}{2\pi i} \int_{(3/4)} L(1+s, E_d) \frac{\Gamma(s)}{s^n} \left( \frac{2\pi}{d^2 N/X} \right)^{-s} ds \\ &= (-1)^{n+1} w_d A_n \left( \frac{d^2 N}{X}, \chi_d \right). \end{aligned}$$

Hence defining

$$G_n(s, d, X) = L(1+s, E_d) \frac{\Gamma(s)}{s^n} \left( \frac{X}{2\pi} \right)^s$$

we obtain by the Cauchy theorem

$$\begin{aligned} \operatorname{res}_{s=0} G_n(s, d, X) &= \frac{1}{2\pi i} \int_{(3/4)} G_n(s, d, X) ds - \frac{1}{2\pi i} \int_{(-3/4)} G_n(s, d, X) ds \\ &= A_n(X, \chi_d) + (-1)^n w_d A_n \left( \frac{d^2 N}{X}, \chi_d \right). \end{aligned}$$

By the definition of  $\mathbb{D}$  we have  $w_d = w\chi_d(-N) = -w$  for any  $d \in \mathbb{D}$ . Hence letting  $X = d\sqrt{N}$  we obtain

$$\operatorname{res}_{s=0} G_n(s, d, d\sqrt{N}) = \begin{cases} 2A_n(d\sqrt{N}, \chi_d) & \text{if } n \equiv \frac{w+1}{2} \pmod{2} \\ 0 & \text{if } n \equiv \frac{w-1}{2} \pmod{2}. \end{cases} \quad (8)$$

On the other hand the residuum of  $G_n$  may be expressed in terms of the derivatives  $L^{(k)}(s, E_d)$ ,  $k = 0, 1, 2, \dots$ , by means of the Laurent expansions

$$\Gamma(s) = s^{-1} + \gamma_0 + \gamma_1 s + \dots,$$

$$\left(\frac{X}{2\pi}\right)^s = \sum_{k=0}^{\infty} b_k s^k,$$

where  $-\gamma_0$  is the Euler constant and  $b_k = b_k(X) = \frac{1}{k!} \left(\ln \frac{X}{2\pi}\right)^k$ .

Namely we prove

**Lemma.** For any  $n \geq 0$  we have

$$\begin{aligned} \operatorname{res}_{s=0} G_n(s, d, X) &= b_0 \frac{L^{(n)}(1, E_d)}{n!} \\ &+ \sum_{l=0}^{n-1} \left( b_{l+1} + \sum_{m=0}^l \gamma_m b_{l-m} \right) \frac{L^{(n-1-l)}(1, E_d)}{(n-1-l)!}. \end{aligned}$$

**Proof.** We have

$$\begin{aligned} G_n(s, d, X) &= s^{-n} \left( \sum_{k=0}^{\infty} \frac{L^{(k)}(1, E_d)}{k!} s^k \right) \left( s^{-1} + \sum_{k=0}^{\infty} \gamma_k s^k \right) \left( \sum_{k=0}^{\infty} b_k s^k \right) \\ &= s^{-n} \left( \sum_{k=0}^{\infty} \frac{L^{(k)}(1, E_d)}{k!} s^k \right) \left( \sum_{k=0}^{\infty} b_k s^{k-1} + \sum_{k=0}^{\infty} f_k s^k \right), \end{aligned}$$

where

$$f_k = \sum_{l=0}^k b_l \gamma_{k-l}.$$

The expression in the second bracket is equal to

$$\frac{b_0}{s} + \sum_{k=0}^{\infty} b_{k+1} s^k + \sum_{k=0}^{\infty} f_k s^k = \frac{b_0}{s} + \sum_{k=0}^{\infty} (f_k + b_{k+1}) s^k.$$

Hence

$$G_n(s, d, X) = s^{-n} \left( \frac{b_0}{s} \sum_{k=0}^{\infty} \frac{L^{(k)}(1, E_d)}{k!} s^k + \sum_{k=0}^{\infty} c_k s^k \right),$$

where

$$c_k = \sum_{l=0}^k (f_l + b_{l+1}) \frac{L^{(k-l)}(1, E_d)}{(k-l)!}.$$

Therefore

$$\begin{aligned} \operatorname{res}_{s=0} G_n(s, d, X) &= b_0 \frac{L^{(n)}(1, E_d)}{n!} + c_{n-1} \\ &= b_0 \frac{L^{(n)}(1, E_d)}{n!} + \sum_{l=0}^{n-1} (b_{l+1} + f_l) \frac{L^{(n-1-l)}(1, E_d)}{(n-1-l)!}, \end{aligned}$$

as required.

Since  $b_l(X) = \frac{1}{l!} \left( \ln \frac{X}{2\pi} \right)^l$  we obtain asymptotically

$$\operatorname{res}_{s=0} G_n(s, d, X) \sim \sum_{l=0}^n b_l(X) \frac{L^{(n-l)}(1, E_d)}{(n-l)!}, \quad \text{as } X \rightarrow \infty. \quad (9)$$

By (8) and the Lemma we obtain the formula

$$\begin{aligned} \sum_{l=0}^n b_l(d\sqrt{N}) \frac{L^{(n-l)}(1, E_d)}{(n-l)!} &\sim \\ &\sim \begin{cases} 2A_n(d\sqrt{N}, \chi_d) & \text{if } n \equiv \frac{w+1}{2} \pmod{2} \\ 0 & \text{if } n \equiv \frac{w-1}{2} \pmod{2}, \end{cases} \quad (10) \end{aligned}$$

as  $d \rightarrow \infty$ . Next we sum both sides of the above equality multiplied by the weighted function  $F\left(\frac{d}{D}\right)$  over the numbers  $d \in \mathbb{D}$ . The contribution of the right-hand side of (10) is evaluated on the basis of the results on square-free sieve obtained in § 7-§ 9 of [Iw]. Precisely we have

$$\sum_{d \in \mathbb{D}}^* 2A_n(d\sqrt{N}, \chi_d) F\left(\frac{d}{D}\right) \sim cD \int F(t) B_n(tD\sqrt{N}) dt, \quad (11)$$

where

$$B_n(X) = \frac{1}{2\pi i} \int_{(3/4)} L(s+1) \frac{\Gamma(s)}{s^n} \left(\frac{X}{2\pi}\right)^s ds,$$



the constant  $c$  is defined by (5) and the series  $L(s)$  is defined in §9 of [Iw].

In order to find the asymptotical behaviour of the right-hand side of (11) we denote

$$\tilde{G}_n(s, X) = L(s+1) \frac{\Gamma(s)}{s^n} \left(\frac{X}{2\pi}\right)^s$$

and apply the contour integration to obtain

$$\begin{aligned} B_n(X) &= \frac{1}{2\pi i} \int_{(3/4)} \tilde{G}_n(s, X) ds \\ &= \operatorname{res}_{s=0} \tilde{G}_n(s, X) + \frac{1}{2\pi i} \int_{(-1/4)} \tilde{G}_n(s, X) ds \\ &= \operatorname{res}_{s=0} \tilde{G}_n(s, X) + O(X^{-1/4}). \end{aligned}$$

Applying the asymptotical equality (9) with  $G_n$  replaced by  $\tilde{G}_n$  we obtain

$$B_n(X) \sim \sum_{l=0}^n b_l(X) \frac{L^{(n-l)}(1)}{(n-l)!} \sim b_n(X) L(1) \sim \frac{1}{n!} (\ln X)^n L(1), \quad \text{as } X \rightarrow \infty.$$

Hence the right-hand side of (11) is asymptotically equal to

$$L(1) \frac{c_0}{n!} D \int F(t) \ln^n(tD\sqrt{N}) dt \sim L(1) \frac{c_0}{n!} \int F(t) dt D(\ln D)^n.$$

Therefore in view of (10) we obtain the asymptotic equality

$$\begin{aligned} &\sum_{d \leq D} \sum_{l=0}^n b_l(d\sqrt{N}) \frac{L^{(n-l)}(1, E_d)}{(n-l)!} F\left(\frac{d}{D}\right) \sim \\ &\sim \begin{cases} \frac{c}{n!} L(1) \int F(t) dt D(\ln D)^n & \text{if } n \equiv \frac{w+1}{2} \pmod{2} \\ 0 & \text{if } n \equiv \frac{w-1}{2} \pmod{2}. \end{cases} \end{aligned}$$

Hence we obtain immediately that the constant  $c_0$  in Theorem 2 is equal to  $\left(\frac{1-w}{2}\right) c$ , where  $c$  is the constant defined by (5).

Assuming that

$$\sum_{d \in \mathbb{D}} L^{(k)}(1, E_d) F\left(\frac{d}{D}\right) \sim c_k L(1) \int F(t) dt D(\ln D)^k$$

where  $c_k$  are some constants we see that they have to satisfy the equality

$$\sum_{k=0}^n \binom{n}{k} c_k = \begin{cases} c & \text{if } n \equiv \frac{w+1}{2} \pmod{2} \\ 0 & \text{if } n \equiv \frac{w-1}{2} \pmod{2}. \end{cases} \quad (12)$$

To complete the proof of Theorem 2 it remains to prove that (12) holds with the constants  $c_k$  defined by (4). Indeed we have

$$\begin{aligned} \sum_{k=0}^n \binom{n}{k} c_k &= \left(\frac{1-w}{2}\right) c + \sum_{k=1}^n \binom{n}{k} c_k \\ &= c \left( \frac{1-w}{2} - w \left( \frac{-1}{2} \sum_{k=1}^n \binom{n}{k} (-2)^k \right) \right) \\ &= c \left( \frac{1-w}{2} - \frac{w}{2} ((-1)^n - 1) \right) \\ &= \begin{cases} c & \text{if } n \equiv \frac{w+1}{2} \pmod{2} \\ 0 & \text{if } n \equiv \frac{w-1}{2} \pmod{2} \end{cases} \end{aligned}$$

as it is claimed. This completes the proof of Theorem 2.

**Acknowledgement.** I wish to thank Professor H. Iwaniec for some very useful conversations on the subject.

#### REFERENCES

- [B-S] B. Birch and H. Swinnerton-Dyer, *Elliptic curves and modular functions*, in *Modular functions of one variable IV*, Lecture Notes in Mathematics, Springer-Verlag, vol. 476, 1975, pp. 2–32.
- [B-F-H] D. Bump, S. Friedberg and H. Hoffstein, *Non-vanishing theorems for  $L$ -functions of modular forms and their derivatives*, *Invent. Math.* **102** (1990), 543–618.
- [HB] D.R. Heath-Brown, *A mean value estimate for real character sum*, *Acta Arith.* **72** (1995), 235–275.
- [Iw] H. Iwaniec, *On the order of vanishing of modular  $L$ -functions at the critical point*, *Séminaire de Théorie des Nombres de Bordeaux* **2** (1990), 365–376.
- [M-M] M.R. Murty and V.K. Murty, *Mean values of derivatives of modular  $L$ -series*, *Ann. of Math.* **133** (1991), 447–475.

- [Mo] H.L. Montgomery, *Topics in Multiplicative Number Theory*, Lecture notes in Mathematics, Springer-Verlag, vol. 227, 1971.
- [Ko] V. A. Kolyvagin, *Finiteness of  $E(Q)$  and  $III(E(Q))$  for a subclass of Weil curves*, Math. USSR Izvest. **32** (1989), 523–542.
- [P-P] A. Perelli and J. Pomykała, *Averages over twisted elliptic  $L$ -functions*, Acta Arith. **80** (1997), 149–163.
- [P-S] J. Pomykała and J. Szmidt, *On the order of vanishing of  $n$ -th derivatives of  $L$ -functions of elliptic curves*, Biuletyn Wojskowej Akademii Technicznej **12 42/496** (1993).
- [Wa] J.-L. Waldspurger, *Sur les coefficients de Fourier des formes modulaires de poids demi-entier*, J. Math. Pures Appl. **60** (1981), 375–484.

Jacek POMYKAŁA  
Institute of Mathematics  
Warsaw University  
ul. Banacha 2  
02-097 Warsaw, Poland  
e-mail : pomykala@impan.gov.pl