

JOURNAL

de Théorie des Nombres
de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Ritabrata MUNSHI

Density of rational points on cyclic covers of \mathbb{P}^n

Tome 21, n° 2 (2009), p. 335-341.

<http://jtnb.cedram.org/item?id=JTNB_2009__21_2_335_0>

© Université Bordeaux 1, 2009, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

Density of rational points on cyclic covers of \mathbb{P}^n

par RITABRATA MUNSHI

RÉSUMÉ. Nous obtenons une majoration de la densité des points rationnels sur les revêtements cycliques de \mathbb{P}^n . Quand $n \rightarrow \infty$ notre estimation tend vers la majoration conjecturale de Serre.

ABSTRACT. We obtain upper bound for the density of rational points on the cyclic covers of \mathbb{P}^n . As $n \rightarrow \infty$ our estimate tends to the conjectural bound of Serre.

1. Introduction

Let \mathbb{P}^n denote the n -dimensional projective space over the field of rational numbers. For a finite cover $f : X \rightarrow \mathbb{P}^n$ over \mathbb{Q} , one can define a counting function

$$N(f, B) = |\{P \in X(\mathbb{Q}) : H(f(P)) \leq B\}|,$$

where H is the standard multiplicative height function defined on \mathbb{P}^n . If the degree of f is at least two, Serre [10] proved that

$$N(f, B) \ll B^{n+\frac{1}{2}}(\log B)^\gamma$$

with $\gamma < 1$. Here the implied constant depends both on f and γ . Serre's proof is based on an analogous result proved by Cohen [3] in the affine case. The proof is a beautiful application of the large sieve inequalities. Serre [10] also conjectures that one should have

$$N(f, B) \ll B^n(\log B)^c,$$

for some c . Broberg [2] almost settled Serre's conjecture for covers of the projective line and the projective plane. More precisely, he proved:

(1) If $f : X \rightarrow \mathbb{P}^1$ is a cover of degree d , then for every $\varepsilon > 0$ we have

$$N(f, B) \ll_{\varepsilon, f} B^{\frac{2}{d}+\varepsilon}.$$

(2) If $f : X \rightarrow \mathbb{P}^2$ is a cover of degree $d > 2$, then for every $\varepsilon > 0$ we have

$$N(f, B) \ll_{\varepsilon, f} B^{2+\varepsilon}.$$

In the case $d = 2$, we have

$$N(f, B) \ll_{\varepsilon, f} B^{\frac{9}{4}+\varepsilon}$$

for every $\varepsilon > 0$.

Broberg’s method is based on that introduced by Heath-Brown in [7]. These methods, though elementary, have proved to be powerful in many situations. In fact the last result of Broberg for degree two covers of \mathbb{P}^2 , is the best known result towards Manin’s conjecture, which predicts the exponent to be $1 + \varepsilon$, for del Pezzo surfaces of degree 2. Here we will give a slightly better estimate albeit at the cost of employing heavy machinery.

To this end let $F(x_0, \dots, x_n)$ be an irreducible homogenous polynomial of degree md , with $d > 1$. Moreover suppose that the projective hypersurface defined by $F(\mathbf{x}) = 0$ is smooth. Then the equation

$$y^d = F(x_0, \dots, x_n)$$

defines a variety X in the weighted projective space $\mathbb{P}(m, 1, \dots, 1)$, where y is given weight m and each x_i is given weight 1. This variety can be viewed as a cyclic d -sheeted cover of the projective space \mathbb{P}^n , via the natural map

$$\begin{aligned} f : X &\rightarrow \mathbb{P}^n; \\ (y, x_0, \dots, x_n) &\mapsto (x_0, \dots, x_n). \end{aligned}$$

This cover is ramified above the hypersurface $F(\mathbf{x}) = 0$. Using adjunction formula one easily verifies the canonical sheaf of X to be

$$\omega_X = O_X(-m + md - n - 1).$$

Hence, if the parameters n, m and d are such that

$$md - m < n + 1$$

then the variety X is a Fano variety, i.e. the anticanonical sheaf is ample. In this case we have a more precise conjecture due to Manin et al ([1], [5]), regarding the density of rational points.

In this paper, we employ the power sieve together with Poisson summation formula, and deep results of Deligne [4], and of Katz [9] about cancellation in mixed character sums over finite fields, to estimate the number of d -powers in the set

$$\{F(x_0, \dots, x_n) : -B < x_i < B\}$$

with some smooth weight W . This will give us our main result.

Theorem 1.1. *Let $f : X \rightarrow \mathbb{P}^n$ be a finite cyclic cover of \mathbb{P}^n over \mathbb{Q} . Then we have*

$$N(f, B) \ll_f B^{n + \frac{1}{n+2}} (\log B)^{\frac{n+1}{n+2}}.$$

Remark. In the special case of cyclic covers our result is better than that of Serre [10]. Also it is exciting to note that as the dimension n gets larger our bound comes closer to the conjectured bound of Serre.

Remark. For $n = 2$ and $d = 2$ our result gives

$$N(f, B) \ll B^{\frac{9}{4}} (\log B)^{\frac{3}{4}},$$

which improves the result of Broberg by an ε . However, for $n = 2, d > 2$ our result is worse.

2. Power sieve

In this section we briefly recall the concept of d -power sieve (see [6] for $d = 2$). Let p be a prime such that $p \equiv 1 \pmod{d}$. Then for every non-zero element a in the finite field \mathbb{F}_p , we can define a d -th root of unity by $a^{\frac{p-1}{d}}$. Now since \mathbb{F}_p^* is cyclic, we can define a non-canonical isomorphism

$$\theta_p : \mathbb{F}_p^* \rightarrow \mu_{p-1},$$

where μ_{p-1} denotes the group of $(p - 1)$ -th roots of unity in \mathbb{C}^* . Using this we can define a primitive Dirichlet character modulo p ,

$$\chi_p(n) = \theta_p\left(\bar{n}^{\frac{p-1}{d}}\right)$$

for $(p, n) = 1$, and $\chi_p(n) = 0$ for $p|n$. The crucial property of this character is the following:

$$\chi_p(n) = 1 \quad \text{if } (p, n) = 1, \text{ and } n = m^d \text{ for some } m.$$

Now let $\mathcal{A} = (a(n))$ be a finite sequence of non-negative quantities. We are interested in estimating the sum over the d -powers

$$S(\mathcal{A}) = \sum_n a(n^d).$$

Let \mathcal{P} be a set of P primes each having residue 1 modulo d . Suppose that $a(n) = 0$ if $|n| \geq e^P$, and consider the expression

$$S = \sum_n a(n) \left| \sum_{p \in \mathcal{P}} \chi_p(n) \right|^2$$

Each n is counted with a non-negative weight, and if $n = m^d$ and $0 < |n| < e^P$ then

$$\sum_{p \in \mathcal{P}} \chi_p(n) = \sum_{\substack{p \in \mathcal{P} \\ (p,n)=1}} 1 \gg P.$$

Hence $P^2 S(\mathcal{A}) \ll S$. Moreover

$$\begin{aligned} S &= \sum_{p,q \in \mathcal{P}} \sum_n a(n) \chi_p(n) \bar{\chi}_q(n) \\ &\leq P \sum_n a(n) + \sum_{p \neq q \in \mathcal{P}} \left| \sum_n a(n) \chi_p(n) \bar{\chi}_q(n) \right|. \end{aligned}$$

Hence we obtain the following:

Lemma 2.1. *Let \mathcal{P} be a set of P primes each having residue 1 modulo d . Suppose that $a(n) = 0$ for $n = 0$ and for $|n| \geq e^P$. Then*

$$S(\mathcal{A}) \ll P^{-1} \sum_n a(n) + P^{-2} \sum_{p \neq q \in \mathcal{P}} \left| \sum_n a(n) \chi_p(n) \bar{\chi}_q(n) \right|.$$

3. Proof of Theorem 1

Let $W : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ be a non-negative smooth function supported in the dyadic box $[-B, B]^{n+1}$, and such that the partial derivatives satisfy the following bound:

$$\left| \frac{d^{i_0 + \dots + i_n} W(x_0, \dots, x_n)}{dx_0^{i_0} \dots dx_n^{i_n}} \right| \ll B^{-(i_0 + \dots + i_n)}.$$

Then via integration-by-parts we note the following bound for the Fourier transform

$$\hat{W}(\mathbf{u}) = \int_{\mathbb{R}^{n+1}} W(\mathbf{x}) e(-\langle \mathbf{x}, \mathbf{u} \rangle) d\mathbf{x} \ll B^{n+1} \prod_{i=0}^n (1 + |u_i|B)^{-2}$$

where $\mathbf{u} = (u_0, \dots, u_n)$.

Then we define a sequence of non-negative numbers by

$$a(k) = \sum_{\mathbf{x} \in \mathbb{Z}^{n+1}, F(\mathbf{x})=k} W(\mathbf{x}).$$

We wish to obtain an upper bound for

$$S(\mathcal{A}) = \sum_k a(k^d)$$

using the d -power sieve. Now clearly we have

$$\sum_k a(k) \ll B^{n+1}.$$

Hence to apply the lemma we need to estimate the sum

$$S(p, q) = \sum_k a(k) \chi_p(k) \bar{\chi}_q(k) = \sum_{\mathbf{x} \in \mathbb{Z}^{n+1}} W(\mathbf{x}) \chi_p(F(\mathbf{x})) \bar{\chi}_q(F(\mathbf{x})),$$

where p and q are two distinct primes. Using Poisson summation formula we obtain

$$\begin{aligned} S(p, q) &= \sum_{\mathbf{a} \pmod{pq}} \chi_p(F(\mathbf{a})) \bar{\chi}_q(F(\mathbf{a})) \sum_{\mathbf{x} \equiv \mathbf{a}} W(\mathbf{x}) \\ &= (pq)^{-(n+1)} \sum_{\mathbf{u} \in \mathbb{Z}^{n+1}} g(\mathbf{u}; p, q) \hat{W}\left(\frac{\mathbf{u}}{pq}\right), \end{aligned}$$

where

$$g(\mathbf{u}; p, q) = \sum_{\mathbf{a} \pmod{pq}} \chi_p(F(\mathbf{a})) \bar{\chi}_q(F(\mathbf{a})) e\left(\frac{\langle \mathbf{a}, \mathbf{u} \rangle}{pq}\right)$$

is a mixed character sum. Then we observe that the above sum satisfies the following multiplicative property

$$g(\mathbf{u}; p, q) = g(\bar{q}\mathbf{u}; \chi_p) g(\bar{p}\mathbf{u}; \bar{\chi}_q),$$

where $p\bar{p} \equiv 1 \pmod{q}$, $q\bar{q} \equiv 1 \pmod{p}$ and

$$g(\mathbf{u}; \chi_p) = \sum_{\mathbf{a} \pmod{p}} \chi_p(F(\mathbf{a})) e\left(\frac{\langle \mathbf{a}, \mathbf{u} \rangle}{p}\right).$$

So our job is reduced to estimating the mixed character sum over the finite field \mathbb{F}_p . Let V be the hypersurface defined by $F(\mathbf{x}) = 0$ over the field of rationals. Let V^* be its dual variety, which in this case is again a hypersurface. Assuming that the prime p is such that the reduction $V_p = V \pmod{p}$ is smooth, we have three situations:

(1) First suppose that the vector $\mathbf{u} \in \mathbb{Z}^{n+1}$ is such that $\mathbf{u} \equiv \mathbf{0} \pmod{p}$, then

$$g(\mathbf{u}; \chi_p) = \sum_{\mathbf{a} \pmod{p}} \chi_p(F(\mathbf{a})) \ll p^{\frac{n+1}{2}},$$

where the implied constant depends only on n and the degree of F . This follows from the multiplicative analogue of Deligne’s bound established by Katz [8].

(2) If $\mathbf{u} \pmod{p}$ is non-zero and the associated hyperplane $\langle \mathbf{u}, \mathbf{x} \rangle = 0$ is not a tangent to the hypersurface V_p , then by Theorem 1.1 in [9] we have square-root cancellation i.e.

$$g(\mathbf{u}; \chi_p) \ll p^{\frac{n+1}{2}}.$$

Here again the implied constant depends only on n and the degree of F . In particular, it does not depend on \mathbf{u} .

(3) If $\mathbf{u} \pmod{p}$ is non-zero and the associated hyperplane $\langle \mathbf{u}, \mathbf{x} \rangle = 0$ is a tangent to the hypersurface V_p , then using the primitivity of the character we get

$$g(\mathbf{u}; \chi_p) = \frac{\chi_p(-1)\tau(\chi_p)}{p} \sum_{\mathbf{b} \pmod{p}} \bar{\chi}_p(\mathbf{b}) \sum_{\mathbf{a} \pmod{p}} e\left(\frac{\mathbf{b}F(\mathbf{a}) + \langle \mathbf{a}, \mathbf{u} \rangle}{p}\right),$$

where $\tau(\chi)$ stands for the Gauss sum of the character χ . Then we apply Deligne’s bound [4] to the inner sum to obtain

$$g(\mathbf{u}; \chi_p) \ll \sqrt{pp}^{\frac{n+1}{2}}.$$

Again the implied constant depends only on n and the degree of F .

In the first two cases we say that the vector $\mathbf{u} \in \mathbb{Z}^{n+1}$ is ‘good’ modulo p . In the third case we say that \mathbf{u} is ‘bad’ modulo p .

We first compute the contribution to $S(p, q)$ from those vectors $\mathbf{u} \in \mathbb{Z}^{n+1}$ which are ‘good’ modulo both p and q . Using square-root cancellation we see that this is bounded by

$$\ll B^{n+1} \frac{(pq)^{\frac{n+1}{2}}}{(pq)^{n+1}} \sum_{\mathbf{u} \in \mathbb{Z}^{n+1}} \prod_{i=0}^n \left(1 + \frac{|u_i|B}{pq}\right)^{-2} \ll (pq)^{\frac{n+1}{2}}.$$

In the last inequality we are assuming that $pq \geq B$, which follows from our choice of the set \mathcal{P} . Now the contribution of those vectors $\mathbf{u} \in \mathbb{Z}^{n+1}$ which are ‘bad’ for the prime p but ‘good’ for the prime q is given by

$$(pq)^{-(n+1)} \sum_{\mathbf{v} \in V_p^*(\mathbb{F}_p)} \sum_{\lambda=1}^p \sum_{\mathbf{u} \equiv \lambda \mathbf{v} \pmod{p}}^* g(\mathbf{u}; p, q) \hat{W}\left(\frac{\mathbf{u}}{pq}\right).$$

The $*$ above the sum indicates that the sum is restricted over those vectors $\mathbf{u} \in \mathbb{Z}^{n+1}$ which are ‘good’ modulo q . The above expression is bounded by

$$\begin{aligned} & B^{n+1} \frac{p^{\frac{n+2}{2}} q^{\frac{n+1}{2}}}{(pq)^{n+1}} \sum_{\mathbf{v} \in V_p^*(\mathbb{F}_p)} \sum_{\lambda=1}^p \sum_{\mathbf{u} \equiv \lambda \mathbf{v} \pmod{p}}^* \prod_{i=0}^n \left(1 + \frac{|u_i|B}{pq}\right)^{-2} \\ & \ll B^{n+1} \frac{p^{\frac{n+2}{2}} q^{\frac{n+1}{2}}}{(pq)^{n+1}} \sum_{\mathbf{v} \in V_p^*(\mathbb{F}_p)} \sum_{\lambda=1}^p \left(\frac{q}{B}\right)^{n+1} \ll p^{\frac{n}{2}} q^{\frac{n+1}{2}}. \end{aligned}$$

In the last inequality we have used the fact that $|V_p^*(\mathbb{F}_p)| \ll p^{n-1}$. Similarly one can show that the contribution of those vectors $\mathbf{u} \in \mathbb{Z}^{n+1}$ which are ‘bad’ for both p and q , is bounded by $(pq)^{\frac{n}{2}}$. Hence we conclude that

$$S(p, q) \ll (pq)^{\frac{n+1}{2}}.$$

Now for the set \mathcal{P} we pick P consecutive primes in the progression $(1 + dk)_k$ of size $\sim P \log P$. Then using the d -power sieve inequality, we get

$$S(\mathcal{A}) \ll \frac{B^{n+1}}{P} + (P \log P)^{n+1}.$$

Then choosing

$$P = \left(\frac{B}{\log B}\right)^{\frac{n+1}{n+2}}$$

the theorem follows.

Acknowledgements

I thank the organizers of the 25th Journées Arithmétiques for inviting me to this exciting conference, and for the support.

References

- [1] V.V. BATYREV; Y.I. MANIN, *Sur le nombre des points rationnels de hauteur borné des variétés algébriques*. Math. Ann. **286** (1990), 27–43.
- [2] N. BROBERG, *Rational points on finite covers of \mathbb{P}^1 and \mathbb{P}^2* . J. Number Theory **101** (2003), 195–207.
- [3] S.D. COHEN, *The distribution of Galois groups and Hilbert’s irreducibility theorem*. Proc. London Math. Soc. (3) **43** (1981), 227–250.
- [4] P. DELIGNE, *La conjecture de Weil. I*. Inst. Hautes Études Sci. Publ. Math. No. **43** (1974), 273–307.
- [5] J. FRANKE; Y.I. MANIN; Y. TSCHINKEL, *Rational points of bounded height on Fano varieties*. Invent. Math. **95** (1989), 421–435.
- [6] D.R. HEATH-BROWN, *The square sieve and consecutive square-free numbers*. Math. Ann. **266** (1984), 251–259.
- [7] D.R. HEATH-BROWN, *The density of rational points on curves and surfaces*. Ann. of Math. (2) **155** (2002), 553–595.
- [8] N. KATZ, *Estimates for nonsingular multiplicative character sums*. Int. Math. Res. Not. (2002), 333–349.
- [9] N. KATZ, *Estimates for nonsingular mixed character sums*. Int. Math. Res. Not. (2007), vol. 2007, article ID rnm069, 19 pages, doi:10.1093/imrn/rnm069.
- [10] J-P. SERRE, *Lectures on the Mordell-Weil theorem*. Friedr. Vieweg & Sohn, Braunschweig (1989).

Ritabrata MUNSHI
Rutgers University
110, Frelinghuysen Road
Piscataway NJ 08854, USA
E-mail: rmunshi@math.rutgers.edu
URL: <http://www.math.rutgers.edu/~rmunshi>