

A. BENDALI

**Approximation of a degenerated elliptic boundary value problem by a finite element method**

*RAIRO. Analyse numérique*, tome 15, n° 2 (1981), p. 87-99

[http://www.numdam.org/item?id=M2AN\\_1981\\_\\_15\\_2\\_87\\_0](http://www.numdam.org/item?id=M2AN_1981__15_2_87_0)

© AFCET, 1981, tous droits réservés.

L'accès aux archives de la revue « RAIRO. Analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## APPROXIMATION OF A DEGENERATED ELLIPTIC BOUNDARY VALUE PROBLEM BY A FINITE ELEMENT METHOD (\*) (\*\*)

par A. BENDALI (<sup>1</sup>)

Communique par P G CIARLET

Résumé — On considère un problème aux limites elliptique dégénéré, provenant d'un problème de Dirichlet ayant une symétrie de révolution. Pour approcher la solution, nous construisons une méthode d'éléments fins basée sur une triangulation usuelle du domaine, mais avec des fonctions approchantes adaptées. Cet aspect a déjà été mis en relief, en dimension un, par Crouzeix et Thomas. Par une technique d'approximation de domaine et d'estimation a priori, nous donnons une démonstration directe, dans le cadre bidimensionnel, de la régularité de la solution. Ceci permet de retrouver un comportement asymptotique de l'erreur, identique au cas standard.

Abstract — We consider a degenerated elliptic boundary value problem arising in the (concrete) axisymmetric Dirichlet problem. We construct a finite element approximation of the solution using a classical triangulation, but with special approximating functions. This aspect has already been pointed out in the one-dimensional case by Crouzeix and Thomas. Here we give a direct proof of the regularity of the solution in the two-dimensional case, using a technique of approximated domain and a priori estimates. This leads to an asymptotic error behaviour which is the same as in the usual case.

### 1. STATEMENT OF THE PROBLEM

Let  $\Omega$  be a convex bounded domain with a polygonal boundary contained in  $\mathbb{R}_+^2 = \{ (r, z) \in \mathbb{R}^2 ; r > 0 \}$ ;  $\Gamma = \partial\Omega \cap \mathbb{R}_+^2$  is the boundary of  $\Omega$  in  $\mathbb{R}_+^2$ . We assume that it intersects the  $z$ -axis at right angles, i.e. there exists  $R > 0$  such that

$$\Omega_R = \{ (r, z) \in \Omega ; 0 < r < R \} = ]0, R[ \times ]0, H[ .$$

(\*) Reçu en décembre 1979

(\*\*) Ce travail a été financé par un contrat avec l'O N R S (Algérie)

(<sup>1</sup>) Centre de Mathématiques Appliquées, Ecole Polytechnique, France




Figure 1.

Let  $f$  be a given function defined in  $\Omega$ ; we consider the following boundary problem :

$$\left. \begin{aligned} -\Delta_r u &= -\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial u}{\partial r} \right) - \frac{\partial^2 u}{\partial z^2} = f \quad \text{in } \Omega; \\ u &= 0 \quad \text{on } \Gamma. \end{aligned} \right\} \quad (1.1)$$

This problem has a (weak) variational formulation which needs the introduction of some functions spaces in order to be stated. The notations are those used in the partial differential equations theory (see e.g. Lions-Magenes [8]).

Let us denote by  $L_1^2(\Omega)$  the weighted  $L^2$  space of all Lebesgue measurable functions  $u$  such that

$$\int_{\Omega} |u|^2 r \, dr \, dz \quad \text{is finite}$$

and by

$$H_1^m(\Omega) = \{ u \in L_1^2(\Omega); D^\alpha u \in L_1^2(\Omega); \forall \alpha; |\alpha| \leq m \},$$

the related Sobolev space of order  $m$ . All these spaces, equipped with their natural scalar products and norms, are Hilbert spaces. If  $A$  is a measurable subset of  $\Omega$ , we shall use the semi-norm of  $H_1^m(\Omega)$  defined for  $v$  in  $H_1^m(\Omega)$  by

$$|v|_{j,A} = \left\{ \sum_{|\alpha|=j} \int_A |D^\alpha v|^2 r \, dr \, dz \right\}^{1/2} \quad \text{for } j = 0, 1, \dots, m.$$

Then, if  $V$  is the closure of  $\mathcal{D}(\Omega)$  in the  $H_1^1(\Omega)$  norm, we can give a precise statement of (1.1).

Let  $f$  in  $\mathcal{D}'(\Omega)$  be such that  $rf$  ranges in  $V'$ ; the problem is

$$\left. \begin{aligned} &\text{Find } u \text{ in } V; \forall v \in V; \\ &a(u, v) = \langle rf, v \rangle, \end{aligned} \right\} \tag{1.2}$$

where

$$a(u, v) = \int_{\Omega} (\text{grad } u) \cdot (\text{grad } v) r \, dr \, dz, \tag{1.3}$$

$\langle \cdot, \cdot \rangle$  is the duality pairing between  $V'$  and  $V$ .

It is easily seen by a Poincaré's inequality (we shall later see that in a particular and fundamental aspect) that the continuous bilinear form given by (1.3) is  $V$ -elliptic. Therefore, the Lax-Milgram theorem insures the existence and uniqueness of the solution of problem (1.2).

*Remark*

1) Problem (1.2) solves (1.1) in the following sense :

$$\left\{ \begin{aligned} &-\Delta_r u = f \quad \text{in } \mathcal{D}'(\Omega), \\ &u \in V. \end{aligned} \right.$$

2) If  $v$  belongs to  $H_1^1(\Omega)$ , the trace of  $u$  on  $\Gamma$  is defined as an element of  $H_{loc}^{1/2}(\Gamma)$ .  $u$  is in  $V$  iff  $u = 0$  (in the sense of  $H_{loc}^{1/2}(\Gamma)$ ); for all this, see Amirat [1], Atik [2], Lailly [7].

Now, with a variational formulation, we can make various conformal approximations of the space  $V$  in order to approximate the solution of (1.2). For the one considered here, we have an error behaviour which is the same as in the standard case.

We consider the following mesh of  $\Omega$  denoted by  $\mathcal{T}^h$ ;  $h$  is the maximum of diameters of the elements of  $\mathcal{T}^h$  and is destined to tend to 0.




Figure 2.

$K$  will be a generic element of  $\mathcal{T}^h$ ,  $\mathcal{N}$  is the set of all vertices of the elements  $K$  which do not lie on the  $z$ -axis.

We consider three kinds of approximating functions.

*Case 1* :  $K$  is a rectangle with a side on the  $z$ -axis ; we have

$$P_K = \{ \alpha + \beta z ; \alpha, \beta \in \mathbb{R} \} ;$$

$\Sigma_K$  represents the vertices of  $K$  which are not on the  $z$ -axis.

*Case 2* :  $K$  is a rectangle and has not the above property ; then

$$P_K = \{ \alpha + \beta z + \gamma \log r + \delta r \log r ; \alpha, \beta, \gamma, \delta \in \mathbb{R} \} ;$$

$\Sigma_K$  represents the set of the vertices of  $K$ .

*Case 3* :  $K$  is a triangle (which happens far from the  $z$ -axis) ; we have

$$P_K = \mathbb{P}_1 = \{ \alpha + \beta z + \gamma r ; \alpha, \beta, \gamma \in \mathbb{R} \} ;$$

$\Sigma_K$  represents the vertices of  $K$ .

In each case, if  $v$  is a continuous function around the elements of  $\Sigma_K$ , we can define in a unique way the interpolate  $\pi v$  of  $v$  by

$$\begin{cases} \pi v \in P_K, \\ \pi v(a) = v(a) ; \forall a \in \Sigma_K. \end{cases}$$

Then, we define

$$S^h(\Omega) = \{ v^h \in C^0(\overline{\Omega}) ; v^h|_K \in P_K ; \forall K \in \mathcal{T}^h \} .$$

It is easily seen that we have the following result :

**PROPOSITION 1.1** :  $S^h(\Omega)$  is a subspace of  $H_1^1(\Omega)$ . Every  $v^h$  in  $S^h(\Omega)$  is biuniquely determined by the collection of nodal parameters  $v^h(a)$ ,  $a \in \mathcal{N}$ .

If  $V^h$  denotes the subspace formed by  $v^h$  in  $S^h(\Omega)$  such that  $v^h(a) = 0$  when  $a$  lies in  $\Gamma \cap \mathcal{N}$ , then  $V^h$  is a subspace of  $V$ . ■

Now, we state the discrete problem :

$$\text{Find } u^h \in V^h ; \forall v^h \in V^h ; \text{ such that } \left. \begin{aligned} a(v^h, v^h) &= \langle rf, v^h \rangle. \end{aligned} \right\} \quad (1.4)$$

Cea's lemma leads to

$$\| u - u^h \|_{H_1^1(\Omega)} \leq c \| u - v^h \|_{H_1^1(\Omega)} ; \forall v^h \in V^h, \quad (1.5)$$

where  $c$  is a constant independent of  $u$  and  $h$ .

In the sequel,  $c$  will denote various pure constants and we shall indicate which elements it does not depend on, in each case.

To give the error estimate, we need some regularity results.

**2. REGULARITY OF THE SOLUTION**

We suppose that  $f$  is in  $L^2_1(\Omega)$  (i.e. the data in the axisymmetric three-dimensional problem is  $L^2$ ). We have the following main result : we set

$$D^2_1(\Omega) = \left\{ u \in H^2_1(\Omega); \frac{1}{r} \frac{\partial u}{\partial r} \in L^2_1(\Omega) \right\},$$

$$E(\Omega) = D^2_1(\Omega) \cap V;$$

then, we have

THEOREM 2.1 : *The operator*

$$-\Delta_r : E(\Omega) \rightarrow L^2_1(\Omega)$$

is an isomorphism of Banach spaces. ■

*Remark* : Since  $f$  is in  $L^2_1(\Omega)$ , we have  $\langle rf, v \rangle = (f, v)$  where  $(f, v)$  denotes the scalar product of  $f$  and  $v$  in  $L^2_1(\Omega)$ , for  $v$  in  $V$ .

The proof of theorem 2.1 will be reduced to the non degenerated case by the following à priori estimates.

For  $\varepsilon > 0$  small enough,  $\Omega_\varepsilon$  is the approximated (convex) domain given by cutting in  $\Omega$  a band of width  $\varepsilon$  around the  $z$ -axis :

$$\Omega_\varepsilon = \{ (r, z) \in \Omega; r > \varepsilon \}$$

$$\Gamma_\varepsilon^D = \Gamma \cap \partial\Omega_\varepsilon$$

$$\Gamma_\varepsilon^N = \partial\Omega_\varepsilon \setminus \Gamma_\varepsilon^D$$

$$E_\varepsilon = \left\{ v \in H^2(\Omega_\varepsilon); \frac{\partial v}{\partial r} = 0 \text{ on } \Gamma_\varepsilon^N; v = 0 \text{ on } \Gamma_\varepsilon^D \right\}.$$

For suitable  $v$ , we shall use the following notation :

$$|v|_{j, \Omega_\varepsilon} = \left\{ \sum_{|\alpha|=j} \int_{\Omega_\varepsilon} |D^\alpha v|^2 r \, dr \, dz \right\}^{1/2}$$

LEMMA 2.1 : *There exists a constant  $c$  independent of  $\varepsilon > 0$  and of  $v$  in  $E_\varepsilon$ , such that*

$$|v|_{0, \Omega_\varepsilon}^2 + |v|_{1, \Omega_\varepsilon}^2 \leq c^2 |\Delta_r v|_{0, \Omega_\varepsilon}^2. \quad \blacksquare \tag{2.1}$$




Figure 3.

*Proof* : Since  $v$  is in  $E_\epsilon$  and using Green's formula, we obtain

$$-\int_{\Omega_\epsilon} (\Delta_r v) v r dr dz = -\int_{\Omega_\epsilon} v \operatorname{div} (r \operatorname{grad} v) dr dz = |v|_{1, \Omega_\epsilon}^2 .$$

Let  $\Phi : ]0, H[ \rightarrow \mathbb{R}$  be the continuous piecewise affine function describing  $\Omega$  by

$$\Omega = \{ (r, z) \in \mathbb{R}^2 ; 0 < z < H ; 0 < r < \Phi(z) \} ; \tag{2.2}$$

we have

$$|v(r, z)| \leq \int_r^{\Phi(z)} \left| \frac{\partial v}{\partial r}(s, z) \right| ds ,$$

$$\sqrt{r} |v(r, z)| \leq \int_r^{\Phi(z)} \sqrt{s} \left| \frac{\partial v}{\partial r}(s, z) \right| ds \leq c \left\{ \int_r^{\Phi(z)} \left| \frac{\partial v}{\partial r} \right|^2 s ds \right\}^{1/2} .$$

Here  $c$  can only be the square root of the diameter of  $\Omega$  and does not depend on  $\epsilon$  and  $v$  (see the above remark on the coerciveness of the bilinear form of problem (1.2)). Integrating in  $z$  gives (2.1) (with another constant). ■

LEMMA 2.2 : For all  $v \in E_\epsilon$ , we have

$$\left| \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial v}{\partial r} \right) \right|_{0, \Omega_\epsilon}^2 + \left| \frac{\partial^2 u}{\partial r \partial z} \right|_{0, \Omega_\epsilon}^2 + \left| \frac{\partial^2 u}{\partial z^2} \right|_{0, \Omega_\epsilon}^2 \leq |\Delta_r u|_{0, \Omega_\epsilon}^2 . \quad \blacksquare \tag{2.3}$$

*Proof* : Note that it is sufficient to prove (2.2) for  $v$  in  $H^3(\Omega_\epsilon) \cap E_\epsilon$ .

By a density argument (see Grisvard [6]), the general case is true. Then, if  $v$  is in  $H^3(\Omega_\varepsilon) \cap E_\varepsilon$ , we have

$$|\Delta_r v|_{0,\Omega_\varepsilon}^2 = \left| \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial v}{\partial r} \right) \right|_{0,\Omega_\varepsilon}^2 + \left| \frac{\partial^2 v}{\partial z^2} \right|_{0,\Omega_\varepsilon}^2 + 2 \int_{\Omega_\varepsilon} \frac{\partial}{\partial r} (rv_1) \frac{\partial v_2}{\partial z} dr dz,$$

where

$$v_1 = \frac{\partial v}{\partial r}; \quad v_2 = \frac{\partial v}{\partial z}. \tag{2.4}$$

Let us consider the one differential form  $\bar{\omega} = rv_1 dv_2$ ; by differentiating, we obtain

$$d\bar{\omega} = \frac{\partial}{\partial r} (rv_1) \frac{\partial v_2}{\partial z} dr \wedge dz - \frac{\partial}{\partial z} (rv_1) \frac{\partial v_2}{\partial r} dr \wedge dz.$$

Using Stokes' formula, we can write

$$\int_{\Omega_\varepsilon} \frac{\partial}{\partial r} (rv_1) \frac{\partial v_2}{\partial z} dr dz = \int_{\partial\Omega_\varepsilon} \bar{\omega} + \int_{\Omega_\varepsilon} \frac{\partial v_1}{\partial z} \frac{\partial v_2}{\partial r} r dr dz.$$

By (2.4) and the boundary value of  $v$  in  $E_\varepsilon$ , the curvilinear integral is reduced to the integration of  $\bar{\omega}$  along the oriented piecewise regular path given by

$$\gamma : ]O, H[ \rightarrow \mathbb{R}^2; \quad \gamma(z) = (\Phi(z), z).$$

There is a subdivision of  $]O, H[$ ,  $\alpha_0 = 0 < \alpha_1 < \dots < \alpha_{p+1} = H$ , such that

$$\Phi_i = \Phi|_{[\alpha_i, \alpha_{i+1}]} \text{ is linear affine.}$$

If  $\gamma_i$  is the oriented path associated with  $\Phi_i$ , we have

$$\int_{\partial\Omega_\varepsilon} \bar{\omega} = \sum_{i=0}^p \int_{\gamma_i} \bar{\omega}.$$

But we have  $v = 0$  on  $\gamma_i$ ; hence

$$v_1 \Phi'_i + v_2 = 0.$$

Since  $\Phi_i$  is linear affine, its derivative  $\Phi'_i$  is constant and we obtain

$$\int_{\gamma_i} \bar{\omega} = -\Phi'_i \int_{\gamma_i} rv_1 dv_1 = \frac{1}{2} \Phi'_i \int_{\gamma_i} v_1^2 dr.$$



using the fact that  $v_1$  vanishes at the endpoints of  $\gamma_i$ , since the gradient of  $v$  vanishes in two noncolinear directions. Finally, noticing that  $dr = \Phi'_i dz$  on  $\gamma_i$ , we obtain in all cases :

$$\int_{\gamma_i} \bar{\omega} = \frac{1}{2} \int_{\gamma_i} v_2^2 dz ,$$

and then

$$\int_{\partial\Omega_\varepsilon} \bar{\omega} = \frac{1}{2} \int_0^H v_2^2(\Phi(z), z) dz \geq 0 .$$

This inequality gives (2.3). ■

*Remark :* The proof is valid in all the cases where  $\Omega$  is represented by (2.2). Geometrically this means that the axisymmetric domain of  $\mathbb{R}^3$ , obtained by the rotation of  $\Omega$  around the  $z$ -axis, is convex. The three-dimensional regularity is then completely recovered.

LEMMA 2.4 : For all  $v$  in  $E_\varepsilon$ , we have

$$\left| \frac{1}{r} \frac{\partial v}{\partial r} \right|_{0, \Omega_\varepsilon}^2 + \left| \frac{\partial^2 v}{\partial r^2} \right|_{0, \Omega_\varepsilon}^2 \leq c^2 \left| \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial v}{\partial r} \right) \right|_{0, \Omega_\varepsilon}^2 \tag{2.5}$$

where  $c$  is a constant independent of  $v$  and  $\varepsilon$ . ■

*Proof :* We set

$$w = r \frac{\partial v}{\partial r}; \quad w = 0 \quad \text{on} \quad \Gamma_\varepsilon^N;$$

extending  $w$  by 0 in  $\Omega$ , we can assume that  $w$  is in  $H^1(\Omega)$ , and write

$$w(r, z) = \int_0^r \frac{\partial w}{\partial r}(s, z) ds .$$

An easy majoration gives

$$r^{-3/2} |w(r, z)| \leq \frac{1}{r} \int_0^r \frac{1}{\sqrt{s}} \left| \frac{\partial w}{\partial r} \right| ds .$$

Denote by  $\tilde{w}$  et  $\partial\tilde{w}/\partial r$  the extension by 0 of  $w$  and  $\partial w/\partial r$  respectively ; Hardy's inequality leads to

$$\int_0^{+\infty} r^{-3} |\tilde{w}(r, z)|^2 dr \leq \int_0^{+\infty} \frac{1}{r} \left| \frac{\partial\tilde{w}}{\partial r} \right|^2 dr .$$

After integrating in  $z$  and taking into account that  $\tilde{w}$  and  $\partial\tilde{w}/\partial r$  vanish outside of  $\Omega_\varepsilon$ , we obtain (2.5). ■

*Proof of theorem 2.1 :*  $-\Delta_r$  is a continuous injective operator from  $E(\Omega)$  into  $L_1^2(\Omega)$ . Theorem 2.1 will be proved if we show  $-\Delta_r$  is surjective as well. At the same time, we will give an estimate of the continuity constant of the inverse operator.

Let  $f$  be in  $L_1^2(\Omega)$ . For  $\varepsilon > 0$  small enough, we denote by  $f_\varepsilon = f|_{\Omega_\varepsilon}$  the restriction of  $f$  to  $\Omega_\varepsilon$ .

We know (see Grisvard [6] ; at this step, the convexity of  $\Omega$  is needed)  $-\Delta$  is an isomorphism of Banach spaces from  $E_\varepsilon$  onto  $L^2(\Omega_\varepsilon)$ .

$-\Delta_r = -\Delta - \frac{1}{r} \frac{\partial}{\partial r}$  appears as a perturbation of an isomorphism by a compact operator  $-\frac{1}{r} \frac{\partial}{\partial r}$ ; hence, it is a Fredholm operator of index 0. Since the kernel of  $-\Delta_r$  is reduced to 0,  $-\Delta_r$  is also onto. Then, the problem

$$\begin{cases} -\Delta_r u_\varepsilon = f_\varepsilon & \text{in } \mathcal{D}'(\Omega_\varepsilon), \\ u_\varepsilon \in E_\varepsilon, \end{cases}$$

has one (and only one) solution. Using the estimates given by the above lemmas, we have

$$\sum_{j=0}^2 |u_\varepsilon|_{j,\Omega_\varepsilon}^2 + \left| \frac{1}{r} \frac{\partial u_\varepsilon}{\partial r} \right|_{0,\Omega_\varepsilon}^2 \leq c^2 |f_\varepsilon|_{0,\Omega_\varepsilon}^2 \leq c^2 \|f\|_{L_1^2(\Omega)}^2, \quad (2.6)$$

where  $c$  is a constant independent of  $f$  and  $\varepsilon > 0$ .

Extending by 0 all the partial derivatives of  $u_\varepsilon$  to  $\Omega$ , and using the well-known properties of weak compactness of finite balls in Hilbert spaces, as well as the continuity of partial derivatives in  $\mathcal{D}'(\Omega)$ , we obtain that there exists a function  $u$  in  $D_1^2(\Omega)$  which satisfies  $-\Delta_r u = f$  in  $\mathcal{D}'(\Omega)$ .

The compactness of the injection of  $H_{loc}^2(\Omega)$  into  $H_{loc}^1(\Omega)$  implies also that  $u$  is in  $V$  (hence, it is in  $E(\Omega)$ ). Finally, the inferior semi-continuity of the norm of  $L_1^2(\Omega)$  insures that

$$\|u\|_{D_1^2(\Omega)} \leq c \|f\|_{L_1^2(\Omega)},$$

where  $c$  is a constant depending only on the diameter of  $\Omega$  by Poincaré's inequality. The constant of Hardy's inequality is independent of  $\Omega$ . ■

### 3. ERROR ESTIMATE

The solution  $u$  of problem (1.2) belongs to  $D_1^2(\Omega)$  if  $f$  is in  $L_1^2(\Omega)$ ; hence, it is continuous on  $\Omega \cup \Gamma$ . This enables us to define the interpolate of  $u$  by

$$\left. \begin{aligned} \pi^h u &\in V^h, \\ \pi^h(a) &= u(a); \quad \forall a \in \mathcal{N} \end{aligned} \right\}. \tag{3.1}$$

Thus, to estimate  $\| u - u^h \|_{H_1^1(\Omega)}$ , we need only approximating the interpolate error  $\| u - \pi^h u \|_{H_1^1(\Omega)}$ .

Let us denote by  $h_K$  and  $\rho_K$  the diameter of  $K \in \mathcal{T}^h$  and the maximum of radius of the balls inscribed in  $K$ , respectively. Assume the regularity hypothesis on the mesh  $\mathcal{T}^h$  :

$$\max_{K \in \mathcal{T}^h} \frac{h_K}{\rho_K} \leq c, \tag{3.2}$$

where  $c$  is a constant independent of  $h$  and  $h = \max_{K \in \mathcal{T}^h} h_K$ . Then we have :

**THEOREM 3.1 :** *Under the condition (3.2), if  $u$  is the solution of problem (1.2) with the data  $f$  in  $L_1^2(\Omega)$  and if  $u^h$  denotes the solution of the discrete problem (1.4), we have*

$$\| u - u^h \|_{H_1^1(\Omega)} \leq ch \| f \|_{L_1^2(\Omega)}$$

where  $c$  is a constant independent of  $h$  and  $f$  in  $L_1^2(\Omega)$ . ■

Standard techniques give the proof, using the following lemmas.

**LEMMA 3.2 :** *Let  $K = ]0, h[ \times ]0, h[$ . There exists a constant  $c$  independent of  $v$  in  $D_1^2(K)$  and of  $h$  such that*

$$| v - \pi v |_{j,K} \leq ch^{2-j} \left( | v |_{2,K}^2 + \left| \frac{1}{r} \frac{\partial v}{\partial r} \right|_{0,K}^2 \right)^{1/2}; \quad j = 0, 1. \tag{3.3}$$

*Proof :* We set  $\hat{K} = ]0, 1[ \times ]0, 1[$ . With a linear change of each variable, the proof is a consequence of

There exists a constant  $c$  independent of  $\hat{v} \in D_1^2(\hat{K})$  such that

$$\| \hat{v} \|_{D_1^2(\hat{K})}^2 \left( \leq c^2 | \hat{v}(\hat{a}_1) |^2 + | \hat{v}(\hat{a}_2) |^2 + | \hat{v} |_{2,K}^2 + \left| \frac{1}{r} \frac{\partial \hat{v}}{\partial r} \right|_{0,\hat{K}}^2 \right), \tag{3.4}$$

where  $\hat{a}_1 = (1, 0)$ ,  $\hat{a}_2 = (1, 1)$  are the vertices of  $\hat{K}$  out of the  $z$ -axis. To prove (3.4), we assume the contrary. Then, there exists a sequence  $\{ \hat{v}_n \}$  such that

$$\| \hat{v}_n \|_{D_1^2(\hat{K})} = 1; \tag{3.5}$$

$$\lim | \hat{v}_n(\hat{a}_i) | = 0; \quad i = 1, 2; \tag{3.6}$$

$$\lim | \hat{v}_n |_{2,\hat{K}} = 0; \tag{3.7}$$

$$\lim \left| \frac{1}{r} \frac{\partial \hat{v}_n}{\partial r} \right|_{0, \bar{K}} = 0. \tag{3.8}$$

Using a subsequence, we can assume by (3.5) that there exists a function  $\hat{v}$  in  $D_1^2(\hat{K})$  such that

$$\lim \hat{v}_n = \hat{v} \text{ weakly in } D_1^2(K).$$

Using (3.7), we obtain

$$\hat{v} = \alpha r + \beta z + \gamma, \quad \alpha, \beta, \gamma \in \mathbb{R}.$$

(3.8) gives  $\alpha = 0$  and (3.6),  $\beta = \gamma = 0$ . Therefore, the sequence  $\{ \hat{v}_n \}$  converges weakly to 0 in  $D_1^2(\hat{K})$ . The compactness of the unit ball of  $H_1^2(\hat{K})$  in  $H_1^1(\hat{K})$  (see e.g. El Kolli [5]) gives

$$\lim \hat{v}_n = 0 \text{ in } H_1^1(\hat{K}) \text{ strongly.}$$

But this result, together with (3.7) and (3.8) contradicts (3.5). Thus, (3.4) and the lemma are proved. ■

LEMMA 3.3 : Let  $K = ]\bar{r}, \bar{r} + h[ \times ]0, h[$

$$h \leq c\bar{r}, \tag{3.9}$$

where  $c$  is independent of  $h$  and  $\bar{r}$ . Then, there exists a constant  $c$  independent of  $h$  and  $\bar{r}$  such that

$$|v - \pi v|_{j,K} \leq ch^{2-j} \left( |v|_{2,K}^2 + \left| \frac{1}{r} \frac{\partial v}{\partial r} \right|_{0,K}^2 \right)^{1/2}; \quad j = 0, 1,$$

for all  $v$  in  $D_1^2(K)$ . ■

*Proof* : We prove the assertion for  $j = 1$ , the proof for  $j = 0$  being analogous. We set (Euler's variable change) :

$$r = e^s; \quad z = \mu t,$$

where  $\mu$  is a constant chosen in such a way that  $K$  is transformed in a square  $\hat{K}$ , i.e.,

$$\mu = h/\log \left( 1 + \frac{h}{\bar{r}} \right) = \bar{r} + \theta h; \quad 0 < \theta < 1.$$

$w = v - \pi v$  is transformed in  $\hat{w}$  by  $w(r, z) = \hat{w}(s, t)$  and  $\hat{w} = \hat{v} - \hat{\pi}\hat{v}$ ,  $\hat{\pi}$  being the usual interpolation operator by polynomials of partial degree less than 1

from the values on the vertex. Then, we have

$$|w|_{1,\hat{K}}^2 = \mu \int_{\hat{K}} \left( \left| \frac{\partial \hat{w}}{\partial s} \right|^2 + \left( \frac{e^s}{\mu} \right)^2 \left| \frac{\partial \hat{w}}{\partial t} \right|^2 \right) ds dt.$$

Since

$$\frac{e^s}{\mu} \leq \frac{(\bar{r} + h)}{\bar{r}} \leq c; c \text{ independent of } h \text{ and } \bar{r};$$

we obtain

$$|w|_{1,K}^2 \leq c\mu \int_{\hat{K}} \left( \left| \frac{\partial \hat{w}}{\partial s} \right|^2 + \left| \frac{\partial \hat{w}}{\partial t} \right|^2 \right) ds dt.$$

Using some standard results of error interpolation (see Ciarlet [4], Raviart [9]), we have

$$|w|_{1,K}^2 \leq c^2 \log^2 \left( 1 + \frac{h}{\bar{r}} \right) \mu \int_{\hat{K}} \left( \left| \frac{\partial^2 \hat{v}}{\partial s^2} \right|^2 + \left| \frac{\partial^2 \hat{v}}{\partial s \partial t} \right|^2 + \left| \frac{\partial^2 \hat{v}}{\partial t^2} \right|^2 \right) ds dt.$$

Taking again the previous variables, we obtain

$$\begin{aligned} |w|_{1,K}^2 &\leq c^2 \log^2 \left( 1 + \frac{h}{\bar{r}} \right) \times \\ &\times \int_{\hat{K}} \left\{ r^2 \left| \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial v}{\partial r} \right) \right|^2 + \mu^2 \left| \frac{\partial^2 v}{\partial r \partial z} \right|^2 + \left( \frac{\mu}{r} \right)^2 \mu^2 \left| \frac{\partial^2 v}{\partial z^2} \right|^2 \right\} r dr dz. \end{aligned}$$

Using

$$\frac{\mu}{r} \leq \frac{\bar{r} + \theta h}{\bar{r}} \leq c; c \text{ constant independent of } h \text{ and } \bar{r},$$

and with some easy majorations, we come to

$$|w|_{1,K}^2 \leq c^2 (\bar{r} + h)^2 \log^2 \left( 1 + \frac{h}{\bar{r}} \right) \left( |v|_{2,K}^2 + \left| \frac{1}{r} \frac{\partial v}{\partial r} \right|_{0,K}^2 \right),$$

where  $c$  is a constant independent of  $h$ ,  $\bar{r}$  and  $v$ . Finally, (3.10) follows from

$$\bar{r} \left( 1 + \frac{h}{\bar{r}} \right) \log \left( 1 + \frac{h}{\bar{r}} \right) \leq \left( 1 + \frac{h}{\bar{r}} \right) h. \quad \blacksquare$$

LEMMA 3.4 : *If  $K$  is a triangle satisfying the following conditions :*  
 $h_K$  : *diameter of  $K$  less than  $h$  ; least interior angle of  $K$  greater than  $\theta_0 > 0$  ;*  
 $\theta_0$  *independent of  $h$  ;  $K$  : contained in the half space  $\{(r, z) \in \mathbb{R}^2 ; r \geq R\}$ ,*  
 $R$  *independent of  $h$  ; then, there exists a constant  $c$  independent of  $h$  such that*

$$|v - \pi v|_{j,K} \leq ch^{2-j} |v|_{2,K}; \quad j = 0, 1, \quad (3.11)$$

for all  $v$  in  $D_1^2(K)$ . ■

*Proof* : It is sufficient to remark that both semi-norms with and without weight are uniformly equivalent far from the  $z$ -axis and then to apply standard error interpolation estimates. ■

The proof of theorem 3.1 results from the above lemmas by standard techniques of majoration of the error over each element which can be reduced by a linear affine transformation keeping the estimates under assumption (3.2) to the cases (3.3), (3.10) and (3.11). ■

#### REFERENCES

1. Y. AMIRAT, *Une méthode d'optimisation de domaine dans le problème du cône d'eau stationnaire*, Thèse de 3<sup>e</sup> cycle, Alger, 1977.
2. Y. ATIK, *Sur une formulation variationnelle et approximation numérique du problème de cône d'eau stationnaire*, Thèse de 3<sup>e</sup> cycle, Alger, 1977.
3. M. CROUZEIX, J. M. THOMAS, *Éléments finis et problèmes aux limites elliptiques dégénérés*, R.A.I.R.O., Analyse Numérique, Paris, décembre 1973.
4. P. G. CIARLET, *The finite element method for elliptic problems*, North Holland, Amsterdam and New York, 1978.
5. A. EL KOLLI,  *$n$ -ième épaisseur dans les espaces de Sobolev et applications*, Thèse d'État, Alger, 1977.
6. P. GRISVARD, *Alternative de Fredholm relative au problème de Dirichlet dans un polygone ou un polyèdre*, BUMI, Bologna, 1972.
7. P. LAILLY, *Résolution numérique des équations de Stokes en symétrie de révolution par une méthode d'éléments finis non conformes*, Thèse de Docteur-Ingénieur, Paris XI, 1976.
8. J. L. LIONS, E. MAGENES, *Problèmes aux limites non homogènes et applications*, vol. 1, Dunod, Paris, 1968.
9. P. A. RAVIART, *Méthode des éléments finis*, Cours de 3<sup>e</sup> cycle, Paris VI, 1972.

