

JEAN BOURGAIN

HARRY FURSTENBERG

YITZHAK KATZNELSON

DONALD S. ORNSTEIN

Appendix on return-time sequences

Publications mathématiques de l'I.H.É.S., tome 69 (1989), p. 42-45

http://www.numdam.org/item?id=PMIHES_1989__69__42_0

© Publications mathématiques de l'I.H.É.S., 1989, tous droits réservés.

L'accès aux archives de la revue « Publications mathématiques de l'I.H.É.S. » (<http://www.ihes.fr/IHES/Publications/Publications.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

APPENDIX

by J. Bourgain, H. Furstenberg, Y. Katznelson, D. S. Ornstein

Return Times of Dynamical Systems

Let (X, \mathcal{B}, μ, T) be an ergodic system and let $A \in \mathcal{B}$ be of positive measure $\mu(A) > 0$. For $x \in X$, consider the return time sequence $\Lambda_x = \{n \in \mathbf{Z}_+ \mid T^n x \in A\}$. By Birkhoff's pointwise ergodic theorem, the sequence Λ_x has positive density for μ -almost all $x \in X$. This fact refines the classical Poincaré recurrence principle (cf. [Fu]). An even stronger statement is given by the Wiener-Wintner theorem: there is a set X' of X of full measure such that the sums

$$\frac{1}{N} \sum_{1 \leq n \leq N} \chi_A(T^n x) z^n$$

converge for all z in the unit circle $\mathbf{C}_1 = \{z \in \mathbf{C} \mid |z| = 1\}$ and $x \in X'$. Thus from general theory of unitary operators, this fact may be reinterpreted by saying that almost all sequences Λ_x satisfy the L^2 , hence the mean ergodic theorem. Our purpose here is to prove the following fact, answering a question open for some time.

Theorem. — *With the notation above, Λ_x satisfies almost surely the pointwise ergodic theorem, i.e., the averages*

$$\frac{1}{N} \sum_{\substack{1 \leq n \leq N \\ n \in \Lambda_x}} S^n g$$

converge almost surely for any measure preserving system (Y, \mathcal{D}, ν, S) and $g \in L^1(Y)$.

The argument given next actually yields a more precise condition on the point x .

Let $f \in L^\infty(X)$ be obtained by projecting χ_A on the orthogonal complement of the eigenfunctions of T . It clearly suffices to prove that for almost all $x \in X$, $\{f(T^n x)\}$ is a "summing sequence", i.e.,

$$(*) \quad \frac{1}{N} \sum_{1 \leq n \leq N} f(T^n x) g(S^n y) \rightarrow 0 \quad \text{a.e. } y \in Y$$

for any measure preserving system (Y, \mathcal{D}, ν, S) and $g \in L^\infty(Y)$. (The contribution of the eigenfunctions is taken care of by Birkhoff's theorem.)

Observe the equivalence of the following statements:

- (i) f has continuous spectral measure,
- (ii) $\langle T^n f, f \rangle = \hat{\sigma}_f(n)$, σ a continuous measure,
- (iii) $(1/N) \sum_{1 \leq n \leq N} f(T^n x) f(T^n \xi) \rightarrow 0$ a.e. in (x, ξ) as $N \rightarrow \infty$.

Proof of (ii) \Rightarrow (iii). — Write $F = \lim (1/N) \sum_1^N f(T^n x) f(T^n \xi)$, a limit which exists by the ergodic theorem, and $\|F\|^2 = \lim (1/N^2) \sum_{m, n=1}^N (\hat{\sigma}_f(n-m))^2 = 0$.

Proposition. — Assume x generic for f and $(1/N) \sum f(T^n x) f(T^n \xi) \rightarrow 0$, a.e. in ξ (!). Then $\{f(T^n x)\}$ is a summing sequence.

Proof. — I) Assume that for some (Y, \mathcal{D}, ν, S) and $g \in L^\infty(Y)$ there is a set B^* of positive measure for which the limsup of $(*)$ is positive. Then there exists $a > 0$, $B \subset B^*$, $\nu(B) > 0$ and a sequence of intervals $R_j = (L_j, M_j)$ (called “ranges”) such that for every $y \in B$ and every j there exists $n_j \in \mathbf{R}_j$ ($n_j = n_j(y)$) such that

$$(**) \quad \sum_{n=1}^{n_j} f(T^n x) g(S^n y) > an_j.$$

II) Given $\delta > 0$, there exists $K = K(N, \delta)$ such that

$$\nu\left(\bigcup_1^K S^j B\right) > 1 - \delta.$$

III) Write φ for the indicator function of $\bigcup_{j=1}^K S^j B$. If M_0 is large enough, and if we denote by G the set $G = \{y : |(1/n) \sum_1^n \varphi(S^j y) - 1| < 2\delta \text{ for all } n > M_0\}$, then $\nu(G) > 1 - \delta$.

IV) For notational convenience we assume that f has finite range, and we denote by B_n the set of all n -blocks for f , i.e., the set of words $w_k^{(n)} = (f(T^{k+1}x), \dots, f(T^{k+n}x))$; $w_k^{(n)}$ appears with density $p(w_k^{(n)})$.

Given $\delta > 0$ (δ can be chosen once and for all as a function of a and $\nu(B)$ in I) let N_δ be such that for each set $A_\delta \subset X$, $\mu(A_\delta) > 1 - \delta$, $|(1/N) \sum f(T^n x) f(T^n \xi)| < \delta$ for all $\xi \in A_\delta$ and $N > N_\delta$ (cf. assumption (!)).

Given a range (L, M) with $L > N_\delta$, set $N = N(M)$ so that in any interval on the integers of length $\geq N$ the statistics of the n -blocks (for f) with $n \leq M$ is correct. Denote by B_n^* the n -blocks that have the form $(f(T\xi), \dots, f(T^n \xi))$ with $\xi \in A_\delta$ (we are interested in $n \in (L, M)$). For $L < n < M$ the total probability (= density) of the blocks in B_n^* exceeds $1 - \delta$ (in any interval of length $\geq N(M)$). Notice also that heads of M -blocks which are in B_M^* are in the appropriate B_n^* .

V) A sequence of ranges $\{(L_j, M_j)\}$ is *properly spaced* if $L_{j+1} > N(M_j)$. (We also assume $L_1 > N_\delta$. Another assumption on L_1 is that it is $> M_0$ (recall the definition of G in III) and assume that K (II)) is $\ll L_1$.) *Going back to I*), we select a properly spaced sequence of ranges $\{(L_j, M_j)\}_{j=1}^J$ (J depending on a) and N large enough so that $N \geq N(M_j)$.

Recall B from I) and G from III).

For any $y \in B \cap G$ we define a sequence $\{c_n(y)\}_{n=1}^N$ which is a sum of J sequences (layers) $\{c_n^j(y)\}$ having the following properties:

(α) For all j, n and y , $c_n^j(y)$ is in the range of f (in particular uniformly bounded)

(β) For $j_1 \neq j_2$, $|(1/N) \sum_{n=1}^N c_n^{j_1}(y) c_n^{j_2}(y)| < \delta$

(γ) $(1/N) \sum_{n=1}^N c_n^j(y) g(S^n y) > a - \delta$, $j = 1, \dots, J$

(α) and (β) together imply $[(1/N) \sum_{n=1}^N (c_n(y))^2]^{1/2} = O(\sqrt{J} + \delta J)$, and (γ) implies $(1/N) \sum_{n=1}^N c_n(y) g(S^n y) > J(a - \delta)$. Contradiction.

We construct $\{c_n^j\}$ in reverse order on j . The number $c_n^J(y)$ is defined as follows: $\ell_1(y)$ is the first index $k > 0$ such that $S^k y \in B$; on the interval $(\ell_1(y), \ell_1(y) + n_J(S^{\ell_1(y)} y))$ we set

$$c_n^J(y) = f(T^{n - \ell_1(y)} x),$$

$\ell_2(y)$ is the index of the first point in the S -orbit of y after $\ell_1(y) + n_J(S^{\ell_1(y)} y)$ which is in B , and on the interval $(\ell_2(y), \ell_2(y) + n_J(S^{\ell_2(y)} y))$ we copy again $\{f(T^k x)\}_{k=1}^{n_J(S^{\ell_2(y)} y)}$ etc. The intervals on which we copy those starting n_J blocks fill most of $[1, N]$. We refer to these as the basic intervals of the J -layer. Outside of these, set $c_n^J(y)$ arbitrarily.

We now define $c_n^{J-1}(y)$ in a similar manner within every basic interval of the J -layer, with the additional restriction on the starting place of the new basic blocks that (in addition to the fact that the corresponding point in the orbit of y is in B) the matching piece of the basic J -layer block in is B^* , i.e., more or less orthogonal to the “new” basic block; see IV). Since the “orthogonal” blocks have density $> 1 - \delta$, the new basic blocks cover more than $1 - 3\delta$ of $[1, N]$. We continue with $c_n^{J-2}(y), \dots, c_n^1(y)$, working each time within the basic blocks of the previous level and introducing blocks which are “orthogonal” to all previous levels.

Remarks.

- (i) The condition that $(1/N) \sum_{n=1}^N f(T^n x) f(T^n \xi) \rightarrow 0$ a.e. in $\xi(!)$ is a special case of (*) and hence necessary. One can construct examples showing that it is not a consequence of the genericity of x .
- (ii) One may construct a sequence $\Lambda = \{k_n\}$, $k_n = o(n)$, and a weakly mixing system (Y, S) such that $(1/N) \sum_{n=1}^N g(S^{k_n} y)$ does not converge a.e., for some $g \in L^\infty(Y)$. (This question was considered in [Fu], p. 96.)

REFERENCES

- [B₁] J. BOURGAIN, On the maximal ergodic theorem for certain subsets of the integers, *Israel J. Math.*, **61** (1) (1988), 39-72.
- [B₂] J. BOURGAIN, On the pointwise ergodic theorem on L^p for arithmetic sets, *ibid.*, 73-84.

- [B₃] J. BOURGAIN, An approach to pointwise ergodic theorems, *Springer LNM*, **1317** (1988), 204-223.
- [B₄] J. BOURGAIN, *Return time sequences of dynamical systems*, preprint IHES, 3/1988.
- [B₅] J. BOURGAIN, Almost sure convergence and bounded entropy, *Israel J. Math.*, **63** (1) (1988), 79-97.
- [B₆] J. BOURGAIN, Temps de retour pour les systèmes dynamiques, *CRASc Paris*, Ser I, **306** (1988), 483-485.
- [Fu] H. FURSTENBERG, *Recurrence in ergodic theory and combinatorial number theory*, Princeton UP, 1981.
- [Ga] A. GARSIA, *Martingale Inequalities*, Benjamin, 1970.
- [K-W] Y. KATZNELSON, B. WEISS, A simple proof of some ergodic theorems, *Israel J. Math.*, **42** (4) (1982), 391-395.
- [Lé] D. LÉPINGLE, La variation d'ordre p des semi-martingales, *Z. Wahrscheinlichkeitstheorie verw. Gebiete*, **36** (1976), 295-316.
- [Ma] J. M. MARSTRAND, On Khinchine's conjecture about strong uniform distribution, *Proc. London Math. Soc.*, **21** (1970), 540-556.
- [Nev] J. NEVEU, *Martingales à temps discret*, Masson, 1972.
- [Ra] D. A. RAIKOV, On some arithmetical properties of summable functions, *Mat. Sb.*, **1** (43) (1936), 377-384.
- [Ri] F. RIESZ, Sur la théorie ergodique, *Comment. Math. Helv.*, **17** (1945), 221-239.
- [S] R. SALEM, *Collected works*, Hermann, Paris, 1967.
- [St] E. STEIN, On limits of sequences of operators, *Ann. Math.*, **74** (1961), 140-170.
- [Vaug] R. C. VAUGHAN, *The Hardy-Littlewood method*, Cambridge Tracts, **70** (1981).
- [Vin] I. M. VINOGRADOV, *The method of trigonometrical sums in the theory of numbers*, Interscience N. Y., 1954.
- [W] B. WEISS, Private communications.
- [Wl] M. WIERDL, Pointwise ergodic theorem along the prime numbers, *Israel J. Math.*, **64** (1988), 315-336.

J. B.:
Institut des Hautes Études Scientifiques
35, route de Chartres
91440 Bures-sur-Yvette
France

H. F.:
Hebrew University
91904 Jerusalem
Israel

Y. K. and D. O.:
Department of Mathematics
Stanford University
Stanford, California 94305
États-Unis

Manuscrit reçu le 30 décembre 1988.