

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

GUEORGUIY V. BABIKOV

On the structure of Clifford algebras

Rendiconti del Seminario Matematico della Università di Padova,
tome 33 (1963), p. 267-269

http://www.numdam.org/item?id=RSMUP_1963__33__267_0

© Rendiconti del Seminario Matematico della Università di Padova, 1963, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE STRUCTURE OF CLIFFORD ALGEBRAS

*Nota ** di GUEORGUIY V. BABIKOV (a Sverdlovsk)

The principal result of the paper by G. B. Rizza published in this journal ¹⁾ may be defined as following:

« Clifford algebra C_n of index n over the field of real numbers is direct sum of 2^{n-2} fields of quaternions. »

This result is wrong. The mistake may be found in the inattentive application of the method of mathematical induction.

After the above given result has been considered and proved for $n = 3$ the process of proovement for $n > 3$ follows in such a way.

Let $x = \alpha_1 + \alpha_2 i_n \in C_n$ and $\alpha_1, \alpha_2 \in C_{n-1}$.

Suppose that $C_{n-1} = C_{\frac{1}{2}}^{(1)} + C_{\frac{1}{2}}^{(2)} + \dots + C_{\frac{1}{2}}^{(2^{n-2})}$ where $C_{\frac{1}{2}}^{(k)}$ is isomorphic to a field of quaternions.

Then $x = \sum_{\frac{1}{2}}^{2^{n-2}} [\alpha_1^{(k)} + \alpha_2^{(k)} i_n]$ and the sets $\{\alpha_1^{(k)} + \alpha_2^{(k)} i_n\}$ are isomorphic to C_3 when $\alpha_1^{(k)}$ and $\alpha_2^{(k)}$ run through all possible real quaternions. This last statement is wrong for $n \geq 4$. If a set $\{\alpha_1^{(k)} + \alpha_2^{(k)} i_n\}$ is isomorphic to $C_3 = C_{\frac{1}{2}}^{(1)} + C_{\frac{1}{2}}^{(2)}$, then i_n and base units of $C_{\frac{1}{2}}^{(k)}$ have specific forms depending on base units of $C_{\frac{1}{2}}^{(1)}$ and $C_{\frac{1}{2}}^{(2)}$.

The result by Rizza disproves also the following contradictory example.

***)** Pervenuta in redazione il 5 gennaio 1963.

Indirizzo dell'A.: Pedagogical institute, Sverdlovsk (U.R.S.S.)

Suppose we have algebra C_n ($n > 3$) over the field of real numbers. Let us take an element $x = i_1 + i_2 i_3 i_4$.

Let $x = q_1 \dot{+} q_2 \dot{+} \dots \dot{+} q_{2^{n-2}}$, where q_k belongs to some field of quaternions $C_4^{(k)}$.

Then $x^2 = q_1^2 \dot{+} q_2^2 \dot{+} \dots \dot{+} q_{2^{n-2}}^2 = (i_1 + i_2 i_3 i_4)^2 = 0$. As the above sum is direct, we have $q_1^2 = q_2^2 = \dots = q_{2^{n-2}}^2 = 0$. Consequently $q_1 = q_2 = \dots = q_{2^{n-2}} = 0$, i.e. $x = 0$.

We come to a contradiction.

The structure of Clifford algebras defined by the following theorem 2):

« Clifford algebras over the arbitrary field F of characteristic $\neq 2$ under different n (with exactness to isomorphism) are exhausted by the following list:

- 1) C_{8m} is algebra of matrices of power 2^{4m} over F .
- 2) C_{8m+1} is direct product $A \times K$, where A is algebra of matrices of power 2^{4m} over F and K is algebra of complex numbers over F .
- 3) C_{8m+2} is direct product $A \times Q$, where A is algebra of matrices of power 2^{4m} over F and Q is algebra of quaternions over F .
- 4) C_{8m+3} is algebra $Q \times A \dot{+} Q \times B$, where A and B are algebras of matrices of power 2^{4m} over F and Q is algebra of quaternions over F .
- 5) C_{8m+4} is direct product $A \times Q$, where A is algebra of matrices of power 2^{4m+1} over F and Q is algebra of quaternions over F .
- 6) C_{8m+5} is direct product $A \times K$, where A is algebra of matrices of power 2^{4m+2} over F and K is algebra of complex numbers over F .
- 7) C_{8m+6} is algebra of matrices of power 2^{4m+3} over F .
- 8) C_{8m+7} is direct sum $A \dot{+} B$, where A and B are algebras of matrices of power 2^{4m+3} over F . »

Relying upon this it's possible to solve the question about the distribution of divisors of zero in Clifford algebras.

- [1] RIZZA G. B.: *Sulla struttura delle algebre di Clifford*. Rend. sem. mat. Univ. Padova, 1954, 23, N. 1, 91-99.
- [2] BABIKOV G. V.: *Predstavleniya algebr Clifforda nad proizvol'nimi polyanii*. Uchenie zapiski Uralskogo Universiteta. 1960, Vip. 23 N. 2. tetr. 3. s. 19-26