

RENDICONTI  
*del*  
SEMINARIO MATEMATICO  
*della*  
UNIVERSITÀ DI PADOVA

A. S. VATSALA

R. L. VAUGHN

**Existence of extremal solutions and comparison results  
for delay differential equations in abstract cones**

*Rendiconti del Seminario Matematico della Università di Padova,*  
tome 64 (1981), p. 1-14

[http://www.numdam.org/item?id=RSMUP\\_1981\\_\\_64\\_\\_1\\_0](http://www.numdam.org/item?id=RSMUP_1981__64__1_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1981, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## Existence of Extremal Solutions and Comparison Results for Delay Differential Equations in Abstract Cones.

A. S. VATSALA and R. L. VAUGHN (\*)

### 1. - Introduction.

The study of Cauchy problem for ordinary differential equations in a Banach space has been extensive [2, 4, 10]. It is of interest to look at the corresponding problem for delay differential equations since such equations occur in many physical problems. Existence of solutions of such equations are considered in [8, 9, 6] using monotonicity conditions and dissipative conditions.

In this paper our objective is to prove the existence of extremal solutions for the delay differential equation

$$(1.1) \quad x'(t) = f(t, x, x_t), \quad x_{t_0} = \varphi_0$$

relative to a cone  $k$  of the Banach space  $E$ . For this purpose, we begin by proving an existence result under a simple set of conditions without assuming uniform continuity on  $f$ , we then develop needed theory of differential inequalities and utilize this to show the existence of extremal solutions for (1.1). Several useful comparison theorems are then proved including a flow invariance result. Our results generalize some of the recent results for equations without delay [5, 7].

(\*) Indirizzo degli AA.: A. S. Vatsala, Department of Mathematics, University of Texas at Arlington, Arlington, Texas 76019; R. L. Vaughn, Department of Mathematics, Texas Christian University, Forth Worth, Texas 76129.

Research partially supported by U.S. Army Research Grant No. DAAG29-77-G0062.

## 2. – Preliminaries and definitions.

Let  $\tau > 0$  be a given real number and let  $E$  be a Banach space with norm  $\|\cdot\|$ . Let  $\mathcal{C} = C[[-T, 0], E]$  denote the Banach space of continuous functions with the norm of  $\varphi \in \mathcal{C}$  given by

$$\|\varphi\|_0 = \max_{-\tau \leq s \leq 0} \|\varphi(s)\|.$$

If  $t_0 \in R^+$  and  $x \in C[[t_0 - \tau, \infty), E]$ , then for any  $t \in [t_0, \infty)$ , we let  $x_t \in \mathcal{C}$  be defined by

$$x_t(s) = x(t + s), \quad -\tau \leq s \leq 0.$$

Also let

$$\mathcal{C}_\varrho = \{\varphi \in \mathcal{C}: \|\varphi\|_0 < \varrho\}.$$

Similarly if  $A \subset C[[t_0 - \tau, \infty), E]$  and for  $J \subset [t_0, \infty)$  we will let

$$A_J = \{x_t: x \in A, t \in J\}.$$

To establish existence criteria for the Cauchy problem (1.1) we require that  $f$  satisfies a compactness condition. The compactness condition for this paper will be given in terms of Kurtowski measure of noncompactness  $\alpha$ . The measure of noncompactness  $\alpha(s)$  is defined by

$$\alpha(s) = \inf \{d > 0: s \text{ can be covered by a finite number of sets of diameter } d\}$$

for each bounded subset of  $E$ . We denote  $\alpha_E, \alpha_B, \alpha_C$  to denote the Kurtowski measure of noncompactness defined relative to the Banach spaces  $E, B$  and  $C$  respectively.

A cone  $k$  is a proper subset of  $E$  such that if  $v, \omega \in k, \lambda \in R^+$  then  $v + \omega, \lambda v \in k$ . Throughout this paper we will consider a closed cone  $k$  and its interior  $k^0$ . These cones induce orderings on  $E$  defined by

$$\begin{aligned} x \leq y & \quad \text{if } y - x \in k \\ x < y & \quad \text{if } y - x \in k^0. \end{aligned}$$

For some of the fundamental properties of  $\alpha$  and the cone  $k$  see [2, 3, 4, 10].

Let  $k^*$  be the set of all continuous linear functionals  $c$  on  $E$  such that  $c(x) \geq 0$  for all  $x \in k$ , and let  $k_0^*$  be the set of all continuous linear functionals  $c$  on  $E$  such that  $c(x) > 0$  for  $x \in k^0$ .

A function  $f \in C[J \times E \times C, E]$  is said to be quasimonotone nondecreasing in  $x$  for fixed  $t, \varphi$  if  $x \leq y$  and  $c(x) = c(y)$  for  $c \in k_0^*$  then  $c(f(t, x, \varphi)) \leq c(f(t, y, \varphi))$ .

A function  $f \in C[J \times E \times C, E]$  is said to be monotone nondecreasing in  $\varphi$  for fixed  $t, x$ , if for  $c \in k_0^*$   $\varphi_1(s) \leq \varphi_2(s)$  then  $c(f(t, x, \varphi_1)) \leq c(f(t, x, \varphi_2))$ .

A function  $f \in C[J \times E \times C, E]$  is said to satisfy one sided Lipschitz's condition if there exists an  $L > 0$  such that

$$f(t, x, \varphi_1) - f(t, y, \varphi_2) \leq L(x - y) + \sup_{s \in [-\tau, 0]} \{\varphi_1(s) - \varphi_2(s)\}$$

whenever  $x \geq y$  and  $\varphi_1 \geq \varphi_2$ .

The function  $f(t, x, \varphi)$  is said to be quasinonnegative if  $x \geq 0, \varphi \geq 0, c(x) = 0$  for  $c \in k_0^*$  implies  $c(f(t, x, \varphi)) \geq 0$ . If in the above definition if the inequalities are reversed then  $f$  is said to be quasinonpositive.

In what follows when we say  $x_t \leq y_t$  we mean  $x_t(s) \leq y_t(s), -\tau \leq s \leq 0$ .

A closed set  $F \subset E$  is said to be flow invariant relative to the system (1.1) if for every solution  $x(t)$  of (1.1), we have

$$x_{t_0} \in F, \quad \text{implies } x(t) \in F \text{ for } t \geq t_0.$$

We state below the Darbo fixed point theorem [1] and Mazur's theorem [5] which are needed in our existence theorem and on the results on differential inequalities respectively.

**THEOREM 2.1 (Darbo).** Let  $E$  be a Banach space and  $A$  be a closed, bounded convex, nonvoid subset of  $E$ . If  $T \in C[A, A]$  is such that  $\alpha(T(B)) \leq k\alpha(B)$  where  $k < 1$  for each bounded subset  $B$  of  $A$ , then  $T$  has a fixed point.

**THEOREM 2.2 (Mazur's).** Let  $k$  be a cone with nonempty interior  $k_0$ . Then

- (i)  $x \in k$  is equivalent to  $c(x) \geq 0$  for all  $c \in k^*$ ,
- (ii)  $x \in \partial k$  implies that there exists a  $c \in k_0^*$  such that  $c(x) = 0$ .

### 3. - Existence.

In this section we prove the existence of a solution of the Cauchy problem (1.1)

**THEOREM 3.1.** Let  $f \in C[J \times E \times C_e, E]$  where  $J = [t_0, t_0 + a]$ ,  $a > 0$  and suppose the following conditions are satisfied.

$$(a_1) \quad \|f(t, x, \varphi)\| \leq M \text{ for all } t, x, \varphi \in J \times E \times C_e$$

$$(a_2) \quad \alpha_E(f(J \times A_1 \times A_2)) \leq \beta \max(\alpha_E(A_1), \alpha_C(A_2))$$

where  $\beta > 0$  and  $A_1$  and  $A_2$  are any bounded subsets of  $E$  and  $C$  respectively. Then given an initial function  $\varphi_0 \in C_e$  at  $t = t_0 \geq 0$  there exists a  $\gamma > 0$  such that there is a solution  $x(t, t_0, \varphi_0)$  of (1.1) existing on  $[t_0 - \tau, t_0 + \gamma)$ .

**PROOF.** Define  $y \in C[[t_0 - \tau, t_0 + a], E]$  as follows:

$$y(t) = \begin{cases} \varphi_0(t - t_0) & t_0 - \tau \leq t \leq t_0 \\ \varphi_0(0) & t_0 \leq t \leq t_0 + a. \end{cases}$$

Then  $f(t, y, y_t)$  is a continuous function of  $t$  on  $[t_0, t_0 + a]$  and hence  $\|f(t, y, y_t)\| \leq M$  by  $(a_1)$ . We can show [11] that there exists a constant

$$b \in (0, \varrho - \|\varphi_0(0)\|)$$

such that

$$\|f(t, x, \psi) - f(t, y, y_t)\| \leq 1$$

whenever  $t \in [t_0, t_0 + a]$ ,  $x_t = \psi \in C_e$  and  $\|\psi - y_t\|_0 \leq b$ .

If now follows that  $\|f(t, x, \psi)\| \leq M + 1$  whenever  $t \in [t_0, t_0 + a]$ ,  $x_t = \psi \in C_e$  and  $\|\psi - y_t\|_0 \leq b$ . Choose  $\gamma = \min(a, (b/M + 1), 1)$ .

Let  $B$  denote the space of continuous functions from  $[t_0 - \tau, t_0 + \gamma]$  into  $E$ . For an element  $x \in B$  define the norm

$$\|x\|_0 = \max_{t_0 - \tau \leq t \leq t_0 + \gamma} \|x(t)\|.$$

Then  $B$  is a Banach space with respect to this norm. Let  $S \subset B$  be

defined as follows

$$S = \begin{cases} x \in B & \text{(i) } x(t) = \varphi_0(t - t_0) & t_0 - \tau \leq t \leq t_0 \\ & \text{(ii) } \|x(t_1) - x(t_2)\| \leq M|t_1 - t_2| & t_1, t_2 \in [t_0, t_0 + \gamma]. \end{cases}$$

clearly  $S$  is closed, bounded, convex and equicontinuous.

We now define a mapping on  $S$  as follows. For an element  $x \in S$ , let

$$\begin{aligned} \text{(i) } T(x_{t_0}) &= \varphi_0 \\ \text{(ii) } T(x(t)) &= \varphi_0(0) + \int_{t_0}^t f(s, x(s), x_s) ds, \quad t_0 \leq t \leq t_0 + \gamma. \end{aligned}$$

If  $x \in S$ ,  $T(x_{t_0}) = \varphi_0$

$$\|(Tx)(t_1) - T(x)(t_2)\| \leq \int_{t_1}^{t_2} \|f(s, x(s), x_s)\| ds \leq M|t_2 - t_1| \quad t_2 > t_1.$$

Consequently  $TS \subseteq S$  and  $T$  is bounded.

To show  $T$  is continuous let  $\{x_n\} \subseteq S$  be a sequence converging to  $x$ . By the continuity of  $f$  we have

$$f(t, x_n(t), x_n) \rightarrow f(t, x(t), x_t).$$

Further  $\|f(t, x_n(t), x_n)\| \leq M$ . By applying the bounded convergence theorem  $Tx_n \rightarrow Tx$ .

Now let  $\Sigma \subset S$ . Then using the properties of  $\alpha$  for  $t > t_0$ ,

$$\begin{aligned} \alpha_E(T\Sigma(t)) &= \alpha_E\left(\left\{\varphi_0(0) + \int_{t_0}^t f(s, x(s), x_s) ds : x \in \Sigma\right\}\right) \\ &\leq \alpha_E\left(\left\{\int_{t_0}^t f(s, x(s), x_s) ds : x \in \Sigma\right\}\right) \\ &\leq |t - t_0| \alpha_E\left(\overline{\text{co}} f([t_0, t], \Sigma([t_0, t]), \Sigma_{[t_0, t]})\right) \\ &\leq |t - t_0| \max\left\{\alpha_E(\Sigma([t_0, t])), \alpha_C(\Sigma_{[t_0, t]})\right\}. \end{aligned}$$

Since  $S$  is equicontinuous and  $\Sigma \subset S$ , then  $\Sigma$  is also equicontinuous.

Furthermore  $\Sigma_{[t_0, t]}$  is an equicontinuous subset of  $\mathcal{C}$ . Hence we have

$$\alpha_{\mathcal{C}}(\Sigma_{[t_0, t]}) = \alpha_E\left(\Sigma_{[t_0, t]}([- \tau, 0])\right) = \sup_{\sigma \in [- \tau, 0]} \alpha_E(\Sigma_{[t_0, t]}(\sigma)).$$

But  $\Sigma_{[t_0, t]}(\sigma) = \Sigma([t_0 + \sigma, t + \sigma])$ , and so

$$\alpha_{\mathcal{C}}(\Sigma_{[t_0, t]}) \leq \alpha_B(\Sigma([t_0 - \tau, t])) = \alpha_E(\Sigma([t_0, t])).$$

We may then write

$$\begin{aligned} \alpha_B(T\Sigma(t)) &\leq |t - t_0| \alpha_B(\Sigma([t_0, t])) \leq \frac{1}{2} \alpha_E(\Sigma([t_0, t])) \\ &\leq \frac{1}{2} \alpha_B(\Sigma) \end{aligned}$$

Consequently,

$$\alpha_B(T\Sigma) = \sup_{t \in [t_0, t]} \alpha_B(T\Sigma(t)) \leq \frac{1}{2} \alpha_B(\Sigma).$$

By the Darbo fixed point Theorem there exists a fixed point  $x$  of  $T$ . Such a fixed point is a solution of (1.1).

#### 4. - Differential inequalities.

In this section we develop the theory of differential inequalities which is used as a tool to prove the existence of extremal solutions and a comparison result. All the inequalities are relative to a cone  $k$  of the Banach space  $E$ .

**THEOREM 4.1.** Let  $f \in C[J \times E \times \mathcal{C}, E]$  (where  $J = (t_0, \infty)$ ) and  $f$  is quasimonotone nondecreasing in  $x$  for fixed  $(t, \varphi)$  and monotone nondecreasing in  $\varphi$  for fixed  $(t, x)$ . Let  $x, y \in C[[t_0 - \tau, \infty), E]$  and

$$x_{t_0} < y_{t_0}.$$

Assume further that

$$(4.1) \quad D_- x(t) \leq f(t, x(t), x_t)$$

$$(4.2) \quad D_- y(t) \geq f(t, y(t), y_t)$$

for  $t \in (t_0, \infty)$ . Then

$$x(t) < y(t) \quad t \in [t_0, \infty)$$

provided one of the above inequalities is taken strict.

PROOF: Suppose that the assertion of the Theorem is false. Then there exists a  $t_1 > t_0$  such that  $y(t_1) - x(t_1) \in \partial k$  and  $y(t) - x(t) \in k_0$ ,  $t \in [t_0 - \tau, t_1)$ . Thus by Mazur's Theorem, there exists a  $c \in k_0^*$  with  $c(y(t_1) - x(t_1)) = 0$ . Setting  $m(t) = c[y(t) - x(t)]$ , we see that  $m(t) > 0$  for  $t_0 - \tau \leq t < t_1$  and  $m(t_1) = 0$ . Consequently  $D_- m(t_1) \leq 0$ .

Also

$$(4.3) \quad \begin{aligned} D_- m(t_1) &= C[D_- y(t_1) - D_- x(t_1)] \\ &> C[f(t_1, y(t_1), y_{t_1}) - f(t_1, x(t_1), x_{t_1})] \end{aligned}$$

by (4.1), (4.2). Using the quasimonotonicity of  $f$  in  $x$  and monotonicity of  $f$  in  $\varphi$  it follows that

$$(4.4) \quad c(f(t_1, y(t_1), y_{t_1})) \geq c(f(t_1, x(t_1), x_{t_1})).$$

Now (4.3), (4.4) together imply

$$D_- m(t_1) > 0.$$

Hence a contradiction and the theorem is proved.

REMARK 4.1. The above result is valid if in the inequalities (4.1), (4.2)  $D_-$  is replaced by any other diniderivative. See [11] for details.

REMARK 4.2. The conclusion of the above theorem is not valid if one of the inequalities in (4.1), (4.2) is not taken strict. However the conclusion is valid if further  $f$  satisfies a onesided Lipschitz condition.

THEOREM 4.2. Let  $f \in C[J \times E \times C, E]$  satisfy the assumptions of Theorem 4.1. Further let  $f$  satisfy the onesided Lipschitz's condition. Then  $x_{t_i} \leq y_{t_i}$  together with (4.1) (4.2) imply  $x(t) \leq y(t)$ ,  $t \in J$ .

PROOF. Consider  $\tilde{x}(t) = x(t) - \varepsilon \exp[3L(t - t_0)]$  where  $\varepsilon \in k^0$ , certainly  $\tilde{x}(t) < x(t)$ ,  $\tilde{x}_{t_i} < x_{t_i}$  for  $t \geq t_0$ .

Also consider

$$\begin{aligned} D_-\tilde{x}(t) &= D_-x(t) - 3L\varepsilon \exp[3L(t-t_0)] \\ &\leq f(t, x, x_t) - 3L\varepsilon \exp[3L(t-t_0)] \end{aligned}$$

by (4.1). Using the on-sided Lipschitz's condition it follows that

$$f(t, x, x_t) \leq f(t, \tilde{x}, \tilde{x}_t) + 2L\varepsilon \exp[3L(t-t_0)].$$

Now we can conclude that

$$(4.5) \quad D_-\tilde{x}(t) < f(t, \tilde{x}, \tilde{x}_t) \quad \text{and} \quad \tilde{x}_{t_0} < x_{t_0} \leq y_{t_0},$$

Now applying Theorem 4.1 to (4.5) and (4.2), it follows that

$$(4.6) \quad \tilde{x}(t) < y(t).$$

Taking the limit as  $\varepsilon \rightarrow 0$  in (4.6) we get the required result.

## 5. - Existence of extremal solution.

In this section we prove the existence of maximal solution of 1.1 only. The existence of minimal solutions can be proved on similar lines.

**THEOREM 5.1.** Let  $f \in C[J \times E \times C_\rho, E]$  and suppose the assumptions  $(a_1)$  and  $(a_2)$  of Theorem 3.1 are satisfied and also further  $f(t, x, \varphi)$  is quasimonotone nondecreasing in  $x$  for fixed  $(t, \varphi)$  and monotone nondecreasing in  $\varphi$  for fixed  $(t, x)$  then given an initial function  $\varphi_0 \in C_\rho$  at  $t = t_0 \geq 0$  there exists a  $\gamma > 0$  such that there is a maximal solution  $r(t, t_0, \varphi_0)$  of (1.1) existing on  $[t_0 - \tau, t_0 + \gamma)$ .

**PROOF.** Consider the delay differential equation

$$(5.1) \quad x'(t) = f(t, x, x_t) + \frac{1}{n}y_0, \quad x_{t_0} = \varphi_0 + \frac{1}{n}y_0 = \varphi_0^n$$

where  $y_0 \in k^0$  and  $\|y_0\| = 1$  for  $n = 1, 2, 3, \dots$

The solutions of (5.1) for  $n = 1, 2, 3, \dots$  exists on some interval  $[t_0 - \tau, t_0 + \gamma)$ ,  $\gamma > 0$  from Theorem 3.1.

Let  $\{x_n\}$  be any sequence of solutions for (5.1) for  $n = 1, 2, 3, \dots$  respectively. That is  $x_n$  is a solution of

$$x'(t) = f(t, x, x_t) + \frac{1}{n} y_0, \quad x_{t_0} = \varphi_0 + \frac{1}{n} y_0 = \varphi_0^n$$

and  $x_{n+1}$  is a solution of

$$x'(t) = f(t, x, x_t) + \frac{1}{n+1} y_0, \quad x_{t_0} = \varphi_0 + \frac{1}{n+1} y_0 = \varphi_0^{n+1}.$$

Using Theorem 4.1 we can conclude  $x_n > x_{n+1}$  i.e.  $\{x_n\}$  is a monotone decreasing sequence bounded by  $x(t)$  the solution of (1.1). It can be easily shown that  $\{x_n\}$  is equicontinuous, uniformly bounded and also  $\alpha(\{x_n\}) = 0$ . That is  $\{\overline{x_n}(t)\}$  is compact. Hence by Asoli's lemma there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  which converges. Suppose  $\{x_{n_k}\}$  converges to  $r(t)$ . By assumption  $(a_1)$ ,  $\|f(t, x_n(t), x_{n_t})\| \leq M$  and thus the bounded convergence theorem implies

$$\int_{t_0}^t \left\{ f(s, x_{n_k}(s), x_{n_{k_t}}) + \frac{1}{n} y_0 \right\} ds \rightarrow \int_{t_0}^t f(s, r(s), r_s) ds.$$

This proves that  $r(t)$  is a solution of (1.1).

If  $x(t)$  is any solution of (1.1) then by theorem 4.1 it follows that

$$x_{n_k}(t) > x(t) \quad \text{for } t \in [t_0, t_0 + \gamma].$$

This implies that

$$x(t) \leq \lim_{n_k \rightarrow \infty} x_{n_k}(t) = r(t) \quad \text{for } t \in [t_0, t_0 + \gamma].$$

Thus  $r(t)$  is the maximal solution of (1.1) on  $[t_0, t_0 + \gamma]$ .

## 6. – Comparison results.

In this section we give some comparison theorems related to the system 1.1.

**THEOREM 6.1.** Assume that  $f \in C[J \times E \times \mathbb{C}, E]$  be quasinonnegative and that  $f$  satisfies the onesided Lipschitz's condition. Then the closed set  $\bar{Q}$  is flow invariant relative to the system (1.1) where  $Q = [x \in E, x > 0]$ .

**PROOF.** Set  $m(t) = x(t) + \varepsilon \exp [3Lt]y_0$ , where  $x(t)$  is any solution of (1.1) such that  $x_{t_0} \in \bar{Q}$ ,  $y_0 \in k^0$  and  $\varepsilon > 0$  is arbitrarily small number.

Suppose the conclusion is false, then there exists a least  $t_1$  and  $c \in k_0^*$  such that  $m(t_1) \geq 0$  and  $c(m(t_1)) = 0$  and  $m(t) \in k^0$  for  $t \in [t_0 - \tau, t_1]$ . This implies  $c(m(t_1) - m(t_1 - h)) < 0$  for small  $h > 0$  and therefore  $c(m'(t)) < 0$ .

Also

$$\begin{aligned} c(m'(t_1)) &= c(x'(t_1) + 3\varepsilon \exp [3Lt]y_0) \\ &= c(f(t_1, x(t_1), x_{t_1}) + 3\varepsilon \exp [3Lt]y_0) \\ &\geq c(f(t_1, m(t_1), m_{t_1}) + \varepsilon \exp [3Lt]y_0) \end{aligned}$$

using the Lipschitz's condition. Further using the quasimonotonicity of  $f$  we can conclude

$$c(f(t_1, m(t_1), m_{t_1})) \geq 0$$

which proves that

$$c(m'(t_1) \geq c(\varepsilon \exp [3Lt]y_0)) > 0 .$$

This leads to a contradiction. Thus  $m(t) > 0$  for  $t \in [t_0 - \tau, \infty)$ . Taking the limit as  $\varepsilon \rightarrow 0$  the conclusion follows.

**REMARK 6.1.** Theorem 4.2 can be obtained as a consequence of Theorem 6.1.

For this purpose set  $d = y - x$  so that

$$d' = F(t, d, d_t) \equiv f(t, x + d, (x + d)_t) - f(t, x, x_t) + P(t)$$

where  $P(t) = y' - f(t, y, y_t) - x' + f(t, x, x_t)$ .

Clearly  $d_{t_0} \geq 0$  and  $F$  satisfies the onesided Lipschitz's condition. Furthermore if  $d(t) \geq 0$ ,  $c(d(t)) = 0$ ,  $d_t \geq 0$  for  $c \in k_0^*$  then

$$c(F(t, d, d_t)) \geq c(f(t, x + d, (x + d)_t) - f(t, x, x_t)) .$$

Using the form of  $P(t)$  and also using the quasimonotonicity property of  $f$  in  $x$  and monotonicity of  $f$  in  $\varphi$  we can conclude

$$c(f(t, x + d, (x + d)_i) - f(t, x, x_i)) \geq 0$$

whenever  $d_i \geq 0$   $c(d(t)) = 0$ .

This proves that  $F$  is quasimonotone. The claim now follows from Theorem 6.1.

**COROLLARY 6.1.** Assume that  $f$  is quasimonotone and that  $f$  satisfies the one-sided Lipschitz's condition. Then the closed set  $\bar{Q}$  is flow invariant relative to the system 1.1 where  $Q = [x \in E, x < 0]$ .

**COROLLARY 6.2.** Assume  $f \in C[J \times E \times \mathbb{C}, E]$  satisfies the one-sided Lipschitz condition. Assume also that the following condition holds:

If

$$a \leq b, \varphi(s) \leq b, c(x) = c(b) \text{ for } c \in k_0^* \text{ for } s \in [-\tau, 0],$$

$$\text{then } c(f(t, x, \varphi)) \leq 0$$

and if

$$a \leq x, a < \varphi(s), c(a) = c(x) \text{ for } c \in k_0^* \text{ for } s \in [-\tau, 0],$$

$$\text{then } c(f(t, x, \varphi)) \geq 0.$$

Then the closed set  $\bar{W}$ , where  $W = [x \in E, a < x < b, a, b \in E]$  is flow invariant relative to the system 1.1.

We shall next give a comparison result which yields upper and lower bounds for the solutions of 1.1.

**THEOREM 6.2.** Assume that

(i)  $g_1, g_2 \in C[J \times E \times \mathbb{C}, E]$  are quasimonotone nondecreasing in  $x$  and monotone nondecreasing in  $\varphi$  relative to  $k$  and for  $t, x, \varphi \in J \times E \times \mathbb{C}$ ,

$$g_2(t, x, \varphi) \leq f(t, x, \varphi) \leq g_1(t, x, \varphi).$$

(ii)  $r(t), \varrho(t)$  are solutions of

$$r' = g_1(t, r, r_i), \quad r_{t_0} = \psi_0, \quad \varrho' = g_2(t, \varrho, \varrho_i), \quad \varrho_{t_0} = \chi_0$$

respectively existing on  $[t_0, \infty)$ .

(iii)  $f$  satisfies the on-sided Lipschitz's condition, then if  $x(t)$  is any solution of 1.1 existing on  $[t_0, \infty)$  we have

$$\varrho(t) \leq x(t) \leq x(t), \quad t \geq t_0$$

whenever  $\chi_0 \leq \varphi_0 \leq \psi_0$ .

PROOF. Setting  $m(t) = x(t) - \varrho(t)$ , we see  $m(t)$  satisfies the differential equation

$$(6.1) \quad m'(t) = F(t, m(t), m_t), \quad m_{t_0} = \varphi_0 - \chi_0 \geq 0.$$

where

$$\begin{aligned} F(t, m, m_t) &= f(t, x(t), x_t) - g_2(t, \varrho, \varrho_t) \\ &= f(t, m + \varrho, (m + \varrho)_t) - g_2(t, \varrho, \varrho_t). \end{aligned}$$

It is enough to verify that  $F$  satisfies the assumptions of Theorem 6.1. so that the closed set  $\bar{Q}$  is flow invariant relative to system (6.1). Let  $m \geq 0, m_t \geq 0, c(m) = 0$  for some  $c \in K_0^*$ . Then

$$c(F(t, m, m_t)) = c(f(t, m + r, (m + r)_t) - g_2(t, \varrho, \varrho_t)).$$

Since  $g_2$  is quasimonotone nondecreasing in  $\varrho$  and monotone nondecreasing in  $\chi$ , we have

$$c(g_2(t, \varrho, \varrho_t)) \leq c(g_2(t, m + \varrho, (m + \varrho)_t))$$

and this implies together with assumption (i) that

$$c(F(t, m, m_t)) \geq 0$$

proving that  $F$  is quasinnegative. Clearly  $F$  satisfies the on-sided Lipschitz's condition and hence by Theorem 6.1 we get  $x(t) \geq \varrho(t)$  or  $\varrho(t) \leq x(t), t \geq t_0$ .

A similar argument with  $m(t) = x(t) - r(t)$  would yield  $x(t) \leq r(t), t \geq t_0$  completing the proof.

**COROLLARY 6.3.** If  $\bar{W}$  is flow invariant relative to system 1.1 then there exist functions  $g_1, g_2$  which are quasimonotone nondecreasing

in  $x$  for  $x \in E$  and monotone decreasing in  $\varphi$  for  $\varphi \in C$  provided  $E = R^n$ ,  $k = R_+^n$ .

PROOF. We construct  $g_1, g_2$  as follows: for each  $i$ ,  $1 \leq i \leq n$

$$g_{1i}(t, x, \varphi) = \sup [f_i(t, v, \psi): a_j \leq v_j \leq x_j, v_i = x_i \text{ and } a_i \leq \psi_i(s) \leq \varphi_i(s)]$$

$$g_{2i}(t, x, \varphi) = \inf [f_i(t, v, \psi): x_j \leq v_j \leq b_j, x_i = v_i \text{ and } \varphi_i(s) \leq \psi_i(s) \leq b_i].$$

Then clearly the functions  $g_1, g_2$  are quasimonotone nondecreasing in  $x$  and monotone decreasing in  $\varphi$  and satisfy (i) of Theorem 6.2.

Finally we give a comparison theorem which is a consequence of the results in sect. 4 and 5.

THEOREM 6.3. Let  $f \in C[J \times E \times C, E]$  satisfy assumptions  $(a_1), (a_2)$  and  $f$  be quasimonotone nondecreasing in  $x$  for fixed  $(t, \varphi)$  and monotone decreasing in  $\varphi$  for fixed  $(t, x)$ . Let  $r(t)$  be the maximal solution of (1.1) on  $[t_0, t_0 + \gamma)$ . Then if  $u \in C[J \times E \times C, E]$  is such that

$$(6.2) \quad D_- u \leq f(t, u, u_i), \quad u_{i_0} \leq \varphi_0$$

Then  $u(t) \leq r(t)$  for  $t \in [t_0, t_0 + \gamma)$ .

PROOF. Consider

$$(6.3) \quad x'(t) = f(t, x, x_i) + \frac{1}{n} y_0, \quad x_{i_0} = \varphi_0 + \frac{1}{n} y_0.$$

If  $x_n(t)$  is any solution of (6.3) for  $n = 1, 2, 3, \dots$ . Then using Theorem 4.1 we can conclude

$$u(t) < x_n(t), \quad \text{for } n = 1, 2, 3, \dots$$

Then taking the limit as  $n \rightarrow \infty$  we get

$$u(t) \leq \lim_{n \rightarrow \infty} x_n(t).$$

But by Theorem 5.1 we know  $\lim_{n \rightarrow \infty} x_n(t) = r(t)$  the maximal solution of (1.1) and hence the conclusion follows.

## REFERENCES

- [1] DARBO, *Punti uniti in trasformazioni a codominio noncompatto*, Rend. Sem. Mat. Univ. Padova, **24** (1955), pp. 84-92.
- [2] K. DEIMLING, *Ordinary Differential Equations in Banach Spaces*, Lect. Notes, Vol. 596, Springer-Verlag (1977).
- [3] K. KURTOWSKI, *Topology*, Vol. II, Academic Press, New York (1966).
- [4] V. LAKSHMIKANTHAM - S. LEELA, *An introduction to nonlinear differential equations in abstract spaces*, Pergamon Press (1980).
- [5] V. LAKSHMIKANTHAM, *Comparison results for reaction-diffusion equations in Banach space*, Lecture notes of talks delivered at the conference on « A survey of theoretical and numerical trends in nonlinear analysis » at Bari, Italy.
- [6] V. LAKSHMIKANTHAM - A. R. MITCHELL - R. W. MITCHELL, *On the existence of solutions of differential equations of retarded type in a Banach space*, Annals Polonic Mathematics, **35** (1978), pp. 253-260.
- [7] V. LAKSHMIKANTHAM - A. R. MITCHELL - R. W. MITCHELL, *Maximal and minimal solutions and comparison results for differential equations in abstract cones*, Annals Polonic Mathematics, **38** (1977), pp. 317-324.
- [8] V. LAKSHMIKANTHAM - V. LEELA - V. MOURO, *Existence and uniqueness of solutions of delay differential equations on a closed subset of a Banach space*, Nonlinear Analysis, Vol. 2, No. 3, 311-327.
- [9] S. LEELA - V. MOURO, *Existence of solutions in a closed set for delay differential equations in Banach spaces*, Nonlinear Analysis, Vol. 2 (1978), pp. 47-85.
- [10] R. H. MARTIN, *Non linear operators and differential equations in a Banach space*, John Wiley and Sons, New York (1976).
- [11] V. LAKSHMIKANTHAM - S. LEELA, *Differential and integral inequalities*, Vol. I and II, Academic Press (1968).

Manoscritto pervenuto in redazione il 13 novembre 1979.