

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

JAN AMBROSIEWICZ

On the squares of sets of linear groups

Rendiconti del Seminario Matematico della Università di Padova,
tome 75 (1986), p. 253-256

http://www.numdam.org/item?id=RSMUP_1986__75__253_0

© Rendiconti del Seminario Matematico della Università di Padova, 1986, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

On the Squares of Sets of Linear Groups.

JAN AMBROSIEWICZ (*)

The covering problem ($CC \supseteq G$, C -conjugacy class) for the group $PSL(2, q)$ was described in the paper [1]. In this paper we investigate squares of sets $K_w = \{g \in G: |g| = w\}$ of elements of the same order of linear groups. We will say that the group G has the property W if for each w , the set $K_w K_w$ is a subgroup of G (see [2]). We have proved that the set $K_2 K_2$ is not a subgroup in any groups $GL(n, K)$, $SL(n, K)$, $PSL(n, K)$ for $n \geq 3$. We have proved, too, that $K_2 K_2 \leq GL(2, K)$, $K_2 K_2 \leq SL(2, K)$ for $\text{char } K = 2$ and $K_2 K_2 \not\leq GL(2, K)$ for $\text{char } K \neq 2$. The set $K_2 K_2$ of the group $PSL(2, q)$ is described in Theorem 5.

The following Lemma will be useful in the sequel.

LEMMA 1. Let G be a group. An element $g \in K_2^m$ ($m \geq 2$) if and only if there is an element $x \in K_2^{m-1}$, $x \neq g^{-1}$ such that $(gx)^2 = 1$.

PROOF. If $g = s_1 s_2 \dots s_m$, $s_i \in K_2$ then $g^{-1} = x g x$ where $x = s_m s_{m-1} \dots s_2$. Conversely, if $(gx)^2 = 1$ and $x \in K_2^{m-1}$ then $g = x^{-1}(g^{-1}x^{-1})$ and $(g^{-1}x^{-1})^2 = g^{-1}x^{-1}g^{-1}x^{-1} = g^{-1}g = 1$. Since $g^{-1}x^{-1} \neq 1$ by assumption, hence $g \in K_2^m$.

THEOREM 1. The groups $GL(n, K)$, $SL(n, K)$ and $PSL(n, K)$ with $n \geq 3$ over any field K have not property W .

PROOF. Let us consider all possible cases: (i) $|K| > 3$, (ii) $|K| = 2$, (iii) $|K| = 3$.

(*) Indirizzo dell'A.: Institute of Mathematics ul. Dzierzynskiego 15 m 99, 15-099 Białystok - Polonia.

Ad (i). In this case there is an element u such that $u^2 \neq 1$.

If $A = \text{diag}[u^{-1}, u, 1]$, $B = \text{diag}[1, u, u^{-1}]$ with $u^2 \neq 1$ then $A, B \in K_2K_2$, $AB \notin K_2K_2$ by Lemma 1 ($m = 2$). Using this fact we can see that quasideagonal matrices $[A, 1, \dots, 1]$, $[B, 1, \dots, 1]$ belong to the set K_2K_2 but their product do not belong to the set K_2K_2 . In the remain cases we act similarly, namely in the case (ii) one can take

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

and in the case (iii)

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 0 & 2 \\ 0 & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 2 & 2 \end{bmatrix}.$$

Since $A_8 \simeq PSL(4, 2)$, $G_{168} \simeq PSL(3, 2)$ (see [4]) then by Theorem 1 the simple group A_8 and G_{168} have not property W .

REMARK 1. If $X \in K_2 \subset GL(2, K)$ and $\text{char } K = 2$ then $\det X = 1$.

THEOREM 2. If $\text{char } K = 2$ and $|K| > 2$ then $A \in K_2K_2 \subseteq GL(2, K)$ if and only if $\det A = 1$ i.e. $K_2K_2 = SL(2, K)$.

PROOF. Taking into consideration Corollary 4.7 p. 360 of [3] it is sufficient enough to show our proposition for matrices $A = N_1 = [0, 1; -a_2, -a_1]$, $A = N_2 = [k_1, 0; 0, k_2]$.

By Lemma 1

$$(1) \quad N_i \in K_2K_2 \Leftrightarrow \exists_{\substack{X \in K_2 \\ X \neq N_i}} (XN_i = N_i^{-1}X, \det X \neq 0), \quad i = 1, 2.$$

If $N_i \in K_2K_2$ then $\det N_i = 1$ by Remark 1. If $\det N_1 = 1$ then the equation $XN_1 = N_1^{-1}X$ is satisfied by $X = [0, 1; 1, 0]$ for $a_1 \neq 0$ and $X = [u, u + 1; u + 1, u]$ ($u \neq 0, u \neq 1$) for $a_1 = 0$. Therefore $N_1 \in K_2K_2$. The equation $XN_2 = N_2^{-1}X$ has a solution $X \in GL(2, K)$ if and only if $k_1k_2 = 1$ i.e. $\det N_2 = 1$. We can take $X = [0, 1; 1, 0]$ which satisfies other condition of (1).

REMARK 2. If $|K| = 2$ then $K_2K_2 \leq GL(2, K) = SL(2, K) = PSL(2, K)$ what one can verify by calculations.

COROLLARY 2.1. If $\text{char } K = 2$ and $|K| > 2$ then $K_2K_2 = PSL(2, K)$. The proof results from Theorem 2 and from the equality $SL(2, K) = PSL(2, K)$.

THEOREM 3. If $\text{char } K \neq 2$ then $K_2K_2 \not\leq GL(2, K)$.

Indeed, by Lemma 1 and by easy calculations we have

$$[-1, 0; 0, 1], [0, 1; -1, -a_1] \in K_2K_2 (a_1 \neq 0)$$

and

$$[-1, 0; 0, 1] \cdot [0, 1; -1, -a_1] = [0, -1; -1, -a_1] \notin K_2K_2.$$

Theorem 4. The set K_2K_2 is a subgroup of the group $SL(2, K)$.

PROOF. If $\text{char } K = 2$ then Theorem 4 results from Remark 2 and from Theorem 2. If $\text{char } K \neq 2$ then only the matrix $[-1, 0; 0, -1]$ has the order 2 what one can verify by calculations.

THEOREM 5. (a) If in the field K ($\text{char } K \neq 2$) the element -1 is a square then $K_2K_2 = PSL(2, K)$.

(b) If in the field K ($\text{char } K \neq 2, |K| \neq 3$) the element -1 is not a square then $K_2K_2 \not\leq PSL(2, K)$.

PROOF. (a) One can verify that $K_2 = \{[a, b; c, -a], a^2 + bc = -1\}$ in the group $PSL(2, K)$. By Lemma 1 we have

$$(2) \quad A \in K_2K_2 \iff \exists_{\substack{T \in K_2 \\ T \neq A}} A^{-1} = T^{-1}AT.$$

Let $T = [x, y; z, -x], x^2 + yz = -1, A = [a_{11}, a_{12}; a_{21}, a_{22}]$. The condition (2) is equivalent to the solvability of the system equations

$$(3) \quad \begin{cases} x^2 + yz = -1 \\ a_{21}y + a_{12}z = (a_{22} - a_{11})x \end{cases}$$

or to the solvability of the equation

$$(4) \quad [(a_{11} + a_{22})^2 - 4]x^2 + 4a_{12}a_{21} = u^2 \text{ if } a_{12} \neq 0 \text{ or } a_{21} \neq 0.$$

One can verify that in (2) we can take $T = [0, y; y^{-1}, 0]$ if $a_{12} = a_{21} = 0$; $T = [x, (a_{22} - a_{11})a_{21}^{-1}x; 0, -x]$, $x^2 = -1$ if $a_{21} \neq 0$ and $T = [x, 0; (a_{22} - a_{11})a_{12}^{-1}x, -x]$, $x^2 = -1$ if $a_{12} \neq 0$. Therefore $K_2K_2 = PSL(2, K)$.

(b) The matrices $A = [-1, 1; 1, -2]$, $T = [0, 1; -1, 0]$ fulfil the condition (2). Therefore $A \in K_2K_2$ and $A \neq I$.

If $A = [0, 1; -1, \pm 2]$ then the equation (4) has no solution and $A \notin K_2K_2$. Since K_2K_2 is a normal subset and in this case $PSL(2, K)$ is a simple group then $K_2K_2 \not\leq PSL(2, K)$.

REMARK 3. If $|K| = 3$ then K_2K_2 is a subgroup of $PSL(2, K)$ what can be verified by calculations.

COROLLARY 5.1. $K_2K_2 = A_6$ in the alternating group A_6 .

The proof results from the fact $A_6 \simeq PSL(2, K)$ (see [4]) and from (a) of Theorem 5.

REFERENCES

- [1] J. L. BRENNER - L. CARLITZ, *Covering theorems for finite nonabelian simple groups, III*, Rend. Sem. Mat. Uni. Padova, **55** (1976).
- [2] E. AMBROSIEWICZ, *On groups having the property W*, Demonstratio Mathematica, **10** (1977), pag. 2.
- [3] T. W. HUNGEERFORD, *Algebra*, Springer-Verlag, New York, Heidelberg, Berlin, 1974.
- [4] B. HUPPERT, *Endliche Gruppen I*, Springer-Verlag, Berlin, Heidelberg, New York, 1967.

Manoscritto pervenuto in redazione l'8 dicembre 1984 ed in forma modificata il 26 marzo 1985.