

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

PAOLO ZANARDO

On $*$ -modules over valuation rings

Rendiconti del Seminario Matematico della Università di Padova,
tome 83 (1990), p. 193-199

http://www.numdam.org/item?id=RSMUP_1990__83__193_0

© Rendiconti del Seminario Matematico della Università di Padova, 1990, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

On *-Modules Over Valuation Rings.

PAOLO ZANARDO (*)

The problem of investigating *-modules over valuation rings was proposed to the author by C. Menini. We recall the definition of *-module, given by D'Este in [3]. Let R be a ring, ${}_R M$ a left R -module and ${}_R E$ an injective cogenerator of the category of all R -modules; let $S = \text{End}_R(M)$ and $H = \text{Hom}_R({}_R M, {}_R E)$ and denote by $\text{Gen}({}_R M)$ the category of all left R -modules generated by ${}_R M$ and by $\text{Cog}({}_S H)$ the category of all left S -modules cogenerated by H . In this situation, ${}_R M$ is said to be a *-module if there exists an equivalence of categories

$$\text{Gen}({}_R M) \xrightleftharpoons[G]{F} \text{Cog}({}_S H)$$

such that the functor F is naturally isomorphic to $\text{Hom}_R({}_R M, -)$ and the functor G is naturally isomorphic to $M_S \otimes -$ (we shall write $F \approx \text{Hom}_R({}_R M, -)$, $G \approx M_S \otimes -$).

The main motivation for the study of *-modules is the following result by Menini and Orsatti ([8], Theorem 3.1): let R, S be rings; if \mathfrak{S} is a full subcategory of $R\text{-Mod}$ closed under direct sums and factor modules, \mathfrak{D} is a full subcategory of $S\text{-Mod}$ containing ${}_S S$ and closed under submodules, and $\mathfrak{S} \xrightleftharpoons[G]{F} \mathfrak{D}$ is any equivalence with F and G additive functors, then there exists a module ${}_R M$ such that: $S = \text{End}_R({}_R M)$, $\mathfrak{S} = \text{Gen}({}_R M)$, $\mathfrak{D} = \text{Cog}({}_S H)$ (where ${}_S H$ is as above), $F \approx \text{Hom}_R({}_R M, -)$ and $G \approx M_S \otimes -$.

Recent results on *-modules have been obtained by D'Este [3], D'Este and Happel [4], Colpi [1], Colpi and Menini [2].

(*) Lavoro eseguito con il contributo del M.P.I.

Indirizzo dell'A.: Dipartimento di Matematica Pura e Applicata, Università dell'Aquila, 67100 L'Aquila, Italy.

In the present paper we characterize *finitely generated *-modules over a valuation ring R* . Using a theorem by Colpi ([1], Prop. 4.3) and some results in [9] (see also [5], Ch. IX), we prove that a finitely generated module X over a valuation ring R is a $*$ -module if and only if $X \cong (R/A)^n$, for suitable $n \geq 0$ and A ideal of R (Theorem 3). Note that a module of the form $(R/A)^n$ is a $*$ -module for any ring R , as a consequence of the above mentioned result by Colpi. Hence our Theorem 3 shows that the class of finitely generated $*$ -modules over a valuation ring is, in a certain sense, as small as possible.

Note that, at present, there are no examples of rings which admit $*$ -modules not finitely generated; Colpi and Menini in [2] proved that $*$ -modules over artinian rings or noetherian domains with Krull dimension one are necessarily finitely generated. The author feels that the same is true for $*$ -modules over valuation rings. Our final Remark 4 gives a contribution in this direction.

The author thanks R. Colpi, G. D'Este, C. Menini and L. Salce for helpful discussions and comments.

I. – In the sequel, R will always denote a *valuation ring*, i.e. a commutative ring, not necessarily a domain, whose ideals are linearly ordered by inclusion; the maximal ideal of R is denoted by P . For general terminology and results on modules over valuation rings we refer to the book by Fuchs and Salce [5]; the results we need on finitely generated modules can be found in [9] or in Ch. IX of [5].

In the proof of Theorem 2 we shall need the following facts (see [9] or [5], Ch. IX): let X be a finitely generated R -module; then there exists a submodule B of X such that:

- i) B is a direct sum of cyclic submodules;
- ii) B is pure in X ;
- iii) B is essential in X ;

such a B is said to be *basic* in X ; the basic submodules of X are all isomorphic. Moreover, given a basic submodule B of X , there exists a *minimal* set of generators $\mathbf{x} = \{x_1, \dots, x_k, x_{k+1}, \dots, x_n\}$ of X such that:

$$a) B = \langle x_1, \dots, x_k \rangle = \bigoplus_{i=1}^k \langle x_i \rangle;$$

b) if $A_j = \text{Ann}(x_j + \langle x_1, \dots, x_{j-1} \rangle)$ for all $j > k$, we have $A_{k+1} \leq A_{k+2} \leq \dots \leq A_n$;

c) for all $r \in A_{k+1}$ we have the relation

$$(1) \quad rx_{k+1} = r \sum_{i=1}^k a_i^r x_i, \quad \text{for suitable units } a_i^r \in R.$$

The construction of \mathbf{x} needs some explanation: we start with $B = \bigoplus_{i=1}^k \langle x_i \rangle$ basic in X and consider X/B ; if $\{x_{k+1} + B, \dots, x_n + B\}$ is a minimal set of generators of X/B , from the purity of B it follows that $\mathbf{x} = \{x_1, \dots, x_k, x_{k+1}, \dots, x_n\}$ is a minimal set of generators of X ; in view of Lemma 1.1 of [9], we can permute the indexes $k + 1, \dots, n$ to obtain property b). Since B is pure in X , certainly, for all $r \in A_{k+1}$, the relation (1) holds for suitable elements $a_i^r \in R$, not necessarily units. However, if $r \in \text{Ann } x_i$ for some $i \leq k$, obviously we can replace a_i^r with 1; moreover, if there exist $i \leq k$ and $s \in A_{k+1} \setminus \text{Ann } x_i$ such that $a_i^s \in P$, then for all $r \in A_{k+1} \setminus \text{Ann } x_i$ we have $a_i^r \in P$: in fact, if r divides s , from (1) we get $s(a_i^r - a_i^s)x_i = 0$, hence $a_i^s \in P$ implies $a_i^r \in P$; analogously, if s divides r , $r(a_i^r - a_i^s)x_i = 0$ implies $a_i^r \in P$. Let now $F = \{i \leq k: a_i^r \in P \text{ for all } r \in A_{k+1} \setminus \text{Ann } x_i\}$; if we replace x_{k+1} with $x'_{k+1} = x_{k+1} + \sum_{i \in F} x_i$, we obtain that

$$\text{Ann}(x'_{k+1} + B) = \text{Ann}(x_{k+1} + B) = A_{k+1},$$

$$\mathbf{x}' = \{x_1, \dots, x_k, x'_{k+1}, \dots, x_n\}$$

is a minimal set of generators of X , and (1) becomes

$$(1') \quad rx'_{k+1} = r \sum_{i=1}^k b_i^r x_i \quad \text{for all } r \in A_{k+1}$$

where $b_i^r = a_i^r$ if $i \notin F$ and $b_i^r = 1 + a_i^r$ if $i \in F$, so that b_i^r is a unit for all $i \leq k$ and for all $r \in A_{k+1}$. We conclude that there exists a minimal set of generators \mathbf{x} of X which satisfies properties a), b), c), as desired.

Let us now recall Colpi's result (Prop. 4.3 of [1]).

THEOREM 1 (Colpi). *Let R be a ring, ${}_R M$ a left R -module. Then ${}_R M$ is a *-module if and only if the following conditions are satisfied:*

- i) ${}_R M$ is self-small;
- ii) for each exact sequence

$$0 \rightarrow L \rightarrow N \rightarrow N/L \rightarrow 0$$

where N is an object of $\text{Gen}({}_R M)$, the sequence

$$0 \rightarrow \text{Hom}_R(M, L) \rightarrow \text{Hom}_R(M, N) \rightarrow \text{Hom}_R(M, N/L) \rightarrow 0$$

is exact if and only if $L \in \text{Gen}({}_R M)$. ///

We can now prove our main result.

THEOREM 2. *Let R be a valuation ring, let X be a finitely generated R -module and let $\pi: X \rightarrow X/PX$ be the canonical homomorphism. If the map $\varphi: \text{End } X \rightarrow \text{Hom}_R(X, X/XP)$, $\varphi: f \mapsto \pi \circ f$ is surjective, then $X \cong (R/A)^n$ for suitable $n \geq 0$ and A ideal of R .*

PROOF. In the following we assume $X \neq (R/A)^0 = \{0\}$, otherwise all is trivial. First of all, let us prove that X is a direct sum of cyclic submodules. Let B be a basic submodule of X ; it is enough to verify that $B = X$. By contradiction, suppose that $B < X$; let

$$\mathbf{x} = \{x_1, \dots, x_k, x_{k+1}, \dots, x_n\}$$

be a minimal set of generators of X which satisfies conditions a), b), c) above. Note that, since $B < X$, we have $k < n$, hence condition c) and the relation (1) are not trivially satisfied. For all $j \leq n$, let $\bar{x}_j = x_j + PX$; we have $X/PX = \bigoplus_{j=1}^n \langle \bar{x}_j \rangle$. Let us now consider the homomorphism $g: X \rightarrow X/PX$ defined extending by linearity the assignments

$$g: x_j \mapsto 0 \quad \text{if } j \neq k+1; \quad g: x_{k+1} \mapsto \bar{x}_{k+1}.$$

By hypothesis, there exists $f \in \text{End } X$ such that $g = \pi \circ f$. Hence, for $j \neq k+1$, we will have

$$(2) \quad f(x_j) = p \sum_{h=1}^n a_{hj} x_h, \quad \text{with } p \in P, \quad a_{hj} \in R$$

and

$$(3) \quad f(x_{k+1}) = x_{k+1} + q \sum_{h=1}^n b_h x_h, \quad \text{with } q \in P, \quad b_h \in R.$$

From (1), (2), (3), and the linearity of f , it follows, for all $r \in A_{k+1}$

$$(4) \quad r\left(x_{k+1} + q \sum_{h=1}^n b_h x_h\right) = rp \sum_{i=1}^k a_i^r \left(\sum_{h=1}^n a_{hi} x_h \right).$$

Since $A_{k+1} \leq A_t$ for all $t > k + 1$, and B is pure, we deduce that, for all $r \in A_{k+1}$

$$(5) \quad rq \sum_{h=1}^n b_h x_h \in rqB \quad \text{and} \quad rp \sum_{i=1}^k a_i^r \left(\sum_{h=1}^n a_{hi} x_h \right) \in rpB.$$

Let $\bar{p} \in P$ be a common divisor of p and q ; from (4) and (5) we get $rx_{k+1} \in r\bar{p}B$ for all $r \in A_{k+1}$, i.e.

$$(6) \quad rx_{k+1} = r\bar{p} \sum_{i=1}^k c_i^r x_i, \quad \text{with } c_i^r \in R.$$

From (1), (6), and the linear independence of x_1, \dots, x_k we obtain

$$(7) \quad r(a_i^r - \bar{p}c_i^r)x_i = 0 \quad \text{for } i = 1, \dots, k;$$

since a_i^r is a unit for all i and r , we have that $a_i^r - \bar{p}c_i^r$ is a unit, too, hence (7) implies $r \in \text{Ann } x_i$ for all $r \in A_{k+1}$. But this means that $rx_{k+1} \in B$ implies $rx_{k+1} = 0$, from which $\langle x_{k+1} \rangle \cap B = 0$, and B is not essential, against the definition of basic submodule. We conclude that, necessarily, $X = B$, as desired. It remains to prove that, if $A = \text{Ann } X$, then $X \cong (R/A)^n$. By contradiction, let us suppose that $X = \bigoplus_{i=1}^n \langle x_i \rangle$, where, for a suitable $j \leq n$, $\text{Ann } x_j > A$. Let us assume, without loss of generality, that $\text{Ann } x_1 = A$. Let $\eta: X \rightarrow X/PX$ be the homomorphism which extends by linearity the assignments

$$\eta: x_i \mapsto 0 \quad \text{if } i \neq j; \quad \eta: x_j \mapsto \bar{x}_1 = x_1 + PX.$$

If $\theta \in \text{End } X$ is such that $\eta = \pi \circ \theta$, then we have

$$(8) \quad \theta(x_j) = x_1 + p \sum_{i=1}^n a_i x_i, \quad \text{with } p \in P, \quad a_i \in R.$$

Choose now $r \in \text{Ann } x_j \setminus A$; from (8) we obtain

$$(9) \quad 0 = \theta(rx_j) = r(1 + pa_1)x_1 + rp \sum_{i=2}^n a_i x_i,$$

from which $r(1 + pa_1)x_1 = 0$, which is impossible, because $r \notin A = \text{Ann } x_i$. This concludes the proof. $///$

As an easy consequence of the preceding result we obtain the following

THEOREM 3. *Let R be a valuation ring. A finitely generated R -module X is a $*$ -module if and only if $X \cong (R/A)^n$ for suitable $n \geq 0$ and A ideal of R .*

PROOF. For any ring R , modules of the form $(R/A)^n$ are $*$ -modules as a consequence of Theorem 1, observing that $\text{Gen } ((R/A)^n) = R/A - \text{Mod}$, and $\text{Hom}_R((R/A)^n, -) \approx \text{Hom}_{R/A}((R/A)^n, -)$, if $n \geq 1$.

Conversely, let us note that $PX \in \text{Gen } (X)$, as it is easy to check. Therefore, if X is a finitely generated $*$ -module, then, by Theorem 1, X must satisfy the condition in the hypothesis of Theorem 2, hence X has the desired form. $///$

The problem of finding $*$ -modules which are not finitely generated remains open. We actually think that a $*$ -module over a valuation ring must be finitely generated; this opinion is mainly based on the following remark, derived from discussions with L. Salce.

REMARK 4. The simplest non finitely generated R -modules are the *uniserial* ones, i.e. those R -modules whose lattice of submodules is linearly ordered. Fuchs and Salce proved that, if U is a *divisible* uniserial module over a valuation domain R , whose elements have nonzero principal annihilators, then there is an equivalence of categories

$$\text{Gen } (U) \xrightleftharpoons[G]{F} \mathbb{C}$$

where \mathcal{C} is the class of complete torsion-free reduced R -modules, $F \approx \text{Hom}_R(U, -)$ and $G \approx U \otimes_R -$ (see [6]; this equivalence was inspired by Matlis equivalence in [7]; see also [5], p. 99). Moreover, U is small if and only if it is not countably generated. Nevertheless, for any choice of R we notice that U is not a *-module. This is clear if U is countably generated (see Theorem 1). If U is not countably generated, then also Q , the field of fractions of R , is not countably generated as an R -module; in this case we get that \mathcal{C} is not closed for submodules, hence \mathcal{C} cannot be cogenerated by any module. It is worth giving a check of this last fact: let us suppose, by contradiction, that \mathcal{C} is closed for submodules, for a convenient R , with Q not countably generated as an R -module; with these assumptions, R must be complete, and each free R -module F is complete, too, in view of Cor. 2.2 of [6]. Let us consider a short exact sequence

$$0 \rightarrow K \rightarrow F \rightarrow Q \rightarrow 0$$

with F free; then $F \in \mathcal{C}$ implies $K \in \mathcal{C}$, hence K is closed in F and $Q \cong F/K$ must be Hausdorff in the natural topology, i.e. $\{0\} = \bigcap_{r \in R^*} rQ = Q$, a contradiction.

REFERENCES

- [1] R. COLPI, *Some remarks on equivalence between categories of modules*, to appear on Comm. in Algebra.
- [2] R. COLPI - C. MENINI, *On the structure of *-modules*, to appear.
- [3] G. D'ESTE, *Some remarks on representable equivalences*, to appear in the Proceedings of the Banach Center.
- [4] G. D'ESTE - D. HAPPEL, *Representable equivalences are represented by tilting modules*, to appear on Rend. Sem. Mat. Univ. Padova.
- [5] L. FUCHS - L. SALCE, *Modules over valuation domains*, Lecture Notes in Pure Appl. Math., n. 97, Marcel Dekker, New York, 1985.
- [6] L. FUCHS - L. SALCE, *Equivalent categories of divisible modules*, Forum Math., 1 (1989).
- [7] E. MATLIS, *Cotorsion modules*, Mem. Amer. Math. Soc., 49 (1964).
- [8] C. MENINI - A. ORSATTI, *Representable equivalences between categories of modules and applications*, to appear on Rend. Sem. Mat. Univ. Padova.
- [9] L. SALCE - P. ZANARDO, *Finitely generated modules over valuation rings*, Comm. Algebra, 12 (15), (1984), pp. 1795-1812.

Manoscritto pervenuto in redazione il 7 luglio 1989.